

Mental Health Nursing Interventions in Schizophrenia Care: Effectiveness of Nurse-Led Strategies in Managing Schizophrenia Disorder

Chandani Thakur^{1,*}, Bire Rohidas Suresh², Marinder Pal Kaur³

Abstract

Schizophrenia is a chronic and severe mental disorder characterized by disturbances in thought, perception, emotion, and behavior that can significantly impair an individual's functioning and quality of life. Mental health nurses play a pivotal role in the comprehensive management of schizophrenia through patient-centered and evidence-based interventions. This article critically examines the effectiveness of nurse-led strategies in the management of schizophrenia, with particular emphasis on pharmacological support, psychosocial interventions, psychoeducation, therapeutic communication, relapse prevention, and community-based care. Evidence published over the past ten years is synthesized to assess the impact of these interventions on clinical and psychosocial outcomes, including symptom reduction, medication adherence, relapse prevention, functional recovery, and quality of life. The reviewed evidence indicates that nurse-led interventions can contribute significantly to improved treatment adherence, symptom management, patient engagement, self-management, and overall psychosocial functioning. Therapeutic communication and psychoeducation further support patients and caregivers in understanding the illness, managing symptoms, recognizing early warning signs, and preventing relapse. Community-based nursing interventions also promote continuity of care and facilitate reintegration into family and community settings. The findings highlight the essential contribution of mental health nurses to multidisciplinary schizophrenia care and emphasize the need for strengthened nursing education, structured clinical training, continuous professional development, and supportive mental health policies. Integrating evidence-based nurse-led interventions into routine schizophrenia care may improve long-term outcomes and enhance the quality of life of individuals living with schizophrenia.

Keywords: Schizophrenia, mental health nursing, nurse-led interventions, psychosocial care, psychiatric nursing, patient outcomes, community mental health, relapse prevention

*Author for Correspondence

Chandani Thakur

E-mail: healtheducation236@gmail.com

¹Assistant Professor, Department of Nursing, Kailsah Institute of Health and Medical Sciences, Panchkula, Haryana, India.

²Associate Professor, Department of Nursing, Pravara Rural Education Society's Sindhutai Vikhe Patil College of Nursing, Chincholi, Nashik, Maharashtra, India.

³Professor Cum Vice Principal, Department of Nursing, College of Nursing, Adesh Institute of Medical Sciences, Sri Muktsar Sahib, Punjab, India.

Received Date: April 20, 2026

Accepted Date: June 20, 2026

Published Date: June 30, 2026

Citation: Chandani Thakur, Bire Rohidas Suresh, Marinder Pal Kaur. Mental Health Nursing Interventions in Schizophrenia Care: Effectiveness of Nurse-Led Strategies in Managing Schizophrenia Disorder. Journal of Nursing Science & Practice. 2026; 16(2): 1–18p.

INTRODUCTION

Schizophrenia is a complex and chronic psychiatric disorder affecting approximately one percent of the global population and contributes significantly to disability, stigma, and socioeconomic burden across diverse healthcare systems worldwide, with long-term implications for individuals, families, and communities [1]. It is characterized by a heterogeneous constellation of symptoms broadly categorized into positive symptoms, such as hallucinations and delusions; negative symptoms, including apathy, anhedonia, and social withdrawal; and cognitive impairments that affect attention, working memory, and

executive functioning, all of which collectively interfere with daily functioning and quality of life. The multifactorial etiology of schizophrenia involves complex interactions between genetic predisposition, neurodevelopmental abnormalities, environmental stressors, and neurochemical dysregulation, particularly in dopaminergic and glutamatergic pathways. These underlying mechanisms contribute to the variability in symptom presentation and disease progression, necessitating individualized and multidisciplinary approaches to care. The chronic and relapsing nature of schizophrenia requires long-term management strategies that extend beyond pharmacological treatment to incorporate psychosocial interventions, rehabilitation, and community-based support systems aimed at promoting recovery and social reintegration. Mental health nurses play a pivotal role in this continuum of care, acting as primary caregivers who deliver holistic, patient-centered interventions that address the biological, psychological, and social dimensions of the disorder [2]. Their responsibilities include comprehensive assessment of mental status, monitoring of treatment response, administration of antipsychotic medications, identification of side effects, and implementation of evidence-based psychosocial therapies. Figure 1 presents an overview of schizophrenia symptom domains and functional impact to illustrate the multidimensional nature of schizophrenia and its impact on functional outcomes, highlighting the need for integrated care approaches. Over the past decade, the role of mental health nurses has evolved significantly from traditional task-oriented functions to advanced clinical roles encompassing therapeutic communication, psychoeducation, case management, and coordination of multidisciplinary services [3]. This evolution reflects the growing recognition of nursing contributions to mental healthcare delivery and the increasing demand for specialized skills in managing complex psychiatric conditions. Nurse-led interventions have demonstrated considerable effectiveness in improving treatment adherence, reducing relapse rates, and enhancing recovery-oriented outcomes through structured approaches, such as cognitive behavioral therapy, motivational interviewing, and family-focused interventions. Furthermore, nurses play a critical role in psychoeducation by empowering patients and caregivers with knowledge about illness management, medication adherence, early warning signs of relapse, and coping strategies, thereby fostering self-management and resilience [4]. Therapeutic relationships established by nurses are fundamental to patient engagement, as trust and rapport facilitate open communication, reduce stigma, and promote adherence to treatment plans. Community-based mental health nursing interventions, including home visits, outreach programs, and rehabilitation services, have been shown to improve accessibility to care and reduce hospitalization rates among individuals with schizophrenia. Advances in digital health and telepsychiatry have further expanded the scope of nurse-led interventions, enabling remote monitoring, virtual counseling, and continuity of care for underserved populations. Despite these advancements, challenges such as stigma, limited resources, workforce shortages, and variability in training continue to impact the effectiveness of mental health nursing practice, highlighting the need for ongoing professional development and policy support in this field. Evidence from recent studies underscores the importance of integrating recovery-oriented approaches into nursing practice, emphasizing patient autonomy, shared decision-making, and social inclusion as key components of effective care for schizophrenia. Moreover, nurse-led early intervention strategies have shown promise in improving long-term outcomes by reducing the duration of untreated psychosis and facilitating timely access to care. The incorporation of culturally sensitive care and individualized treatment plans further enhances the effectiveness of nurse-led interventions by addressing diverse patient needs and preferences [5].

Pathophysiology and Clinical Presentation of Schizophrenia

Schizophrenia is widely recognized as a complex neurodevelopmental disorder characterized by the dysregulation of multiple neurotransmitter systems, particularly dopamine and glutamate pathways, which play a central role in the manifestation of psychotic symptoms and cognitive dysfunction. Contemporary research highlights that abnormalities in dopaminergic transmission, especially within the mesolimbic and mesocortical pathways, are associated with positive symptoms such as hallucinations and delusions, whereas glutamatergic dysfunction contributes to cognitive impairments and negative symptoms observed in schizophrenia. In addition to neurotransmitter imbalances, structural and functional brain abnormalities, including reduced gray matter volume in the prefrontal

cortex, hippocampus, and temporal lobes, have been consistently reported in individuals with schizophrenia, further supporting the neurobiological basis of schizophrenia. Genetic predisposition plays a significant role in the etiology of schizophrenia, with heritability estimates suggesting that genetic factors account for a substantial proportion of the risk, although no single gene is solely responsible for the disease development. Environmental factors such as prenatal infections, obstetric complications, childhood trauma, urban living, and substance use interact with genetic vulnerability to trigger schizophrenia onset, emphasizing the disorder's multifactorial nature. Figure 2 depicts key neurotransmitter pathways, brain regions, and their interactions, providing a visual understanding of the underlying mechanisms that contribute to symptom development and progression. Clinically, schizophrenia presents with a heterogeneous array of symptoms that vary in severity, duration, and response to treatment, making diagnosis and management particularly challenging for healthcare professionals. Positive symptoms, including hallucinations, delusions, and disorganized thinking, are often the most recognizable features of the disorder and are primarily linked to the hyperactivity of dopaminergic pathways. Negative symptoms, such as emotional blunting, reduced speech, lack of motivation, and social withdrawal, are associated with significant functional impairment and are often more resistant to treatment than positive symptoms. Cognitive deficits represent another critical dimension of schizophrenia, affecting domains such as attention, memory, executive functioning, and information processing speed, which collectively impair an individual's ability to perform daily activities and maintain social and occupational roles. These cognitive impairments are considered core features of schizophrenia and are strongly associated with poor long-term outcomes, highlighting the need for targeted cognitive rehabilitation strategies to address them. The variability in symptom presentation and progression reflects the heterogeneity of schizophrenia, with some individuals experiencing episodic symptoms and others exhibiting a chronic course with persistent symptoms. Early identification and intervention are crucial for improving prognosis, as prolonged duration of untreated psychosis is associated with worse clinical and functional outcomes. Mental health nurses play a vital role in recognizing these diverse symptom dimensions and tailoring interventions accordingly, ensuring that care is individualized and responsive to patients' needs. Understanding the neurobiological and clinical complexity of schizophrenia enables nurses to implement evidence-based interventions that address symptom management and functional recovery. Furthermore, advances in neuroimaging and biomarker research continue to enhance our understanding of schizophrenia pathophysiology, offering the potential for more precise diagnostic and therapeutic approaches in the future. Despite these advancements, challenges remain in translating neurobiological knowledge into clinical practice, particularly in resource-limited settings, where access to specialized care may be restricted. Therefore, healthcare professionals, especially nurses, must comprehensively understand the neurodevelopmental, neurochemical, and clinical aspects of schizophrenia to provide effective patient-centered care and improve overall treatment outcomes [6-9].

Role of Mental Health Nurses in Schizophrenia Care

Mental health nurses play a central and indispensable role in the comprehensive management of schizophrenia by actively participating in the assessment, planning, implementation, and evaluation of individualized care plans that address the complex and multidimensional needs of individuals affected by schizophrenia. Their role begins with a thorough and continuous assessment of patients' mental status, behavioral patterns, symptom severity, and psychosocial functioning, which forms the foundation for evidence-based care planning and intervention. Nurses utilize standardized assessment tools and clinical judgment to identify changes in symptoms, such as hallucinations, delusions, mood disturbances, and cognitive impairments, thereby enabling the early detection of relapse or deterioration. In addition to assessment, mental health nurses are responsible for developing and implementing care plans that integrate pharmacological and non-pharmacological interventions tailored to individual patient needs and recovery goals. Medication management is a critical aspect of nursing practice, involving the administration of antipsychotic drugs, monitoring for therapeutic effectiveness, and identifying and managing potential side effects, such as extrapyramidal symptoms, metabolic syndrome, and sedation. Nurses also play a key role in promoting medication adherence by educating

patients about the importance of consistent treatment and addressing barriers, such as stigma, lack of insight, and fear of side effects. Figure 3 illustrates the diverse responsibilities of mental health nurses across acute care, outpatient services, and community-based settings, emphasizing their role in ensuring continuity and coordination of care [10–13].

Psychoeducation is another essential nursing intervention that empowers patients and their families with knowledge about schizophrenia, treatment options, coping strategies, and relapse prevention, thereby enhancing engagement and self-management. Through structured educational sessions, nurses help patients develop insights into their condition and improve their ability to recognize early warning signs of relapse. Psychosocial interventions delivered by nurses, including cognitive behavioral therapy, social skills training, and supportive counseling, are effective in improving social functioning, reducing symptom severity, and enhancing overall quality of life. Nurses also provide therapeutic communication and emotional support, which are fundamental to building trust and fostering a therapeutic alliance with patients, ultimately improving treatment outcomes. Their role as patient advocates involves safeguarding patient rights, promoting dignity, and ensuring access to appropriate healthcare services, particularly for vulnerable populations who may face stigma and discrimination. Coordination of care is another critical function, as nurses collaborate with psychiatrists, psychologists, social workers, and other healthcare professionals to deliver integrated multidisciplinary care. This coordination extends to community-based services, where nurses facilitate discharge planning, follow-up care, and rehabilitation programs aimed at promoting recovery and social reintegration of patients. Table 1 summarizes key areas of nursing practice, including pharmacological support, psychosocial interventions, psychoeducation, relapse prevention, and community care, highlighting the breadth of nursing responsibilities in managing schizophrenia. The close and continuous interaction between nurses and patients allows for the early identification of subtle changes in behavior or symptoms, enabling timely intervention and prevention of relapse. Furthermore, nurses play a vital role in addressing patients' psychosocial needs by supporting their participation in daily activities, improving social relationships, and enhancing coping skills. Advances in mental health nursing have expanded the scope of practice to include the use of digital technologies and telehealth services, allowing nurses to provide remote monitoring, education, and support to patients in diverse settings. Despite these




Figure 1. Overview of schizophrenia symptom domains and functional impact, including positive, negative, and cognitive symptoms.




Figure 2. Neurobiological basis of schizophrenia showing brain regions and neurotransmitter imbalances.

Positive symptoms	Negative symptoms	Cognitive symptoms
Deviant symptoms. These symptoms are present but should be absent. They indicate that the patient has lost touch with reality. Delusions disorganized behavior & speech	Deficit symptoms. These symptoms reflect the absence of normal characteristics Apathy Anhedonia Blunted affect Poverty of speech	Reflect the patient's abnormal thinking, poor decision-making skill, poor problem-solving skills and ability to communicate and his strange behavior. Thought disorders Bizarre behavior.

Figure 3. Nursing roles in schizophrenia management, including assessment, treatment, and care coordination.

advancements, challenges such as high workload, limited resources, and insufficient training continue to impact the effectiveness of nursing interventions, underscoring the need for ongoing professional development and institutional support [14-17].

Pharmacological Support and Medication Management

Antipsychotic medications remain the cornerstone of schizophrenia treatment because of their effectiveness in reducing psychotic symptoms and stabilizing patients. Mental health nurses play a crucial role in ensuring adherence, monitoring therapeutic responses, and managing potential side effects associated with these medications. These medications primarily act on dopaminergic pathways, helping to control positive symptoms such as hallucinations and delusions, whereas newer atypical antipsychotics also target serotonergic systems to address negative symptoms and cognitive deficits. Nurses are responsible for administering medications accurately and consistently, ensuring that prescribed regimens are followed, and observing patients for clinical improvement or deterioration. A key component of nurse-led medication management is patient education, where nurses provide clear and understandable information about the purpose of medications, expected benefits, duration of treatment, and potential side effects, thereby enhancing patients' understanding and engagement. Education also includes addressing misconceptions about antipsychotics, reducing stigma, and encouraging patients to actively participate in their treatment plans. Figure 4 shows the antipsychotic treatment pathway and nursing monitoring responsibilities, which demonstrates the systematic process of medication management, including prescription, administration, monitoring, and follow-up care, highlighting the integral role of nurses at each stage. Nurses are also responsible for identifying and managing adverse effects associated with antipsychotic medications, such as extrapyramidal symptoms,

weight gain, metabolic syndrome, sedation, and cardiovascular complications, which can significantly impact patient adherence and quality of life. Early detection and prompt management of these side effects are essential to prevent complications and ensure continued adherence to treatment. Motivational strategies employed by nurses, such as motivational interviewing and supportive counseling, are effective in addressing patient concerns, enhancing insight, and promoting adherence to medication. Non-adherence to antipsychotic medication is a major challenge in schizophrenia management and is associated with an increased risk of relapse, hospitalization, and poor long-term outcomes, making nursing interventions critical in mitigating these risks. Nurses also collaborate with multidisciplinary teams to adjust treatment plans based on patient responses and tolerability, ensuring that pharmacological therapy is optimized for each individual. Long-acting injectable antipsychotics have emerged as an important option for improving adherence. Nurses play a key role in administering these medications, monitoring their effects, and educating patients about their benefits. In addition to their clinical responsibilities, nurses provide emotional support and build therapeutic relationships that foster trust and open communication, enabling patients to express concerns about medications and treatments. This therapeutic alliance is essential for improving adherence and overall treatment outcomes. Nurses also monitor for signs of relapse, which may indicate poor adherence or inadequate treatment response, and initiate timely interventions to prevent deterioration. Advances in digital health technologies have further enhanced nurse-led medication management by enabling remote monitoring, reminders, and telehealth consultations, thereby improving adherence and continuity of care. Despite these advancements, challenges such as patient resistance, side effect burden, and limited access to healthcare resources continue to impact medication adherence, highlighting the need for ongoing nursing education and support. Evidence consistently demonstrates that effective nurse-led medication management significantly reduces medication non-adherence, leading to lower relapse rates, decreased hospitalization, and improved quality of life for individuals with schizophrenia [18, 19].

Psychosocial Interventions Led by Nurses

Psychosocial interventions are fundamental components of comprehensive schizophrenia care and are frequently delivered by trained mental health nurses as part of a holistic, recovery-oriented approach aimed at improving functional outcomes and the overall quality of life. These interventions complement pharmacological treatment by addressing the psychological, social, and behavioral dimensions of the disorder, which are often not fully managed by medication alone. Cognitive behavioral therapy is one of the most widely used psychosocial interventions in schizophrenia care, focusing on helping patients identify and modify maladaptive thoughts, beliefs, and behaviors associated with psychotic experiences. Through structured sessions, nurses assist patients in developing coping strategies to manage hallucinations and delusions, reduce distress, and improve their insight into their condition. Social skills training is another key intervention that aims to enhance interpersonal communication, social interaction, and problem-solving abilities, which are often impaired in individuals with schizophrenia. Mental health nurses facilitate these training programs by providing role-playing exercises, feedback, and reinforcement to help patients build their confidence and improve their social functioning. Figure 5 highlights the integration of these nurse-led strategies, illustrating how cognitive, behavioral, and social interventions work together to promote recovery and functional independence of patients with schizophrenia. Rehabilitation programs are also essential components of psychosocial care, focusing on vocational training, supported employment, and daily living skills to help patients reintegrate into society and achieve meaningful roles. Nurses play a central role in coordinating these programs and providing ongoing support to ensure sustained engagement and progress in the program. Family interventions, often delivered by nurses, are crucial in reducing caregiver burden, improving family communication, and enhancing the support system for patients, which, in turn, reduces relapse rates. Psychoeducation is frequently integrated into psychosocial interventions, enabling patients and families to understand the nature of schizophrenia, treatment options, strategies for managing symptoms, and preventing relapse. Evidence indicates that patients who receive structured psychoeducation demonstrate better treatment adherence and improved long-term outcomes. Nurses also utilize motivational interviewing techniques to enhance patient engagement and encourage participation in psychosocial programs, particularly among individuals with low motivation or insight into their

condition. In addition to traditional face-to-face interventions, digital and technology-based psychosocial interventions have gained prominence in recent years, with nurses facilitating online therapy sessions, mobile health applications, and telepsychiatry services to increase the accessibility and continuity of care. These innovations are particularly beneficial for individuals in remote or underserved areas where access to mental health services may be limited. Psychosocial interventions have been shown to significantly reduce symptom severity, particularly negative and cognitive symptoms, which are often resistant to pharmacological treatment. Furthermore, these interventions improve social functioning, enhance quality of life, and promote recovery by enabling patients to develop the skills necessary for independent living. Nurses play a vital role in tailoring psychosocial interventions to individual patient needs, considering factors such as cultural background, illness severity, and personal goals. Continuous monitoring and evaluation of patient progress allow nurses to adjust their interventions and ensure optimal outcomes. Despite their effectiveness, challenges such as limited resources, time constraints, and variability in training can impact the implementation of psychosocial interventions, highlighting the need for ongoing professional development and institutional support [20-24].

Psychoeducation and Family Involvement

Psychoeducation is a critical and evidence-based nursing intervention in the management of schizophrenia that focuses on empowering patients and their families with comprehensive knowledge of the disorder, its symptoms, treatment modalities, and coping strategies, thereby enhancing treatment adherence and overall outcomes. Mental health nurses play a central role in delivering structured psychoeducation sessions tailored to the cognitive and emotional needs of patients, ensuring that information is presented in a clear, accessible, and culturally sensitive manner. These sessions typically include education about the nature of schizophrenia, its causes and risk factors, early warning signs of relapse, medication management, and strategies for coping with stress and daily challenges. By improving patients' understanding of their condition, psychoeducation enhances insight, which is often impaired in patients with schizophrenia and is a major barrier to treatment adherence. Figure 6 illustrates the systematic process of delivering education, including assessment of learning needs, provision of information, reinforcement, and evaluation of outcomes, highlighting the structured approach used by nurses. Family involvement is an essential component of psychoeducation, as caregivers play a crucial role in supporting patients' recovery and ensuring continuity of care within the community. Nurses engage family members in educational sessions to improve their understanding of schizophrenia, reduce misconceptions, and enhance their ability to provide effective support to patients. Family psychoeducation has been shown to reduce relapse rates, decrease caregiver burden, and improve overall family functioning, making it a vital aspect of comprehensive care for patients with schizophrenia. In addition, psychoeducation helps families develop skills to manage challenging behaviors, communicate effectively, and respond appropriately to crises, thereby creating a supportive environment for their recovery. Nurses also address the stigma associated with schizophrenia during psychoeducation sessions, helping patients and families overcome negative beliefs and attitudes that may hinder treatment engagement. The use of interactive teaching methods, such as group discussions, role-playing, and visual aids, enhances learning and retention of information, making psychoeducation more effective in improving mental health. Advances in digital technology have further expanded the scope of psychoeducation, with nurses utilizing online platforms, mobile applications, and telehealth services to deliver education remotely and improve its accessibility. These digital interventions are particularly beneficial for individuals in rural or underserved areas where access to mental health services may be limited. Psychoeducation also plays a key role in promoting self-management, as patients learn to independently monitor their symptoms, adhere to medication regimens, and implement coping strategies. Nurses continuously evaluate the effectiveness of psychoeducation by assessing patient understanding, adherence, and clinical outcomes, and modifying interventions as needed to ensure optimal results. Evidence suggests that patients who receive structured psychoeducation demonstrate improved medication adherence, reduced relapse rates, and better quality of life than those who do not receive such interventions. Furthermore, psychoeducation enhances therapeutic relationships between nurses, patients, and families, fostering trust, collaboration, and shared decision-

making in the care process of patients with schizophrenia. Despite its proven benefits, challenges such as limited time, high workload, and lack of standardized programs can affect the delivery of psychoeducation, highlighting the need for institutional support and training [25].

Table 1. Core domains of mental health nursing interventions in schizophrenia care.

Domain	Key interventions	Nursing activities	Expected outcomes
Pharmacological support	Antipsychotic medication management	Medication administration. Monitoring side effects. Educating patients about adherence	Symptom control. Reduced relapse. Improved adherence
Psychosocial interventions	CBT. Social skills training. Rehabilitation	Conduct therapy sessions. Facilitate group activities. Support behavioral change	Improved social functioning. Reduced symptoms
Psychoeducation	Patient and family education	Provide structured education. Teach coping strategies. Increase illness awareness	Better insight. Increased adherence. Reduced relapse
Therapeutic communication	Nurse–patient relationship building	Active listening. Empathy. Non-judgmental interaction	Reduced anxiety. Improved engagement
Relapse prevention	Early identification and intervention	Monitor warning signs. Crisis management. Follow-up care	Decreased relapse rates. Early intervention
Community care	Home-based and rehabilitation services	Home visits. Coordination of care. Social reintegration support	Reduced hospitalization. Improved quality of life
Care coordination	Multidisciplinary collaboration	Liaise with healthcare team. Discharge planning	Continuity of care. Holistic management
Health promotion	Lifestyle and wellness support	Encourage physical activity. Address substance use. Promote self-care	Improved overall health. Better long-term outcomes




Figure 4. Antipsychotic treatment pathway and nursing monitoring responsibilities.

Therapeutic Communication and Relationship Building

Therapeutic communication is a fundamental component of mental health nursing practice and serves as the cornerstone for building effective nurse–patient relationships in the management of schizophrenia. It enables nurses to engage patients meaningfully and provide individualized emotional and psychological support. Establishing trust and rapport is particularly important in schizophrenia care, as patients often experience paranoia, delusions, and impaired reality testing, which may hinder their willingness to interact with healthcare professionals. Mental health nurses utilize structured communication techniques, such as active listening, which involves giving full attention to the patient, acknowledging their experiences, and responding appropriately to verbal and nonverbal cues, thereby fostering a sense of being understood and valued. Empathy is another essential element of therapeutic communication, allowing nurses to understand patients' feelings and perspectives without judgment, which is crucial for reducing distress and promoting emotional stability. Non-judgmental interaction is particularly important when dealing with patients experiencing hallucinations or delusions, as dismissing or challenging these experiences directly can damage trust and hinder therapeutic engagement. Instead, nurses are trained to acknowledge the patient's experience while gently guiding them toward reality-based thinking, thereby maintaining a supportive and therapeutic environment for patients with dementia. Effective therapeutic communication also involves the use of open-ended questions, clarification, reflection, and summarization techniques to encourage patients to express their thoughts and feelings freely. These techniques help nurses gather comprehensive information about a patient's condition, identify concerns, and tailor interventions accordingly. In schizophrenia care, communication must be adapted to the patient's cognitive abilities and level of functioning, as cognitive deficits may affect their comprehension, attention, and response. Nurses often use simple language, repetition, and visual aids to enhance their understanding and ensure effective communication. Therapeutic communication plays a vital role in reducing patient anxiety, which is commonly experienced due to psychotic symptoms, hospitalization, or social isolation. By providing reassurance, emotional support, and a safe space for expression, nurses help patients feel more secure and less overwhelmed by their symptoms. Effective communication also improves patient engagement in treatment by fostering collaboration and shared decision-making, which are the key principles of recovery-oriented care. Nurses who establish strong therapeutic relationships are better able to motivate patients to adhere to medication regimens, participate in psychosocial interventions, and engage in rehabilitation. Furthermore, therapeutic communication enhances nurses' ability to detect early warning signs of relapse, such as changes in mood, behavior, or thought patterns, enabling timely intervention and prevention of deterioration. Cultural competence is another important aspect of therapeutic communication, as nurses must be sensitive to cultural beliefs, values, and communication styles that may influence patients' perceptions of illness and treatment. Additionally, the use of digital communication tools, including telehealth and mobile applications, has expanded the scope of therapeutic communication, allowing nurses to maintain contact with patients and provide support beyond traditional clinical settings. Despite its importance, barriers such as high workload, time constraints, and lack of training can impact the effectiveness of therapeutic communication, highlighting the need for ongoing education and skill development in this area [26].

Relapse Prevention and Early Intervention

Relapse is a significant and recurrent challenge in the long-term management of schizophrenia and is commonly associated with factors such as poor medication adherence, inadequate social support, substance use, and high levels of psychosocial stress, all of which contribute to symptom exacerbation and increased hospitalization rates. Mental health nurses play a pivotal role in relapse prevention by continuously monitoring patients for early warning signs, including subtle changes in mood, behavior, sleep patterns, and thought processes that may indicate an impending relapse. Early identification of these warning signs enables timely intervention, which is critical for preventing full-blown psychotic episodes and reducing the severity of relapse. Nurses employ structured assessment tools and clinical observations to track symptom progression and detect deviations from baseline functioning, thereby facilitating proactive care. Figure 7 demonstrates the systematic approach used by nurses, including early detection, intervention, and follow-up, highlighting the effectiveness of nurse-led strategies in

reducing the risk of relapse. Medication adherence is a cornerstone of relapse prevention, and nurses play a key role in ensuring that patients follow the prescribed treatment regimens through education, counseling, and regular follow-ups. By addressing barriers such as side effects, lack of insight, and stigma, nurses help improve adherence and reduce the likelihood of relapse in patients. In addition to pharmacological management, nurses implement psychosocial interventions, such as cognitive behavioral therapy and stress management techniques, to enhance coping skills and resilience. Crisis intervention is another critical aspect of relapse prevention, as nurses must be prepared to respond promptly to acute deterioration in mental status by providing immediate support, de-escalation techniques, and coordination of emergency care when necessary. Family involvement is also essential for relapse prevention, as caregivers can help monitor symptoms, provide emotional support, and ensure continuity of care in the community. Nurses educate families about early warning signs and effective coping strategies, enabling them to respond appropriately and seek timely professional assistance. Community-based nursing interventions, including home visits and follow-up care, further enhance relapse prevention by maintaining regular contact with patients and addressing potential risk factors in their environments. Advances in digital health technologies have introduced new opportunities for relapse prevention, with nurses utilizing mobile applications and telehealth platforms to remotely monitor symptoms and provide ongoing support. These digital tools allow real-time tracking of patient progress and early detection of warning signs, thereby improving intervention outcomes. Psychoeducation also plays a crucial role in relapse prevention, as patients who understand their illness and treatment are more likely to adhere to medication and recognize the early signs of relapse. Nurses reinforce this knowledge through continuous education and support, promoting self-management and independence. Despite the effectiveness of nurse-led interventions, challenges such as limited resources, high workload, and patient-related factors can impact relapse prevention efforts, highlighting the need for comprehensive and coordinated care systems to support these initiatives. Evidence consistently shows that structured relapse prevention programs led by nurses significantly reduce the frequency and severity of relapses, improve patient outcomes, and enhance the quality of life [27].




Figure 5. Psychosocial intervention model in schizophrenia care led by nurses.

Understanding Schizophrenia

- Understanding the illness
- Dispelling myths & stigma
- Reducing feelings of guilt & blame

Family Involvement

- Role of the family in recovery
- Effective communication

Early Warning Signs & Relapse Prevention

- Identifying early warning signs
- Developing a relapse prevention plan
- Knowing when to seek help

Early Warning Signs and Relapse Prevention Symptom Management

- Information
- Support
- Empowerment.
- Hope

Family Involvement

- Role of the family in recovery
- Effective communication
- Setting healthy boundaries

Stress and Coping Strategies

- Relaxation techniques
- Problem-solving skills
- Healthy lifestyle habits

Medication Management

- Purpose & benefits of medications
- Addressing side effects
- Adherence & safety

Symptom Management

- Identifying symptoms
- Recognizing triggers
- Teaching coping skills

Figure 6. Psychoeducation framework for patients and families in schizophrenia care.

Community-Based Mental Health Nursing Interventions

Community care is a fundamental component of the long-term management of schizophrenia, as it facilitates continuity of care beyond hospital settings and supports individuals in achieving functional recovery and social reintegration within their natural environments. Mental health nurses play a central role in delivering community-based interventions, including home-based care, follow-up services, and rehabilitation programs tailored to the individual needs of patients. Through regular home visits, nurses assess patients' mental status, monitor medication adherence, and identify environmental or psychosocial stressors that may contribute to relapse, thereby enabling early intervention and prevention. Community-based care models emphasize a recovery-oriented approach that prioritizes patient autonomy, independence, and participation in meaningful activities, which are essential for improving the quality of life. Figure 8 illustrates the community mental health nursing model, which

depicts the continuum of care from acute hospital treatment to community-based support, highlighting the role of nurses in ensuring seamless transitions and ongoing management [28].

Rehabilitation services provided by nurses include vocational training, social skills development, and assistance with daily living activities, all of which are critical for reintegration into society and reducing the risk of long-term disability. Nurses also collaborate with multidisciplinary teams, including psychiatrists, social workers, and occupational therapists, to deliver coordinated and comprehensive care that addresses the diverse needs of individuals with schizophrenia. Community-based interventions have been shown to improve access to mental health services, particularly in underserved and rural areas where hospital-based care may be limited. These interventions also reduce hospitalization rates by providing ongoing support and early management of symptoms, thereby preventing relapses and crisis situations. Table 2 highlights key determinants such as training and education, workload, patient engagement, and availability of support systems, all of which significantly impact the quality and outcomes of community mental health care. Adequate training and education enhance nurses' competence in delivering evidence-based interventions, whereas high workload and limited resources may hinder effective care delivery. Patient Engagement is another critical factor, as active participation in treatment and rehabilitation programs is associated with better outcomes and sustained recovery. Support systems, including family involvement and community resources, play vital roles in facilitating recovery and reducing the burden on healthcare services. Nurses also provide psychoeducation to patients and families in community settings, empowering them with the knowledge and skills to manage the illness effectively and recognize early warning signs of relapse. Advances in digital health technologies have further enhanced community care by enabling remote monitoring, telepsychiatry, and mobile health interventions, which improve accessibility and continuity of care. These innovations allow nurses to maintain regular contact with patients and provide timely support, even in geographically remote areas of the country. Despite the benefits of community-based care, challenges such as stigma, lack of infrastructure, and workforce shortages continue to affect service delivery and patient outcomes in the Philippines. Addressing these challenges requires policy support, increased funding, and investment in mental health services to strengthen the community care systems. Evidence consistently demonstrates that nurse-led community interventions significantly improve clinical outcomes, enhance social functioning, and promote recovery in individuals with schizophrenia [29].




Figure 7. Relapse prevention model highlighting early detection and nursing interventions.




Figure 8. Community mental health nursing model showing continuity of care.

Impact of Nurse-Led Interventions on Patient Outcomes

Research over the past decade has consistently demonstrated that nurse-led interventions play a significant role in improving both clinical and psychosocial outcomes in individuals with schizophrenia by integrating pharmacological management with holistic, patient-centered care approaches that address the complex needs of this population. These interventions contribute to a marked reduction in symptom severity, particularly in managing positive symptoms such as hallucinations and delusions through continuous monitoring, timely intervention, and adherence support provided by mental health nurses. In addition, nurse-led care has been shown to enhance medication adherence, which is a critical determinant of treatment success and relapse prevention, as nurses actively educate patients, address concerns related to side effects, and provide motivational support to maintain long-term compliance. Improved adherence subsequently leads to fewer hospitalizations and better symptom stabilization, thereby reducing the overall burden on healthcare systems. Beyond clinical outcomes, nurse-led interventions significantly improve psychosocial functioning by helping patients develop social skills, coping mechanisms, and strategies for managing daily activities, which are essential for independent living. Nurses facilitate structured psychosocial interventions, such as cognitive behavioral therapy and rehabilitation programs, that promote recovery and reintegration into society. Table 3 summarizes key improvements observed in patients, including reduced symptom severity, decreased relapse rates, enhanced quality of life, and improved social functioning, highlighting the effectiveness of nursing interventions in comprehensive care. Quality of life is another important outcome influenced by nurse-led care, as nurses provide emotional support, therapeutic communication, and psychoeducation that empower patients to manage their condition effectively and improve their overall health. Family involvement, facilitated by nurses, further enhances outcomes by providing a supportive environment that encourages adherence and reduces the risk of relapse. Nurses also play a crucial role in the early identification of relapse warning signs, enabling prompt intervention and prevention of symptom exacerbation, which contributes to sustained recovery. Community-based nurse-led interventions,

including home visits and follow-up care, improve access to services and ensure continuity of care, which is particularly important for patients with chronic conditions, such as schizophrenia. Advances in digital health have further strengthened the impact of nurse-led interventions by enabling remote monitoring, telepsychiatry, and mobile-based support systems that enhance patient engagement and adherence to treatment. These innovations allow nurses to provide continuous care and timely interventions, even in remote or underserved regions. Despite the positive outcomes associated with nurse-led interventions, challenges such as high workload, limited resources, and variability in training can affect the quality and consistency of care delivery, highlighting the need for ongoing professional development and institutional support for nurses. Evidence also suggests that nurses who receive specialized training in mental health are more effective in delivering interventions that improve patient outcomes, emphasizing the importance of education and competency-based training programs in mental health nursing. Furthermore, nurse-led interventions contribute to reducing stigma and improving patient engagement by fostering therapeutic relationships and promoting a recovery-oriented approach to care.

Table 2. Factors influencing effectiveness of nurse-led interventions

Factor	Description	Impact on care	Outcome
Training and Education	Level of knowledge and clinical competence of nurses	Determines the ability to deliver evidence-based interventions	Improved patient outcomes and quality of care
Workload	Number of patients and responsibilities assigned	High workload reduces time for individualized care	Decreased care quality and increased burnout
Patient engagement	Degree of patient involvement in treatment	Active participation enhances adherence and recovery	Better clinical and psychosocial outcomes
Organizational support	Availability of resources, supervision, and policies	Strong support improves efficiency and morale	Enhanced service delivery and continuity of care
Communication skills	Ability to build therapeutic relationships	Effective communication improves trust and engagement	Increased adherence and patient satisfaction
Multidisciplinary collaboration	Coordination with other healthcare professionals	Ensures holistic and integrated care	Comprehensive treatment outcomes
Resource availability	Access to medications, infrastructure, and tools	Limited resources restrict intervention delivery	Poorer outcomes and increased relapse risk
Cultural competence	Sensitivity to patient beliefs and backgrounds	Improves acceptance of care and communication	Enhanced patient cooperation and trust

Table 3. Outcomes of nurse-led interventions in schizophrenia care.

Outcome domain	Description	Nursing contribution	Impact
Symptom reduction	Decrease in positive and negative symptoms	Medication management. Psychosocial support	Improved clinical stability
Medication adherence	Consistent use of prescribed treatment	Patient education. Monitoring. Motivation	Reduced relapse and hospitalization
Relapse rate	Frequency of symptom recurrence	Early detection. Continuous follow-up	Lower relapse incidence
Quality of life	Overall well-being and life satisfaction	Holistic care. Emotional support	Enhanced daily functioning
Social functioning	Ability to interact and maintain relationships	Social skills training. Rehabilitation	Better community integration
Hospitalization	Frequency and duration of hospital stays	Preventive care. Crisis management	Reduced admissions
Insight and Awareness	Understanding of illness and treatment	Psychoeducation	Improved self-management
Functional independence	Ability to perform daily activities	Rehabilitation and community support	Increased autonomy and recovery

Challenges in Mental Health Nursing Practice

Despite their critical role in the management of schizophrenia and other mental health conditions, mental health nurses face numerous challenges that significantly impact their ability to deliver high-quality, patient-centered care, including high workload, stigma, limited resources, and inadequate training, all of which can hinder effective intervention and patient outcomes. High workload is one of the most pressing issues, as nurses often manage large caseloads and complex patient needs, leading to burnout, fatigue, and a reduced capacity to provide individualized care and therapeutic engagement. This excessive workload can compromise patient safety, reduce the time available for meaningful interactions, and limit the implementation of comprehensive psychosocial interventions. In addition, mental health nurses frequently encounter stigma, both from society and within healthcare systems, which can affect their professional identity, job satisfaction, and quality of care provided to patients with schizophrenia. Stigma may also discourage individuals from seeking mental health services, thereby increasing the burden on nurses who must manage more severe and untreated cases of mental illness. Limited resources, including inadequate staffing, insufficient infrastructure, and lack of access to essential medications and therapeutic tools, further exacerbate the challenges faced by mental health nurses, particularly in low- and middle-income countries (LMICs). Resource constraints can lead to delays in treatment, reduced availability of community-based services, and increased reliance on institutional care, which may not always be conducive to recovery. Inadequate training and a lack of continuous professional development opportunities also pose significant barriers, as nurses may not be fully equipped with the latest evidence-based practices and skills required to manage complex psychiatric conditions effectively. The rapidly evolving nature of mental health care necessitates ongoing education and training to ensure that nurses remain competent and confident in their roles. Furthermore, the emotional demands of working with patients with chronic and severe mental illnesses can lead to psychological stress, compassion fatigue, and burnout among nurses, affecting their well-being and job performance. Lack of organizational support, including limited supervision, inadequate recognition, and poor working conditions, can further contribute to job dissatisfaction and high turnover rates among mental health nurses. Communication barriers, cultural differences, and language issues may also impact the effectiveness of care, particularly in diverse populations, where understanding patient perspectives is crucial for successful intervention. Additionally, the integration of mental health services into primary care remains limited in many regions, placing additional pressure on specialized mental health nurses to meet the growing demand. While technological advancements are beneficial, they also present challenges, as nurses must adapt to new systems such as telepsychiatry and digital health tools, which require additional training and resources. Addressing these challenges requires a multifaceted approach that includes policy reforms, increased funding, workforce development, and the implementation of supportive organizational structures that promote nurses' well-being and professional growth. Strengthening mental health education and training programs, improving staffing levels, and fostering a positive work environment are essential strategies for enhancing the effectiveness of mental health nursing practices. Moreover, reducing stigma through public awareness campaigns and promoting the value of mental health nursing can improve both patient outcomes and the professional status of nurses [30].

Implications for Nursing Education and Practice

Strengthening nursing education through evidence-based curricula and simulation-based training is essential for preparing mental health nurses to effectively manage complex conditions, such as schizophrenia, ensuring that they possess the knowledge, skills, and competencies required for high-quality patient care. Modern nursing education must integrate current research findings, clinical guidelines, and best practices into curricula to bridge the gap between theory and practice and promote evidence-based decision-making in clinical settings. Simulation-based training has emerged as a powerful educational tool that allows nursing students and practicing professionals to engage in realistic clinical scenarios, enhancing their critical thinking, clinical judgment, and confidence in managing psychiatric emergencies and providing routine care. Through high-fidelity simulation, nurses can practice communication skills, medication administration, and crisis intervention in a controlled environment without risk to patients, thereby improving their readiness for real-world situations and

enhancing patient safety. Continuous professional development programs are equally important, as they ensure that nurses remain updated on evolving treatment modalities, technological advancements, and evidence-based practices in mental health care. These programs may include workshops, seminars, online courses, and certification programs that focus on enhancing clinical competence and promoting lifelong learning for nurses. Regular training updates also help nurses adapt to new healthcare technologies, such as telepsychiatry, digital monitoring tools, and electronic health records, which are increasingly being integrated into mental health services. In addition, competency-based assessments should be incorporated into professional development programs to evaluate nurses' knowledge and skills, identify learning gaps, and ensure adherence to the established standards of care. Interprofessional education is another key component of strengthening nursing education, as it fosters collaboration and communication among healthcare professionals, leading to more coordinated and effective care. By working alongside psychiatrists, psychologists, and social workers during training, nurses develop a better understanding of multidisciplinary approaches to managing schizophrenia. Furthermore, incorporating recovery-oriented principles into nursing education encourages a patient-centered approach that emphasizes autonomy, empowerment, and social inclusion, which are critical for improving long-term outcomes for patients with SUD. Cultural competence training is also essential, as it enables nurses to provide care that is respectful of and responsive to diverse patient backgrounds, beliefs, and values, thereby enhancing therapeutic relationships and treatment effectiveness. Mentorship and clinical supervision play vital roles in supporting nurses' professional development by providing guidance, feedback, and opportunities for reflective practice that contribute to continuous improvement. Despite the recognized importance of education and training, barriers such as limited funding, time constraints, and lack of institutional support can hinder the implementation of comprehensive development programs, highlighting the need for policy-level intervention. Investment in nursing education and workforce development is essential to address the growing demand for mental health services and ensure that nurses are adequately prepared to meet the challenges of modern healthcare. Evidence suggests that nurses who receive ongoing education and training demonstrate higher competence, improved clinical performance, and better patient outcomes, reinforcing the importance of continuous professional development. Ultimately, strengthening nursing education through evidence-based curricula, simulation training, and lifelong learning initiatives is crucial for enhancing the quality of mental health care, promoting adherence to best practices, and ensuring optimal outcomes for individuals with schizophrenia.

CONCLUSION

Mental health nursing interventions play a crucial role in the effective management of schizophrenia by addressing the complex and multifaceted needs of individuals with this chronic psychiatric disorder. Nurses are at the center of care delivery, implementing comprehensive and patient-centered strategies that integrate pharmacological support, psychosocial interventions, psychoeducation, and community-based care. Through medication management, nurses ensure adherence, monitor therapeutic outcomes, and manage side effects, thereby reducing the risk of relapse and hospitalization. Psychosocial interventions delivered by nurses, including cognitive and behavioral strategies, social skills training, and rehabilitation programs, help improve functional abilities and promote recovery. Psychoeducation further empowers patients and their families by enhancing their understanding of the illness, encouraging active participation in care, and strengthening coping mechanisms. Nurses' community care supports long-term management by facilitating continuity of care, promoting social reintegration, and improving access to mental health services. These nurse-led interventions collectively contribute to improved clinical outcomes, enhanced quality of life, and greater independence among individuals with schizophrenia. However, the effectiveness of these interventions depends on the competence, training, and support available to nursing staff. Strengthening nursing education through evidence-based curricula, enhancing clinical training through simulation and practical exposure, and ensuring adequate organizational support are essential for optimizing care delivery. By investing in the development of mental health nurses and creating supportive healthcare environments, the overall quality and effectiveness of schizophrenia management can be significantly improved, leading to better patient outcomes and sustainable mental healthcare systems.

REFERENCES

1. Addington J, Heinssen RK, Robinson DG, Schooler NR, Marcy P, Brunette MF, et al. Duration of untreated psychosis in community treatment settings in the United States. *Psychiatr Serv*. 2015;66(7):753–756. doi:10.1176/appi.ps.201400124.
2. Alvarez-Jimenez M, Alcazar-Corcoles MA, Gonzalez-Blanch C, Bendall S, McGorry PD, Gleeson JF. Online social media: new data, new horizons in psychosis treatment. *Schizophr Res*. 2015;166(1-3):345–346. doi:10.1016/j.schres.2015.05.006.
3. Barkhof E, Meijer CJ, de Sonnevile LM, Linszen DH, de Haan L. Interventions to improve adherence to antipsychotic medication in patients with schizophrenia: a review of the past decade. *Eur Psychiatry*. 2012;27(1):9–18. doi:10.1016/j.eurpsy.2011.02.005.
4. Bäuml J, Froböse T, Kraemer S, Rentrop M, Pitschel-Walz G. Psychoeducation: a basic psychotherapeutic intervention for patients with schizophrenia and their families. *Schizophr Bull*. 2006;32 Suppl 1:S1–S9. doi:10.1093/schbul/sbl017.
5. Castillo RJ, Guo KL. A framework for cultural competence in health care organizations. *Health Care Manag*. 2011;30:205–214. doi:10.1097/HCM.0b013e318225dfe6.
6. Bighelli I, Salanti G, Huhn M, Schneider-Thoma J, Krause M, Reitmeir C, et al. Psychological interventions to reduce positive symptoms in schizophrenia: systematic review and network meta-analysis. *World Psychiatry*. 2018;17(3):316–329. doi:10.1002/wps.20577.
7. Gleeson JF, Rawlings D, Jackson HJ, McGorry PD. Early warning signs of relapse following a first episode of psychosis. *Schizophr Res*. 2005;80:107–111. doi:10.1016/j.schres.2005.07.019.
8. Bond GR, Drake RE, Becker DR. An update on Individual Placement and Support. *World Psychiatry*. 2020;19(3):390–391. doi:10.1002/wps.20784.
9. Bosanac P, Patton GC, Castle DJ. Early intervention in psychotic disorders: faith before facts? *Psychol Med*. 2010;40(3):353–358. doi:10.1017/S0033291709990341.
10. Brissos S, Veguilla MR, Taylor D, Balanzá-Martinez V. The role of long-acting injectable antipsychotics in schizophrenia: a critical appraisal. *Ther Adv Psychopharmacol*. 2014;4(5):198–219. doi:10.1177/2045125314540297.
11. Cant RP, Cooper SJ. Simulation in the Internet age: the place of web-based simulation in nursing education: an integrative review. *Nurse Educ Today*. 2014;34(12):1435–1442. doi:10.1016/j.nedt.2014.08.001.
12. Chien WT, Bressington D, Yip A, Karatzias T. An international multi-site, randomized controlled trial of a mindfulness-based psychoeducation group programme for people with schizophrenia. *Psychol Med*. 2017;47:2081–2096. doi:10.1017/S0033291717000526.
13. Claxton M, Onwumere J, Fornells-Ambrojo M. Do family interventions improve outcomes in early psychosis? A systematic review and meta-analysis. *Front Psychol*. 2017;8:371. doi:10.3389/fpsyg.2017.00371.
14. Correll CU, Galling B, Pawar A, Krivko A, Bonetto C, Ruggeri M, et al. Comparison of early intervention services vs treatment as usual for early-phase psychosis: a systematic review, meta-analysis, and meta-regression. *JAMA Psychiatry*. 2018;75(6):555–565. doi:10.1001/jamapsychiatry.2018.0623.
15. Corrigan PW, Druss BG, Perlick DA. The impact of mental illness stigma on seeking and participating in mental health care. *Psychol Sci Public Interest*. 2014;15(2):37–70. doi:10.1177/1529100614531398.
16. De Hert M, Detraux J, van Winkel R, Yu W, Correll CU. Metabolic and cardiovascular adverse effects associated with antipsychotic drugs. *Nat Rev Endocrinol*. 2011;8(2):114–126. doi:10.1038/nrendo.2011.156.
17. Shields MC, Trinh NT, Borba CPC, Chen R, Reddy AK, Singer SJ. Former inpatient psychiatric patients' past experiences with traditional frontline staff and their thoughts on the benefits of peers as part of frontline staff. *J Psychosoc Nurs Ment Health Serv*. 2022;60:15–22. doi:10.3928/02793695-20210916-01.
18. Dixon LB, Goldman HH, Bennett ME, Wang Y, McNamara KA, Mendon SJ, et al. Implementing coordinated specialty care for early psychosis: the RAISE connection program. *Psychiatr Serv*. 2015;66:691–698. doi:10.1176/appi.ps.201400281.

19. Drake RE, Bond GR, Essock SM. Implementing evidence-based practices for people with schizophrenia. *Schizophr Bull.* 2009;35:704–713. doi:10.1093/schbul/sbp041.
20. Färdig R, Lewander T, Melin L, Folke F, Fredriksson A. A randomized controlled trial of the illness management and recovery program for persons with schizophrenia. *Psychiatr Serv.* 2011;62(6):606–612. doi:10.1176/ps.62.6.pss6206_0606.
21. Firth J, Siddiqi N, Koyanagi A, Siskind D, Rosenbaum S, Galletly C, et al. The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. *Lancet Psychiatry.* 2019;6:675–712. doi:10.1016/S2215-0366(19)30132-4.
22. Foronda CL, Fernandez-Burgos M, Nadeau C, Kelley CN, Henry MN. Virtual simulation in nursing education: a systematic review spanning 1996 to 2018. *Simul Healthc.* 2020;15(1):46–54. doi:10.1097/SIH.0000000000000411.
23. Freeman D, Taylor KM, Molodynski A, Waite F. Treatable clinical intervention targets for patients with schizophrenia. *Schizophr Res.* 2019;211:44–50. doi:10.1016/j.schres.2019.07.016.
24. Galderisi S, Mucci A, Buchanan RW, Arango C. Negative symptoms of schizophrenia: new developments and unanswered research questions. *Lancet Psychiatry.* 2018;5:664–677. doi:10.1016/S2215-0366(18)30050-6.
25. Galletly C, Castle D, Dark F, Humberstone V, Jablensky A, Killackey E, et al. Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the management of schizophrenia and related disorders. *Aust N Z J Psychiatry.* 2016;50(5):410–472. doi:10.1177/0004867416641195.
26. Happell B, Platania-Phung C, Scott D. Proposed nurse-led initiatives in improving physical health of people with serious mental illness: a survey of nurses in mental health. *J Clin Nurs.* 2014;23(7-8):1018–1029. doi:10.1111/jocn.12371.
27. McCutcheon RA, Keefe RSE, McGuire PK. Cognitive impairment in schizophrenia: aetiology, pathophysiology, and treatment. *Mol Psychiatry.* 2023;28:1902–1918. doi:10.1038/s41380-023-01949-9.
28. Howes OD, Murray RM. Schizophrenia: an integrated sociodevelopmental-cognitive model. *Lancet.* 2014;383(9929):1677–1687. doi:10.1016/S0140-6736(13)62036-X.
29. Kane JM, Robinson DG, Schooler NR, Mueser KT, Penn DL, Rosenheck RA, et al. Comprehensive versus usual community care for first-episode psychosis: 2-year outcomes from the NIMH RAISE early treatment program. *Am J Psychiatry.* 2016;173(4):362–372. doi:10.1176/appi.ajp.2015.15050632.
30. Cognitive-behavioural therapy for the symptoms of schizophrenia: systematic review and meta-analysis with examination of potential bias. Database of Abstracts of Reviews of Effects (DARE): quality-assessed reviews. York (UK): Centre for Reviews and Dissemination, University of York; 2014. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK179939/>.