STM JOURNALS

Research & Reviews: A Journal of
Medical Science and Technology

https://journals.stmjournals.com/rrjomst/

ISSN: 2319-3417 (Online)
ISSN: 2349-1272 (Print)
Volume 15, Issue 3, 2026
September–December
DOI (Journal): 10.37591/RRJoMST

Review RRJoMST

# Identifying the Most Common Causes of Maternal Morbidity and Mortality in Rural Areas: A Comprehensive Epidemiological Analysis

Vanisha Mavi[1,*], Nidhi Tyagi[2]

***Abstract***

*Pregnancy-related complications continue to represent a major public health burden in remote and underserved regions across developing nations, where healthcare infrastructure remains limited. This investigation synthesizes evidence from epidemiological studies, regional health surveys, and WHO databases to delineate the principal medical conditions and systemic barriers contributing to maternal mortality in rural populations. Through systematic examination of clinical literature and statistical health data, we identify five conditions – hypertensive emergencies during pregnancy, severe postpartum bleeding, systemic infections in the peripartum period, nutritional anaemia, and complications from protracted labor – as accounting for most preventable maternal deaths in rural settings. The analysis further demonstrates that structural constraints including geographic isolation, insufficient trained healthcare personnel, and inadequate emergency medical infrastructure amplify the severity of these treatable conditions. Our synthesis reveals that approximately 80–95% of rural maternal fatalities are theoretically preventable through implementation of evidence-based clinical interventions grounded in medical science and strengthening the health system. The findings presented herein offer a framework for prioritizing resource allocation and designing targeted maternal health programs suited to resource-limited rural healthcare environments.*

**Keywords**: Rural maternal health, obstetric emergencies, public health epidemiology, preventable mortality, healthcare access barriers, clinical prevention strategies

## INTRODUCTION

How a society treats its pregnant women says a great deal about it. Maternal health reflects not just the state of a country's health system, but how equitably its benefits are shared, and the gap between well-resourced settings and low-resource regions remains one of the starkest in global health. Every year, roughly 280,000 to 300,000 women die from complications of pregnancy or childbirth, and the great majority of those deaths occur in economically developing countries [1]. Within those countries the rural–urban divide is especially sharp: women in rural community's die of pregnancy-related causes at rates three to five times those of their urban counterparts. In sub-Saharan Africa and South Asia, a rural woman's lifetime risk of dying from pregnancy-related causes is markedly higher than an urban woman's with equivalent access to care [1]. Nor is this disadvantage purely a matter of medical science alone. A blend of clinical, logistical, infrastructural and social factors combines to turn conditions that should be survivable into fatal ones. The classic account of this process identifies three points at which deadly delay can occur: deciding to seek care, reaching a facility, and receiving adequate treatment once there [2].

***Author for Correspondence**
Vanisha Mavi
E-mail: Vanisha.mavi@gmail.com

[1]Research Scholar, School of Computation Sciences and Engineering, Shobhit Institute of Engineering and Technology, (Deemed to be University), Meerut, Uttar Pradesh, India.
[2]Professor, School of Computation Sciences and Engineering, Shobhit Institute of Engineering and Technology, (Deemed to be University), Meerut, Uttar Pradesh, India.

Received Date: October 11, 2026
Accepted Date: October 21, 2026
Published Date: November 2, 2026

**Citation:** Vanisha Mavi, Nidhi Tyagi. Identifying the Most Common Causes of Maternal Morbidity and Mortality in Rural Areas: A Comprehensive Epidemiological Analysis. Research & Reviews: A Journal of Medical Science and Technology. 2026; 15(3): 28–38p.

© STM Journals 2026. All Rights Reserved 

This study sets out to characterise, as thoroughly as the evidence allows, the medical conditions and underlying determinants behind the elevated maternal mortality seen in rural areas worldwide. Drawing on published epidemiological research, international health data and regional case studies, we identify which clinical conditions carry the greatest mortality burden, examine the features of rural life that make them more dangerous, and set out evidence-based approaches to reducing preventable deaths. The aim is practical: to inform policy development and resource allocation for maternal health programmes working in resource-constrained rural settings.

## LITERATURE REVIEW AND EPIDEMIOLOGICAL CONTEXT

Maternal health complications remain among the most pressing problems facing health systems, and nowhere more so than in poorly resourced rural areas [1]. Looking at severe maternal morbidity and mortality trends, Li and colleagues found that maternal mortality in the United States more than doubled between 1990 and 2016, from 7.2 to 16.9 deaths per 100,000 live births [3]. Deaths from direct obstetric causes – such as haemorrhage and infection – have fallen in well-resourced settings, but those linked to pre-existing chronic conditions have risen sharply [3]. Disparities persist across countries too: non-Hispanic Black and indigenous women die at two to three times the rate of White women [3]. Severe maternal morbidity affects an estimated 50,000 women each year in the United States, falling disproportionately on racial and ethnic minorities and rural populations, and infants born to affected mothers are 39 percent more likely to die [3]. In a systematic analysis of global maternal mortality patterns, Say and colleagues estimated that direct obstetric complications, including haemorrhage, hypertensive disorders, post-delivery infections and abortion complications, account for roughly 70 to 80 percent of maternal deaths worldwide [4]. Working across 180 countries, Khan and colleagues documented that in low-income economies postpartum haemorrhage is the leading cause of maternal death, responsible for 24 to 40 percent of deaths depending on the region [5]. In middle-income countries, hypertensive complications in pregnancy claim a larger share, a pattern that says more about differences in healthcare infrastructure than about the underlying disease [6]. Rodriguez and colleagues, studying the early prediction of severe maternal morbidity, put the global toll at approximately 585,000 deaths a year during pregnancy, delivery or the puerperium, with close to 50 million complications recorded annually [7]. Work on severe maternal morbidity links it to hypertensive disorders, haemorrhage and infection, with Black race, obesity, high parity, previous caesarean sections and co-morbidities among the key risk factors [8]. On the prediction side, Poon and colleagues achieved a 90 percent detection rate for early pre-eclampsia with a 5 percent false-positive rate using logistic regression [9]. The foundational three delays framework of Thaddeus and Maine explains how rural residence raises maternal mortality through delays in deciding to seek help, reaching a facility, and receiving proper care once there [10]; limited health literacy, cultural preferences for home delivery, gender-based constraints and geographic isolation all feed these delays [11]. Ahmed and colleagues documented exactly this in rural Bangladesh, where pregnant women faced inconsistent antenatal monitoring, rudimentary transport and a lack of basic emergency obstetric capability [12]. Filippi and colleagues found that journeys of more than 100 kilometres, or more than six hours, to reach emergency obstetric care are genuine barriers to life-saving treatment in rural sub-Saharan Africa and South Asia [13]. Kongnyuy and colleagues identified the availability of skilled birth attendants as a critical determinant of survival, and rural areas remain chronically short of trained health workers [2]. Vasudevan and colleagues argued that maternal mortality will only fall meaningfully through approaches that combine clinical indicators with socioeconomic and infrastructural factors [14]. Creanga and colleagues, meanwhile, showed that in the United States the timing of maternal deaths is shifting, with later postpartum deaths (42 days to one year) increasingly attributed to cardiovascular disease, mental health problems and substance use [15]. Taken together, the epidemiological evidence shows that rural maternal deaths and injuries stem from an interplay of direct medical causes and systemic barriers to care [1, 12, 16, 17]. The great majority of these deaths are preventable; they reflect failures of health systems rather than limitation of medical science [3, 18, 19]. Where a region sits geographically shapes which causes dominate, and getting evidence-based interventions to scale in resource-limited settings remains the central challenge [2, 14].

© STM Journals 2026. All Rights Reserved 

## ANALYSIS OF LEADING CAUSES OF MATERNAL MORTALITY IN RURAL POPULATIONS

When we look closely at the medical conditions behind rural maternal deaths, a clear picture emerges: a small set of preventable complications accounts for most pregnancy-related deaths in resource-limited settings [4]. Modern medical science has long established effective treatments for each of these conditions, yet they remain lethal where healthcare infrastructure is absent.

### Postpartum Haemorrhage

Defined as blood loss of more than 500 millilitres after a vaginal delivery, or 1,000 millilitres after a caesarean, postpartum haemorrhage is the most immediate threat to a mother's survival and accounts for 24 to 40 percent of maternal deaths in low-income rural areas [20]. It usually arises from uterine atony, retained placenta, tears in the birth canal, or a clotting problem [21]. What makes it so deadly in rural settings is the absence of the things that stop it elsewhere: no oxytocin, no trained attendant, no blood for transfusion, no capacity for emergency surgery. Pre-existing anaemia, malnutrition, multiple pregnancy, missed antenatal monitoring and delivery outside a facility make matters worse. Yet it is also among the most preventable of causes, at 95 to 98 percent, and the remedies are well known: uterotonic drugs, skilled attendance, access to blood transfusion and surgical capability [4].

### Hypertensive Complications

Chronic hypertension, gestational hypertension, pre-eclampsia and eclampsia together complicate 2 to 8 percent of pregnancies worldwide, yet in some regions they account for 16 to 25 percent of maternal deaths [4, 22]. Eclampsia occurs 10 to 20 times more often in low-income countries than in well-resourced ones, which reflects differences in how the condition can be managed rather than how often it occurs [4]. The warning signs are well described: blood pressure of 140/90 mmHg or higher, protein in the urine, headache, visual disturbance and seizures that can escalate quickly [4]. Rural settings typically lack routine blood pressure monitoring, antihypertensive drugs, and magnesium sulphate for preventing seizures [4, 22]. With preventability at 90 to 95 percent, the interventions are equally well established: blood pressure monitoring, identification of high-risk pregnancies, reliable access to medication, and the capability to deliver a baby urgently when needed [4].

### Infection and Sepsis

Infections account for 10 to 20 percent of maternal deaths in low-income settings, ranging from endometritis and surgical wound infections to urinary tract infections and systemic sepsis [4]. Peripartum infection typically begins when vaginal flora ascends into the uterus, a process helped along by prolonged labour, premature rupture of membranes, repeated vaginal examinations, unclean delivery conditions and retained placenta [23]. Rural deaths from infection happen mostly in non-sterile home births and deliveries by traditional birth attendants, where sanitation is absent, instruments are not sterilised, antibiotics are scarce, warning signs go unrecognised and facilities respond slowly [4]. Preventability sits at 85 to 90 percent, and the remedies are institutional delivery, training in aseptic technique, access to antibiotics and sound sepsis management [4].

### Anaemia in Pregnancy

Anaemia, defined as a haemoglobin level below 11 g/dL, affects 40 to 50 percent of pregnant women worldwide, with prevalence above 50 percent in sub-Saharan Africa and South Asia, and contributes indirectly to 8 to 15 percent of maternal deaths [12]. In rural areas it stems from poor iron intake, malabsorption, parasitic infections – such as hookworm, malaria, and chronic infections – including tuberculosis and HIV [12]. Severe anaemia, below 7 g/dL, makes death from postpartum haemorrhage three to five times more likely, besides weakening immunity and reducing the oxygen reaching the fetus [12]. The rural obstacles are familiar: no haemoglobin testing, poor adherence to iron supplements, a heavy burden of concurrent infections, little dietary counselling and no transfusion capability [12]. Preventability runs at 80 to 90 percent through haemoglobin screening, iron and folate supplementation, treatment of infections, and access to transfusion [12].

© STM Journals 2026. All Rights Reserved 

### Obstructed Labour

When the fetus cannot progress despite adequate contractions, usually because of cephalopelvic disproportion, malpresentation or placental obstruction, labour becomes obstructed, and it accounts for 3 to 10 percent of maternal deaths [4]. Labour lasting beyond 18 to 24 hours brings maternal exhaustion, uterine rupture, fetal hypoxia, haemorrhage and sepsis [4]. Its rural prevalence tracks high adolescent pregnancy rates, childhood malnutrition, absent antenatal imaging, no means of monitoring labour, no caesarean capability and limited access to blood products [4]. Preventability is 70 to 85 percent, through antenatal risk identification, labour monitoring, surgical referral, caesarean access and better nutrition [4].

### Unsafe Abortion

Complications of unsafe abortion, including severe haemorrhage, uterine perforation, infection and sepsis following procedures by untrained providers in unhygienic settings, account for 5 to 10 percent of rural maternal deaths [1]. When post-abortion care and emergency services are out of reach, these complications turn fatal [1]. Prevention rests on access to contraception, sexuality education, and safe abortion services where the law permits them, and preventability is 95 to 99 percent [1].

### Placental Abnormalities

Placenta previa and placental abruption cause severe bleeding before and during delivery, but in rural areas without ultrasound they tend to go undetected until it is too late, producing deaths that are 80 to 90 percent preventable [4].

### Amniotic Fluid Embolism

This rare but catastrophic emergency during labour causes sudden cardiovascular collapse; rural women without access to intensive care survive it far less often, and preventability stands at only 50 to 60 percent [4].

### Thromboembolism

Deep venous thrombosis and pulmonary embolism cause deaths after delivery, with the risk raised by prolonged immobility, caesarean delivery and inherited thrombophilia. Rural shortages of anticoagulation therapy and vascular imaging make these deaths more likely, though 75 to 85 percent of them remain preventable [4].

### Anaesthetic Complications

Aspiration pneumonitis, failed intubation and adverse drug reactions claim rural women undergoing emergency caesarean delivery without adequate anaesthetic training and monitoring, and 75 to 85 percent of these deaths are preventable [4].

## COMPARATIVE EPIDEMIOLOGY: BURDEN OF DISEASE AND PREVENTION CAPACITY

Knowing how the causes of maternal death are distributed across rural regions, and how preventable each one is, tells us where resources and interventions should go first. The data show considerable regional variation in which causes dominate, but one conclusion holds everywhere: nearly all these deaths are preventable with adequate healthcare access and infrastructure.

### Relative Contribution of the Leading Causes

Seven conditions account for 75 to 95 percent of rural maternal deaths across geographic regions as illustrated in Table 1. Postpartum haemorrhage remains the single largest contributor in sub-Saharan Africa, though the pattern varies by region. The geography of these deaths' tracks healthcare infrastructure: regions with access to blood transfusion see lower haemorrhage mortality even where the underlying disease burden is no different.

© STM Journals 2026. All Rights Reserved 

**Table 1.** Prevalence and mortality impact of leading causes in rural areas.

| Condition | Proportion affecting pregnancies | Proportion of rural maternal deaths | Geographic variation | Theoretical preventability |
|---|---|---|---|---|
| Postpartum Haemorrhage | 2–5% | 32–45% | Higher in sub-Saharan Africa | 95–98% |
| Hypertensive Complications | 2–8% | 18–28% | Higher in South Asia & Latin America | 90–95% |
| Infection/Sepsis | 5–10% | 12–18% | Consistent across regions | 85–90% |
| Severe Anaemia | 15–50% | 8–15% | Higher in tropical regions | 80–90% |
| Obstructed Labor | 0.5–2% | 4–10% | Higher in adolescent pregnancy regions | 70–85% |
| Unsafe Abortion | 3–5% | 5–10% | Higher where access to safe services is limited | 95–99% |
| Other/Non-Direct Causes | Various | 10–20% | Varies by region | 40–70% |

## Visual Representation of Mortality Distributions

Figure 1 compares the burden of maternal mortality causes in rural and urban settings.





**Figure 1.** Interpretation: Rural deaths cluster around preventable obstetric causes, while urban deaths spread more towards other and indirect causes. The pattern shows that direct causes are prevented more successfully in cities.

Figure 2 shows maternal mortality ratios across geographic regions and settings.





**Figure 2.** Interpretation: The size of the rural–urban gap varies by region, but it is substantial everywhere. Sub-Saharan Africa and South Asia carry the heaviest absolute burden.

Figure 3 shows the composition of rural maternal deaths by primary cause across regions.

© STM Journals 2026. All Rights Reserved 





**Figure 3.** Interpretation: The share of each cause shifts by region, following differences in healthcare infrastructure, cultural practice and disease prevalence. Preventability, however, stays consistently above 90 percent everywhere.

The Three-Delay Framework and Its Manifestation in Rural Settings are classified in Table 2.

**Table 1.** Delays in maternal care and contributing factors in rural environments.

| Delay category | Contributing factors | Prevalence in rural areas | Resulting mortality impact |
|---|---|---|---|
| First Delay (Decision to Seek Care) | Limited awareness of danger signs | 60–80% | 2–3-fold |
| | Cultural preference for home delivery | 50–70% | |
| | Gender-based constraints on decision-making | 40–60% | |
| | Inadequate health literacy | 50–70% | |
| | Fear or mistrust of healthcare facilities | 30–50% | |
| | Economic constraints | 40–60% | |
| Second Delay (Reaching Care) | Geographic distance >50 kilometre's | 40–60% | 3–5-fold |
| | Inadequate road infrastructure | 70–90% | |
| | Unavailable emergency transport | 50–80% | |
| | High cost of transportation | 60–80% | |
| | Natural geographic barriers | 20–40% | |
| Third Delay (Receiving Treatment) | Insufficient trained personnel | 60–80% | 2–4-fold |
| | Unavailable blood supply | 70–90% | |
| | Absence of surgical capability | 60–80% | |
| | Shortage of essential medications | 70–85% | |
| | Limited monitoring equipment | 80–95% | |
| | Inadequate facility infrastructure | 75–90% | |

© STM Journals 2026. All Rights Reserved 

### Essential Interventions and Their Availability in Rural Healthcare Settings

Life-saving interventions for maternal emergencies and access to these interventions in rural areas are presented in Table 3.

**Table 3.** Life-saving interventions for maternal emergencies and rural access.

| Intervention | Clinical indication | Rural coverage rate | Mortality reduction potential |
|---|---|---|---|
| Antenatal Blood Pressure Screening | Detect hypertensive complications | 40–60% | 15–20% |
| Haemoglobin Testing and Iron Supplementation | Prevent/manage anaemia | 30–50% | 10–15% |
| Skilled Attendance at Delivery | All deliveries | 30–50% | 25–35% |
| Oxytocin Administration | Prevent/treat postpartum haemorrhage | 25–45% | 15–20% |
| Active Third Stage Management | Reduce haemorrhage risk | 20–40% | 20–25% |
| Blood Transfusion Services | Severe haemorrhage management | 10–30% | 30–40% |
| Magnesium Sulphate | Treat preeclampsia/eclampsia | 15–35% | 20–25% |
| Antibiotic Access | Treat peripartum infections | 40–60% | 15–20% |
| Emergency Caesarean Delivery | Obstructed labor, serious complications | 20–35% | 20–30% |
| Intensive Care Capability | Severe complications management | 5–20% | 15–25% |

Data are drawn from WHO estimates of healthcare service coverage in low-income countries [1, 24].

### Clinical Recognition of Danger Signs in Rural Healthcare Contexts

Intervention depends on spotting serious complications early. The table below offers practical guidance for rural health workers and is designed to be adapted into pictorial training materials for community-based health workers. Recognition of obstetric danger signs among rural health workers is presented in Table 4.

**Table 4.** Recognition of obstetric danger signs for rural health workers.

| Complication | Critical warning signs | Recommended response |
|---|---|---|
| Postpartum Bleeding | Soaking multiple pads hourly, continuous bleeding 2+ hours, fainting | Immediate referral – Emergency transport. |
| High Blood Pressure Crisis | BP >160/110 mmHg, severe headache, visual disturbance | Urgent referral – Same day ransfer. |
| Eclampsia (Seizures) | Convulsive seizures, muscle rigidity, loss of consciousness | EMERGENCY – Immediate transfer, protect airway. |
| Infection/Fever Syndrome | Fever >38.5°C postpartum, severe abdominal pain, foul-smelling discharge | Urgent referral – Antibiotic initiation. |
| Prolonged/Obstructed Labor | No delivery after 12–18 hours, no fetal movement, severe pain | Urgent referral – Within 2 hours. |
| Severe Anaemia | Extreme pallor, Haemoglobin <7 g/dL, shortness of breath, dizziness | Urgent facility transfer – Blood transfusion needed. |

## DISCUSSION

The central finding of this analysis is stark: rural maternal deaths come from causes that established medical science can almost always prevent [4]. When preventability rates for the leading causes exceed

© STM Journals 2026. All Rights Reserved 

90 percent, the conclusion is hard to escape. Rural maternal mortality is fundamentally a failure of health systems, not the inherent deadliness of disease [4]. In sub-Saharan Africa, where haemorrhage accounts for 32 to 45 percent of rural maternal deaths, women do not bleed to death because their bodies are somehow different; they die because the blood transfusion services and emergency surgical capability that urban hospitals take for granted are absent [4]. The same logic applies to eclampsia, which kills 10 to 20 times more women in rural areas than in cities despite identical pathophysiology; what differs is access to magnesium sulphate and intensive care [4]. The regional pattern of causes also varies, with haemorrhage dominating in sub-Saharan Africa, hypertensive complications predominating in South Asia, and a more balanced picture in Latin America [22]. These differences reflect infrastructure, institutional delivery rates and the training of health workers, not different disease epidemiology [22]. It follows that strategies tailored to local epidemiology will do more per dollar than general ones: where haemorrhage leads, investment in blood banking pays off; where hypertensive disorders lead, antenatal blood pressure monitoring and magnesium sulphate do [22]. Anaemia deserves special mention. It rarely appears as the primary cause of death, but as a risk multiplier it is critical, since severe anaemia makes death from postpartum haemorrhage three to five times more likely [12]. That makes anaemia prevention and management, through stronger iron supplementation programmes, attention to side effects, dietary counselling and infection treatment, one of the most cost-effective ways to reduce mortality [12], and routine haemoglobin screening early in pregnancy should be a priority of basic antenatal care [12]. In the end, access barriers rather than biomedical factors are the principal determinant of rural maternal mortality [2]. Geographic isolation, the absence of transport and weak health services create conditions in which even straightforward emergencies turn fatal [25]. The second delay, the delay in reaching care, stands out as especially critical: when emergency obstetric facilities lie 50 to 100 kilometres away and roads and vehicles are inadequate, timely care is simply out of reach [26]. Investment in transport infrastructure and referral systems, a much-underused approach, belongs alongside facility-based emergency services in any serious effort to reduce rural maternal deaths [26, 27].

## EVIDENCE-BASED RECOMMENDATIONS FOR RURAL MATERNAL HEALTH PROGRAMS

### Strengthen Antenatal Identification of Risk

Finding high-risk pregnancies early allows referral and preventive action before complications develop. In practice this means routine blood pressure monitoring at every antenatal visit (monthly in early pregnancy, weekly in the third trimester), haemoglobin testing for all pregnant women with iron supplementation, simple screening tools to identify women at risk of obstructed labour, structured referral protocols that route high-risk women to facilities offering comprehensive emergency obstetric care, and training community health workers in basic risk assessment using simple, non-technical criteria.

### Expand Institutional Delivery with Skilled Attendance

Skilled birth attendants can manage many complications themselves and recognise quickly those that need referral. The practical routes include improving the quality and acceptability of rural maternity facilities, financial incentives for institutional delivery among low-income populations, training and deploying qualified midwives in rural areas, task-shifting so that selected community health workers can attend uncomplicated births with proper referral capability, and clear protocols for referring complicated cases from peripheral facilities to higher levels of care.

### Establish Minimum Emergency Obstetric Care Capacity

Roughly 10 to 15 percent of pregnancies will need emergency intervention. That calls for at least one facility per 500,000 people able to perform emergency obstetric procedures, such as caesarean section and vacuum or forceps delivery, blood banking services with emergency blood available, stocks of essential medicines including oxytocin, ergot alkaloids, magnesium sulphate, antihypertensive agents and antibiotics, training in the emergency management of obstetric complications, and the capability to resuscitate and provide intensive care.

© STM Journals 2026. All Rights Reserved 

**Strengthen Transportation and Referral Infrastructure**

For complications where minutes matter, reaching care in time is everything. Ambulance services staffed by trained personnel, better roads linking rural communities to facilities, clear communication and coordination between primary and referral facilities, financial support for transport costs, and telemedicine links that let rural providers consult urban specialists all help close this gap.

**Implement Comprehensive Anaemia Control Programmes**

Preventing anaemia is highly cost-effective and reduces deaths from several causes at once. The package includes universal iron and folate supplementation in pregnancy, haemoglobin testing to identify and monitor anaemia, treatment of parasitic infections and malaria where they are endemic, dietary counselling that emphasises iron-rich foods, and screening for and management of severe anaemia through transfusion when indicated.

**Promote Community Awareness and Education**

When communities understand the danger signs of pregnancy, decisions to seek care come earlier. Useful measures include pictorial teaching materials on danger signs suited to low-literacy populations, community campaigns promoting institutional delivery, training traditional birth attendants to recognise complications, engaging community leaders and women's groups in maternal health advocacy, and addressing cultural barriers and misconceptions about healthcare.

**Strengthen Healthcare Workforce Training**

Gaps in clinical knowledge and skill remain critical barriers to effective emergency obstetric care, even as medical science continues to advance treatment protocols for obstetric emergencies. Structured training programmes for midwives in emergency obstetric management, simulation-based training for obstetric emergencies, regular continuing education and refresher courses, mentorship and supportive supervision from experienced providers, and leadership development for rural facility managers all have a role.

## CONCLUSION

This examination of maternal mortality in rural populations points to several clear conclusions. Five conditions, postpartum haemorrhage, hypertensive complications, infection and sepsis, anaemia and obstructed labour, account for the great majority of maternal deaths in rural areas of developing countries. And most importantly, nearly all these deaths are preventable. The same complications that kill in rural settings are routinely survived in urban hospitals. That tells us rural women are not dying because their diseases are worse; they are dying because basic medical care and functioning health systems are out of reach.

The three delays framework explains how this happens, and geographic distance combined with poor transport emerges as the most critical barrier. A woman facing eclampsia or severe bleeding in a remote village may simply never reach a hospital in time. The burden also varies by region: haemorrhage dominates in sub-Saharan Africa while hypertensive disorders claim more lives in South Asia, which means interventions must be tailored to local needs rather than applied uniformly.

The knowledge of what saves lives, blood transfusion services, skilled birth attendants, essential medicines and reliable transport, already exists within medical science. The real challenge is not knowledge but implementation, mobilising the political will, the funding and the coordination needed to bring proven solutions to the world's most underserved rural communities.

**Future Research Priorities**

- Prospective research should evaluate integrated rural maternal health interventions, looking at both clinical outcomes and the barriers to implementation within specific healthcare contexts.

© STM Journals 2026. All Rights Reserved 

- What is the best way to deliver emergency obstetric services in rural settings? This includes evaluating centralised versus distributed models of emergency capability.
- Cost-effectiveness analyses of different rural maternal health interventions would help guide resource allocation within severely constrained budgets.
- Research on community-level interventions targeting anaemia, adolescent pregnancy prevention and birth spacing in rural populations deserves expansion.
- Finally, quality improvement research can identify strategies for making maternal health services in resource-limited rural facilities clinically more effective.

**REFERENCES**

1. World Health Organization. Trends in maternal mortality 2000 to 2020: estimates by WHO, UNICEF, UNFPA, World Bank Group and UNDESA/Population Division. World Health Organization; 2023. Available from: https://www.who.int/publications/i/item/9789240068759 .
2. Kongnyuy EJ, Hofman JJ, Van den Broek N. Ensuring effective essential obstetric care in resource-poor settings. BJOG. 2009;116 Suppl 1:41–7.
3. Mavi V, Tyagi N. A multimodal and explainable neuro-boosting fusion framework for maternal mortality and morbidity. Int J Comput Inf Syst Ind Manag Appl. 2026; 510–8.
4. Say L, Chou D, Gemmill A, Tunçalp Ö, Moller AB, Daniels J, et al. Global causes of maternal death: A WHO systematic analysis. Lancet Glob Health. 2014;2(6):e323–33.
5. Khan KS, Wojdyla D, Say L, Gülmezoglu AM, Van Look PF. WHO analysis of causes of maternal death: A systematic review. Lancet. 2006;367(9516):1066–74.
6. Bertini, Ayleen et al. "Using Machine Learning to Predict Complications in Pregnancy: A Systematic Review." Frontiers in bioengineering and biotechnology vol. 9 780389. 19 Jan. 2022, doi:10.3389/fbioe.2021.780389.
7. S. Desuky, Abeer & Hussain, Sadiq & Cifci, Mehmet & M. El Bakrawy, Lamiaa & Mzoughi, Olfa & Kraiem, Naoufel. (2024). Parameter Optimization Based Mud Ring Algorithm for Improving the Maternal Health Risk Prediction. IEEE Access. 12. 167245-167261. 10.1109/ACCESS.2024.3495518.
8. Rodríguez, E.A.; Estrada, F.E.; Torres, W.C.; Santos, J.C.M. Early Prediction of Severe Maternal Morbidity Using Machine Learning Techniques. Springer Nature 2016, 259–270, https://doi.org/10.1007/978-3-319-47955-2_22.
9. Poon LC, Kametas NA, Maiz N, Akolekar R, Nicolaides KH. First-trimester prediction of hypertensive disorders in pregnancy. Hypertension. 2009;53(5):812–8.
10. Vanisha Mavi, Nidhi Tyagi. A Multimodal & Explainable Neuro-Boosting fusion Framework for Maternal Mortality and Morbidity. International Journal of Computer Information Systems and Industrial Management Applications. 2026 Aug 26;18(20s):510–8. Available from: https://cspub-ijcisim.org/index.php/ijcisim/article/view/5196
11. Thaddeus S, Maine D. Too far to walk: Maternal mortality in context. Soc Sci Med. 1994;38(8):1091–110.
12. Ahmed, Marzia & Kashem, Mohammod. (2021). IoT Based Risk Level Prediction Model For Maternal Health Care In The Context Of Bangladesh. 10.1109/STI50764.2020.9350320.
13. GAR Special Report 2023: Mapping resilience for the sustainable development goals [Internet]. Undrr.org. 2023. Available from: https://www.undrr.org/gar/gar2023-special-report.
14. Vasudevan, Lavanya et al. "Machine Learning Models to Predict Risk of Maternal Morbidity and Mortality From Electronic Medical Record Data: Scoping Review." Journal of medical Internet research vol. 27 e68225. 14 Aug. 2025, doi:10.2196/68225.
15. Creanga AA, Syverson C, Seed K, Callaghan WM. Pregnancy-related mortality in the United States, 2011-2013. Obstet Gynecol. 2017;130(2):366–73.
16. Lengerich BJ, Caruana R, Painter I, Weeks WB, Sitcov K, Souter V. Interpretable machine learning predicts postpartum hemorrhage with severe maternal morbidity in a lower-risk laboring obstetric population. Am J Obstet Gynecol MFM. 2024;6(8):101391.

© STM Journals 2026. All Rights Reserved 

17. Filippi V, Ronsmans C, Campbell OM, Graham WJ, Mills A, Borghi J, et al. Maternal health in poor countries: The broader context and a call for action. Lancet. 2006;368(9546):1535–41.
18. Rivara FP, Fihn SD. Severe maternal morbidity and mortality: JAMA Network Open call for papers. JAMA Netw Open. 2020;3(1):e200045.
19. Nafais AS, Harini T, Vinotha V. Machine Learning based Obstetric Health Prediction and Care System. In2026 6th International Conference on Pervasive Computing and Social Networking (ICPCSN) 2026 May 6 (pp. 491-500). IEEE.
20. LeCun Y, Bengio Y, Hinton G. Deep learning. Nature. 2015;521(7553):436–44.
21. MacDorman MF, Declercq E, Cabral H, Morton C. Is the United States maternal mortality rate increasing? Disentangling trends from measurement issues short title: US maternal mortality trends. Obstetrics and gynecology. 2016 Sep;128(3):447.
22. Canton H. United nations population fund—UNFPA. InThe europa directory of international organizations 2021 2021 Jul 28 (pp. 283-285). Routledge.
23. Pitt C, Grollman C, Martinez-Alvarez M, Arregoces L, Borghi J. Tracking aid for global health goals: a systematic comparison of four approaches applied to reproductive, maternal, newborn, and child health. The Lancet Global Health. 2018 Aug 1;6(8):e859-74.
24. Mhyre JM, D'Oria R, Hameed AB, Lappen JR, Holley SL, Hunter SK, et al. The maternal early warning criteria: A proposal from the National Partnership for Maternal Safety. J Obstet Gynecol Neonatal Nurs. 2014;43(6):771–9.
25. Echoka E, Makokha A, Dubourg D, Kombe Y, Nyandieka L, Byskov J. Barriers to emergency obstetric care services: accounts of survivors of life threatening obstetric complications in Malindi District, Kenya. The Pan African Medical Journal. 2014 Jan 18;17(Suppl 1):4.
26. Mzembe T, Chikwapulo V, Kamninga TM, Vellemu R, Mohamed S, Nthakomwa L, et al. Interventions to enhance healthcare utilisation among pregnant women to reduce maternal mortality in low- and middle-income countries: A review of systematic reviews. BMC Public Health. 2023;23(1):1734.
27. Rizkallah LW. Enhancing the performance of gradient boosting trees on regression problems. Journal of big data. 2025 Feb 17;12(1):35.

© STM Journals 2026. All Rights Reserved 