

Enhancing Fiber-Digestibility in Dairy Cows: Influence of Dietary, Management, and Environmental Factors

Md. Emran Hossain*

Abstract

The digestibility of fiber in dairy cows is a critical factor for optimizing their nutritional efficiency and overall productivity. This study explores the key dietary, management, and environmental factors influencing fiber-digestibility in the rumen. Various aspects of feed composition, such as fiber content, protein-to-fiber ratio, and the presence of anti-nutritional factors, play a significant role in determining microbial activity and rumen fermentation efficiency. Physical feed characteristics, including particle size, feed processing methods, and forage maturity, further influence the breakdown of fiber. Environmental factors, such as ambient temperature, humidity, and heat stress, can disrupt rumen function and reduce microbial efficiency, negatively impacting fiber digestion. In addition, management practices – such as feed mixing efficiency, stocking density, and access to quality pasture – are essential for maintaining optimal rumen conditions and supporting effective fiber utilization. The study also highlights the importance of microbial diversity and the presence of specific microbial populations, including fibrolytic bacteria, in enhancing fiber digestion. Effective use of dietary interventions – such as enzyme supplementation, ionophores, and prebiotics – can further improve fiber digestibility. By understanding the interplay of these factors, this review provides insights into strategies for enhancing fiber digestion, which is essential for improving feed efficiency, animal health, and sustainable dairy farming practices.

Keywords: Dairy cows, enzyme supplementation, environmental factors, feed composition, feed management, fiber digestibility

INTRODUCTION

Fiber-digestibility in dairy cows plays a fundamental role in their overall productivity and health [1]. The efficiency with which fiber is broken down in the rumen influences feed intake, nutrient absorption, and ultimately milk production [2]. Ruminants, particularly dairy cows, rely on complex microbial populations in the rumen to degrade structural carbohydrates such as cellulose and hemicellulose [3]. The microbial breakdown of fiber results in the production of volatile fatty acids (VFAs), which serve as a major energy source for the animal. Understanding the factors that affect fiber digestibility is crucial for optimizing dairy nutrition and enhancing production efficiency [4].

*Author for Correspondence

Md. Emran Hossain
E-mail: emran@cvasu.ac.bd

Professor, Department of Animal Science and Nutrition,
Chattogram Veterinary and Animal Sciences University,
Khulshi, Chattogram-4225, Bangladesh

Received Date: May 25, 2026
Accepted Date: June 27, 2026
Published Date: June 28, 2026

Citation: Md. Emran Hossain. Enhancing Fiber-Digestibility in Dairy Cows: Influence of Dietary, Management, and Environmental Factors. Research & Reviews: A Journal of Microbiology and Virology. 2026; 16(2): 37–50p.

Several factors influence fiber digestibility, with dietary composition being a key determinant. The fiber content, presence of lignin, protein-to-fiber ratio, and non-fiber carbohydrate levels within the diet can all affect microbial fermentation in the rumen [5–7]. In addition, the physical characteristics of the feed, such as particle size, forage maturity, and feed processing methods, play a significant role in determining how effectively microbes can break down fiber [8, 9]. Environmental factors, including ambient

temperature, humidity, and heat stress, can also impact microbial activity and feed intake, further affecting fiber degradation [10–13].

This study offers a comprehensive analysis of the complex interactions between dietary, management, and environmental factors that influence fiber-digestibility in dairy cows. It is unique in its holistic approach, integrating multiple factors that have often been studied in isolation. The novelty of this study lies in its detailed exploration of how factors – such as feed mixing efficiency, microbial population diversity, and environmental stressors like heat and cold stress – can collectively affect rumen function and fiber digestion. By considering these variables in conjunction, the study provides novel insights into strategies for improving fiber digestibility, thus contributing to more sustainable and efficient dairy farming practices.

FEED COMPOSITION

Fiber Content

The total fiber content in the feed directly influences ruminal digestion as it determines the availability of structural carbohydrates for microbial degradation [14–16]. Optimal fiber levels stimulate chewing and saliva production, maintaining rumen pH. Excess fiber reduces digestibility, while insufficient fiber fails to provide structural support for microbial adhesion and fermentation, compromising fiber breakdown and nutrient absorption.

Lignin Concentration

Lignin is a non-digestible component of plant cell walls that hinders microbial access to cellulose and hemicellulose. Higher lignin concentration reduces fiber digestibility by forming a physical barrier against enzymatic degradation [2, 17]. Reducing lignin content through plant breeding or pre-treatment methods enhances ruminal fibrolytic activity, allowing greater breakdown of structural carbohydrates and improving nutrient availability for the dairy cow.

Cellulose Content

Cellulose serves as a major carbohydrate source for rumen microbes. Its digestibility depends on its accessibility within plant cell walls. Higher cellulose levels provide energy for microbial growth, but excessive crystalline cellulose reduces breakdown efficiency [18, 19]. Optimal cellulose content ensures sustained microbial activity, promoting the production of volatile fatty acids (VFAs) and improving energy supply for the dairy cow.

Hemicellulose Levels

Hemicellulose, a more soluble fiber fraction, complements cellulose in providing structural carbohydrates for fermentation [18–21]. Its branched structure is more easily degraded by rumen microbes compared to cellulose. Adequate hemicellulose levels improve fiber digestibility by enhancing microbial attachment and enzyme activity. Optimizing hemicellulose proportions in the diet ensures higher ruminal volatile fatty acid production and better nutrient utilization.

Protein-to-Fiber Ratio

A balanced protein-to-fiber ratio supports optimal microbial activity in the rumen. Fibrolytic bacteria require nitrogen from protein for enzyme production, while fiber serves as their substrate. Insufficient protein limits microbial growth, whereas excess protein may cause ammonia toxicity [18–21]. Maintaining the right ratio enhances fiber degradation, improves volatile fatty acid production, and supports efficient energy utilization in the dairy cow.

Non-Fiber Carbohydrate Proportion

Non-fiber carbohydrates (NFCs), including starch and sugars, provide readily fermentable substrates that stimulate microbial activity. A balanced proportion of NFCs in the diet supports fibrolytic bacteria by improving overall rumen fermentation dynamics [22–24]. Excess NFCs can cause ruminal acidosis,

inhibiting fiber digestion. Proper NFC levels ensure synchronization between fiber and carbohydrate fermentation, optimizing microbial growth and fiber digestibility.

Mineral Content

Minerals – like calcium, phosphorus, magnesium, and sulfur – play crucial roles in rumen microbial activity and enzyme function. Adequate mineral content supports fibrolytic bacteria by maintaining rumen fluid osmolarity and pH [25–27]. Specific minerals, such as sulfur, enhance the synthesis of sulfur-containing amino acids essential for fiber-degrading enzymes. Balanced mineral supplementation enhances fiber degradation and improves overall nutrient availability.

Fat Content in Feed

Excessive fat inhibits fiber digestibility by coating feed particles and reducing microbial access. However, optimal fat levels provide energy and support microbial activity without compromising fiber degradation. Protected or bypass fats minimize negative effects on rumen fermentation [28–30]. Proper fat supplementation ensures microbial efficiency, prevents fiber inhibition, and improves the energy density of diets without reducing ruminal fibrolytic activity.

Presence of Anti-Nutritional Factors

Anti-nutritional factors, such as tannins and phytates, inhibit rumen microbial activity by binding nutrients or reducing enzyme availability. Their presence negatively affects fiber digestion by limiting microbial access to structural carbohydrates [13, 31–33]. Strategies – such as feed processing, detoxification, or supplementation with tannin-tolerant microbes – can mitigate these effects, improve rumen fermentation efficiency and enhancing fiber-digestibility in dairy cows.

Water-Soluble Carbohydrates

Water-soluble carbohydrates (WSCs) provide an immediate energy source for rumen microbes, stimulating microbial growth and enzyme production. Sufficient WSC levels enhance the degradation of structural carbohydrates by fibrolytic bacteria [7, 9, 24, 34]. They also buffer fiber digestion by maintaining fermentation balance. Including forages with high WSC content optimizes microbial synergy and ensures effective fiber breakdown, improving energy utilization and productivity in dairy cows.

PHYSICAL FEED CHARACTERISTICS

Particle Size

Particle size significantly affects fiber-digestibility in the rumen by influencing microbial attachment and fermentation rates. Smaller particles increase the surface area for microbial colonization, enhancing enzymatic degradation [35–37]. However, excessively fine particles may reduce rumination and buffering through saliva. Optimizing particle size ensures effective microbial activity, promotes rumen pH stability, and enhances the overall breakdown of crude fiber.

Feed Processing Method

Processing methods – like grinding, pelleting, or extrusion – can alter fiber structure and accessibility. Proper feed processing increases digestibility by reducing particle size, disrupting lignocellulosic bonds, and exposing structural carbohydrates to microbial degradation [36, 37]. Over-processing, however, may negatively impact the rumen environment. Selecting appropriate processing methods enhances fiber utilization, microbial efficiency, and nutrient availability for dairy cows.

Dry Matter Content

Dry matter content influences feed intake and microbial activity in the rumen. High dry matter levels may reduce microbial access to fiber due to lower water availability, while overly wet feeds dilute nutrients and disrupt fermentation [17, 38, 39, 40]. Optimal dry matter content ensures proper microbial attachment, enhances fermentation efficiency, and maintains fiber-digestibility by supporting a balanced rumen environment.

Forage Maturity

Forage maturity directly affects fiber digestibility as lignification increases with plant age, reducing microbial access to structural carbohydrates. Harvesting forages at an optimal growth stage ensures higher cellulose and hemicellulose levels with reduced lignin content [39, 40]. Properly timed harvests maximize digestibility, improve rumen fermentation efficiency, and support better nutrient utilization in dairy cows.

Forage-to-Concentrate Ratio

The forage-to-concentrate ratio determines the balance between fiber and readily fermentable carbohydrates in the diet. Higher forage proportions promote chewing and saliva production, stabilizing rumen pH and supporting fiber digestion [14, 41, 42]. Excessive concentrate levels can cause acidosis, reducing fiber degradation. Balancing the ratio enhances microbial activity, improves volatile fatty acid production, and ensures effective crude fiber breakdown in the rumen.

Silage Chop Length

Silage chop length affects particle size and the physical structure of the feed, influencing rumen fermentation and fiber digestibility [43]. Properly chopped silage facilitates microbial attachment and increases the surface area for enzymatic activity. Overly short or long chops can disrupt rumen function or reduce chewing efficiency. Optimizing chop length enhances microbial efficiency and promotes effective fiber breakdown.

Feed Texture

Feed texture determines the physical characteristics of the diet, affecting chewing behavior and microbial attachment. Coarser textures encourage rumination and saliva production, maintaining optimal rumen pH for fiber digestion [14, 41, 42]. Conversely, excessively fine or compacted feed may impair microbial colonization. Ensuring an appropriate feed texture improves microbial activity, enhances fiber breakdown, and supports overall rumen fermentation dynamics.

Moisture Level of Forage

Forage moisture levels influence fermentation dynamics and microbial activity in the rumen. Excessively wet forage may dilute nutrients and disrupt microbial attachment, while overly dry forage reduces microbial access to structural carbohydrates [39, 40, 44]. Proper moisture levels ensure optimal fermentation conditions, enhance fiber degradation by fibrolytic bacteria, and improve nutrient availability for the dairy cow.

MICROBIAL ACTIVITY

Microbial Population Diversity

A diverse microbial population ensures the availability of specialized microbes capable of degrading various fiber components. Greater diversity promotes resilience and adaptability of the rumen ecosystem, optimizing fiber degradation even under dietary fluctuations [18, 27]. Ensuring microbial diversity through balanced nutrition enhances fermentation efficiency, volatile fatty acid production, and nutrient utilization, improving fiber-digestibility in dairy cows.

Abundance of Fibrolytic Bacteria

Fibrolytic bacteria are primary agents of fiber breakdown in the rumen, producing enzymes like cellulases and hemicellulases [18, 27]. Higher fibrolytic bacterial populations enhance the degradation of structural carbohydrates, increasing volatile fatty acid production. Maintaining optimal environmental conditions, such as rumen pH and nitrogen availability, support fibrolytic bacterial growth and ensures efficient fiber digestion in dairy cows.

Presence of Protozoa

Protozoa assist in fiber digestion by engulfing plant particles and interacting with fibrolytic bacteria. They play a complementary role in breaking down cellulose and hemicellulose, enhancing overall fiber

digestibility [10, 45–47]. However, protozoa also compete for nutrients, which may limit bacterial efficiency. Balancing protozoal populations ensures microbial synergy and promotes effective fiber degradation in the rumen.

Methanogen Activity

Methanogens, while primarily involved in methane production, contribute indirectly to fiber-digestibility by maintaining hydrogen balance in the rumen. Effective hydrogen removal supports fibrolytic microbial activity and prevents fermentation inhibition. Excessive methanogenesis reduces energy efficiency [10, 45–47]. Strategies to modulate methanogen activity, such as dietary additives, can optimize fiber degradation and improve energy utilization in dairy cows.

Microbial Synergism

Microbial synergism involves cooperative interactions among different microbial groups, enhancing fiber degradation. Fibrolytic bacteria degrade complex carbohydrates, while other microbes utilize fermentation by-products, preventing accumulation and maintaining optimal conditions [48, 49–51]. Promoting synergistic interactions through balanced diets ensures efficient microbial activity, maximizes fiber breakdown, and improves nutrient availability for dairy cows.

Competition Among Microbes

Competition among microbial populations affects fiber-digestibility by influencing resource allocation and enzymatic activity. Excessive competition may hinder fibrolytic bacteria, reducing fiber degradation [48–51]. Proper dietary formulation minimizes competition by providing adequate nitrogen and energy substrates. Promoting microbial coexistence supports efficient fermentation and enhances fiber digestion in the rumen.

Rumen Microbiome Stability

A stable rumen microbiome ensures consistent fiber degradation by maintaining an optimal microbial population under dietary or environmental changes. Stability minimizes disruptions in fermentation dynamics, supporting fibrolytic activity and nutrient utilization [48–51]. Achieving microbiome stability through gradual dietary transitions and balanced nutrient supply enhances fiber digestibility and supports rumen health in dairy cows.

CHEMICAL TREATMENTS

Alkali Treatment of Feed

Alkali treatment, such as sodium hydroxide application, breaks down lignocellulosic bonds in fiber, enhancing microbial access to cellulose and hemicellulose [23, 40]. This chemical treatment disrupts the crystalline structure of fiber, improving digestibility. Properly treated feeds provide more fermentable substrates for rumen microbes, increasing volatile fatty acid production and enhancing energy availability for dairy cows.

Acid Treatment of Feed

Acid treatment, using chemicals like sulfuric or acetic acid, helps hydrolyze hemicellulose and disrupts lignin-carbohydrate complexes, improving fiber digestibility. It increases the solubility of structural carbohydrates, facilitating microbial access [52]. While effective, precise application is necessary to avoid rumen pH disturbances. Acid-treated feeds promote efficient microbial fermentation and enhance nutrient utilization in dairy cows.

Ammoniation of Forages

Ammoniation, using anhydrous ammonia or urea, enhances fiber digestibility by breaking lignin-carbohydrate bonds and increasing forage nitrogen content [53, 54]. This treatment improves microbial attachment and enzymatic activity on cellulose and hemicellulose. Ammoniated forages provide additional non-protein nitrogen for microbial growth, boosting fiber degradation and nutrient availability for dairy cows, particularly when fed low-quality forages.

Steam Explosion Process

The steam explosion process uses high-pressure steam followed by rapid decompression to disrupt plant cell walls, breaking lignin bonds and increasing fiber accessibility [52–55]. This method enhances the digestibility of cellulose and hemicellulose, making structural carbohydrates more available for microbial fermentation. Properly processed feed promotes microbial activity, increases volatile fatty acid production, and improves overall energy utilization in dairy cows.

Ensiling Duration

The duration of ensiling affects fiber digestibility as microbial fermentation reduces structural carbohydrate content over time. Longer ensiling periods break down hemicellulose and partially degrade lignin bonds, enhancing forage digestibility. However, overextended ensiling can lead to nutrient losses [52–55]. Optimizing ensiling duration ensures improved fiber utilization, better fermentation quality, and enhanced nutrient availability for rumen microbes.

Use of Chemical Preservatives

Chemical preservatives – like propionic acid or formaldehyde – inhibit spoilage during storage, maintaining fiber integrity and nutrient availability. Some preservatives also modify fiber structure, increasing digestibility by enhancing microbial access to carbohydrates [23, 53, 56]. Proper application of chemical preservatives ensures feed stability, reduces nutrient losses, and supports effective rumen fermentation, improving fiber breakdown in dairy cows.

Urea Treatment of Straws

Urea treatment improves the digestibility of low-quality straws by releasing ammonia, which breaks lignin bonds and increases nitrogen content. This treatment enhances microbial attachment and enzymatic degradation of structural carbohydrates [23, 53, 56]. Urea-treated straws provide a cost-effective, nitrogen-enriched feed source, improving fiber digestibility, microbial efficiency, and overall nutritional value for dairy cows.

ANIMAL FACTORS

Rumen Size

Rumen size influences the volume of feed that can be retained and processed [57–60]. A larger rumen provides greater capacity for microbial fermentation, allowing more time for fiber breakdown. Enhanced rumen size supports better retention of fibrous materials, promoting microbial activity and enzymatic degradation, which improves the digestibility of crude fiber in dairy cows.

Rumen pH

Rumen pH is critical for maintaining an optimal environment for fibrolytic bacteria. A stable pH around 6.0–7.0 enhances microbial activity and fiber digestion. Acidosis or alkalosis inhibits fibrolytic microbes and enzymatic activity [57, 58, 60]. Proper dietary balance, particularly between forages and concentrates, helps stabilize rumen pH, improving fiber digestibility and microbial efficiency in dairy cows.

Retention Time in the Rumen

Retention time affects the extent of fiber fermentation. Longer retention periods allow more thorough microbial attachment and enzymatic activity on fiber particles [57, 58, 60]. Adequate retention time ensures efficient degradation of structural carbohydrates, increasing volatile fatty acid production. Factors – such as particle size and feeding frequency – influence retention time, directly affecting fiber digestibility.

Saliva Secretion Rate

Saliva secretion provides bicarbonates that buffer rumen pH, maintaining an optimal environment for fiber digestion [57, 58, 60]. Increased saliva production, stimulated by chewing fibrous feeds,

enhances microbial activity and prevents acid buildup. Adequate saliva secretion supports rumen health, promotes fibrolytic activity, and improves crude fiber-digestibility in dairy cows.

Chewing Efficiency

Chewing efficiency determines the degree of feed particle breakdown and saliva production. Effective chewing reduces particle size, increasing surface area for microbial attachment and fermentation [57, 58, 60]. It also stimulates saliva secretion, buffering rumen pH. Enhanced chewing efficiency supports microbial activity, accelerates fiber degradation, and ensures better nutrient utilization in dairy cows.

Age of the Animal

Age affects fiber digestibility due to variations in rumen development and microbial populations. Young animals with immature rumen systems may have limited fiber digestion compared to mature cows [57, 58, 60]. Age-specific dietary adjustments, such as gradual fiber inclusion, promote rumen microbial adaptation and improve fiber degradation, optimizing nutrient utilization throughout the animal's lifecycle.

Breed Differences

Breed differences influence fiber digestibility through variations in rumen size, microbial populations, and feed intake capacity [57, 58, 60]. Certain breeds are genetically predisposed to better fiber utilization due to adaptations in their digestive physiology. Understanding breed-specific nutritional requirements ensures tailored feeding strategies that enhance fiber digestibility and overall productivity in dairy cows.

Genetic Predisposition

Genetic predisposition affects rumen function, microbial efficiency, and enzymatic activity. Animals with favorable genetics for fiber digestion exhibit improved nutrient utilization and productivity [57, 58, 60]. Selective breeding programs focused on traits – like rumen efficiency and microbial synergy – enhance fiber digestibility, promoting sustainable and efficient dairy farming practices.

Animal Health Condition

Animal health directly impacts rumen function and microbial activity. Illnesses, such as acidosis or parasitic infections, impair fiber digestion by disrupting the rumen environment [57, 58, 60]. Maintaining optimal health through balanced nutrition, disease prevention, and proper management supports microbial efficiency, promotes fiber breakdown, and enhances overall digestibility of crude fiber in dairy cows.

DIETARY INTERVENTIONS

Addition of Rumen Buffers

Rumen buffers, such as sodium bicarbonate, stabilize rumen pH by neutralizing excess acidity from fermentation [34, 61, 62]. Maintaining an optimal pH prevents inhibition of fibrolytic bacteria, ensuring effective fiber breakdown. Buffers enhance microbial activity and enzymatic efficiency, improving volatile fatty acid production and nutrient utilization. Strategic inclusion of buffers supports fiber digestibility and rumen health in dairy cows.

Inclusion of Feed Additives

Feed additives – like yeast cultures and plant extracts – enhance fiber digestibility by promoting microbial growth and fermentation efficiency. Additives stimulate fibrolytic bacteria, improve enzyme activity, and modulate rumen fermentation dynamics [34, 61, 62]. These interventions optimize the rumen environment, leading to better fiber degradation, increased volatile fatty acid production, and improved energy utilization in dairy cows.

Ionophore Supplementation

Ionophores, such as monensin, improve fiber digestibility by altering rumen microbial populations. They reduce competition from less efficient microbes, allowing fibrolytic bacteria to thrive [34, 61, 62]. Ionophores also enhance energy efficiency by modulating volatile fatty acid production. Proper supplementation increases microbial synergy, promotes fiber degradation, and improves overall feed conversion efficiency in dairy cows.

Use of Prebiotics

Prebiotics, such as fructooligosaccharides, selectively stimulate beneficial microbial populations in the rumen [34, 61, 62]. Enhanced fibrolytic bacterial activity promotes efficient fiber breakdown and fermentation. Prebiotics also support microbial stability, preventing disturbances in rumen function. Including prebiotics in diets improves fiber-digestibility, volatile fatty acid production, and nutrient availability for dairy cows.

Use of Probiotics

Probiotics, including live microbial cultures, enhance fiber digestion by boosting microbial populations and stabilizing the rumen environment. Probiotic supplementation improves enzymatic activity and microbial synergy, ensuring efficient fiber breakdown [34, 61, 62]. These benefits promote better volatile fatty acid production, rumen health, and nutrient absorption, enhancing fiber-digestibility in dairy cows.

Inclusion of Enzymes

Exogenous enzymes, such as cellulases and hemicellulases, directly degrade structural carbohydrates, enhancing fiber digestibility. Enzymes complement microbial activity, accelerating fiber breakdown and increasing nutrient availability [34, 61, 62]. Strategic enzyme inclusion improves volatile fatty acid production, energy utilization, and overall rumen efficiency, particularly when fed low-quality forages.

Mineral Supplements

Minerals – such as calcium, magnesium, and sulfur – are essential for microbial enzyme activity and fiber digestion. Adequate mineral supplementation enhances fibrolytic bacterial function and fermentation efficiency [34, 61, 62]. Balanced mineral intake ensures microbial health, promotes fiber degradation, and supports optimal rumen performance, improving fiber-digestibility in dairy cows.

Feeding Frequency

Frequent feeding stabilizes rumen pH and ensures a consistent supply of nutrients for microbial activity. Reduced fluctuations in fermentation conditions enhance fibrolytic bacterial efficiency and fiber breakdown [13]. Proper feeding intervals promote optimal rumen function, increase volatile fatty acid production, and improve fiber-digestibility in dairy cows.

Feeding Timing

Feeding timing influences rumen fermentation dynamics and microbial efficiency. Providing fiber-rich feeds during peak microbial activity enhances fiber breakdown [13]. Coordinating feed intake with rumen fermentation rhythms optimizes microbial populations and enzymatic activity, promoting efficient fiber digestion and nutrient utilization in dairy cows.

ENVIRONMENTAL FACTORS

Ambient Temperature

Ambient temperature influences microbial activity and feed intake in dairy cows. Optimal temperatures support stable rumen function and fibrolytic bacterial growth [29, 30, 33, 63]. High or low temperatures can disrupt fermentation processes and reduce fiber digestibility. Providing a thermoneutral environment ensures effective microbial activity, improving structural carbohydrate breakdown and nutrient utilization in the rumen.

Humidity Levels

Humidity levels affect feed stability and microbial fermentation efficiency. High humidity promotes feed spoilage and reduces fiber availability, while extremely low humidity can lead to reduced feed palatability [29, 30, 33, 63]. Maintaining moderate humidity supports feed quality and optimal microbial activity, ensuring improved fiber digestibility and rumen fermentation in dairy cows.

Season of Feeding

Seasonal changes influence forage quality, feed intake, and microbial efficiency. During certain seasons, higher fiber or lignin content in forages may reduce digestibility [29, 30, 33, 63]. Properly balancing diets with supplements or additives based on seasonal variations ensures consistent microbial activity and fiber breakdown, optimizing nutrient utilization for dairy cows throughout the year.

Heat Stress

Heat stress disrupts rumen function by reducing feed intake and microbial activity, impairing fiber digestibility. Elevated body temperatures decrease chewing and saliva secretion, leading to inefficient fermentation [29, 30, 33, 63]. Mitigating heat stress through cooling systems, shaded areas, and balanced diets maintains rumen stability, enhances microbial efficiency, and improves fiber utilization in dairy cows.

Cold Stress

Cold stress increases energy requirements, potentially altering rumen fermentation dynamics and reducing fiber digestibility. Inadequate nutrition during cold periods can compromise microbial efficiency and fiber degradation [29, 30, 33, 63]. Providing energy-dense diets and adequate shelter helps maintain rumen function and microbial activity, ensuring effective fiber breakdown and nutrient absorption in dairy cows.

Access to Drinking Water

Access to sufficient drinking water supports rumen fermentation and saliva production, aiding fiber digestion. Water facilitates the movement of feed particles within the rumen, promoting microbial attachment and fermentation [29, 30, 33, 63]. Inadequate water access reduces fiber breakdown and volatile fatty acid production, impairing digestibility. Ensuring unrestricted water availability improves overall rumen efficiency.

Water Quality

Water quality directly affects microbial activity and rumen health. Contaminated or imbalanced water sources may disrupt fermentation, reducing fiber digestibility [29, 30, 33, 63]. Clean and mineral-balanced water supports optimal microbial populations and enzymatic activity, enhancing fiber breakdown and nutrient utilization for dairy cows.

Housing Conditions

Housing conditions influence feed accessibility, comfort, and microbial activity in dairy cows. Proper ventilation, space, and hygiene prevent stress and promote consistent feed intake [29, 30, 33, 63]. Comfortable housing supports rumen health and efficient fermentation, enhancing fibrolytic bacterial activity and fiber digestibility, ultimately improving productivity in dairy systems.

MANAGEMENT PRACTICES

Feed Mixing Efficiency

Efficient feed mixing ensures uniform distribution of nutrients and prevents sorting by dairy cows. Consistent mixing improves the balance between forage and concentrate, promoting optimal microbial fermentation and fiber digestion [13]. Properly mixed rations enhance nutrient availability, support stable rumen function, and maximize fiber breakdown, leading to improved digestibility and overall productivity.

Feed Storage Conditions

Proper feed storage prevents spoilage, mold growth, and nutrient loss, preserving feed quality. High-quality stored feed maintains its fiber structure and nutritional value, ensuring effective microbial fermentation [13]. Controlled storage conditions reduce anti-nutritional factors and toxins, supporting rumen health and enhancing fiber-digestibility in dairy cows.

Feed Bunk Management

Effective feed bunk management ensures cows have constant access to fresh and uncontaminated feed. Regular cleaning and proper feed distribution prevent feed sorting and spoilage [13]. Consistent access to balanced rations promotes stable rumen fermentation, microbial activity, and improved fiber digestibility, contributing to optimal cow health and productivity.

Feed Delivery Method

The feed delivery method impacts feed intake and rumen fermentation dynamics. Frequent and evenly distributed feed delivery reduces competition and ensures consistent nutrient availability [13]. Improved delivery methods enhance microbial efficiency, optimize fiber digestion, and maintain rumen stability, supporting better nutrient utilization and performance in dairy cows.

Stocking Density

Optimal stocking density minimizes competition for feed and promotes adequate feed intake. Overcrowding can lead to reduced feeding time and disrupted rumen fermentation [13]. Properly managed stocking density ensures all animals consume balanced rations, supporting efficient microbial activity and improved fiber-digestibility in dairy cows.

Access to Pasture

Access to pasture provides cows with high-quality forages rich in fiber and nutrients. Grazing stimulates natural feeding behavior, increasing chewing time and saliva production [13]. Pasture access enhances microbial fermentation, promotes efficient fiber digestion, and improves overall nutrient utilization, contributing to healthier and more productive dairy cows.

Grouping Strategies

Grouping strategies based on age, production stage, or nutritional requirements optimize feeding efficiency and reduce competition [13]. Proper grouping ensures animals receive tailored diets that support microbial activity and fiber digestibility. Segregating cows with similar needs promotes uniform feed intake and effective fiber breakdown, enhancing overall rumen performance and productivity.

Feed testing Protocols

Regular feed testing ensures the nutritional quality of feed and identifies imbalances or deficiencies. Accurate testing allows for adjustments in rations to optimize fiber content and microbial activity [13]. Feed testing supports effective fermentation, improves fiber digestibility, and enhances the nutritional value of diets, ensuring sustainable productivity in dairy cows.

CONCLUSION

Fiber-digestibility in dairy cows is influenced by a complex interplay of dietary, management, and environmental factors. Optimizing feed composition, microbial activity, and environmental conditions is essential for improving rumen function and maximizing nutrient utilization. Effective feed management practices, such as proper mixing and storage, along with the use of dietary additives, can enhance fiber breakdown. Environmental factors, including temperature and humidity, should be carefully managed to mitigate stress and support microbial efficiency. This study emphasizes the importance of a holistic approach to fiber digestibility, providing valuable insights for enhancing dairy cow nutrition and promoting sustainable farming practices. Implementing these strategies can lead to improved productivity, animal health, and overall farm efficiency.

Future Directions

Future research should focus on identifying novel dietary additives, such as specific prebiotics, probiotics, and enzymes, that can further enhance fiber-digestibility in dairy cows. Investigating the role of rumen microbiome modulation through dietary interventions could lead to more targeted strategies for improving fiber breakdown. Additionally, exploring the genetic basis of fiber digestion in dairy cows may offer opportunities for selective breeding programs aimed at enhancing rumen efficiency. Further studies on the impact of climate change on environmental factors, such as heat stress and feed quality, are crucial for developing adaptive management strategies. Additionally, improving precision livestock farming technologies to monitor and manage individual cow's rumen function and fiber utilization will provide more personalized approaches to feed and health management. Interdisciplinary research incorporating nutrition, microbiology, genomics, and environmental science will be key to developing sustainable solutions for enhancing fiber digestibility and overall dairy production.

REFERENCES

1. Zebeli Q, Aschenbach JR, Tafaj M, Boguhn J, Ametaj BN, Drochner W. Invited review: Role of physically effective fiber and estimation of dietary fiber adequacy in high-producing dairy cattle. *J Dairy Sci.* 2012;95(3):1041–1056. doi: 10.3168/jds.2011-4421.
2. Zhao XH, Zhang T, Xu M, Yao JH. Effects of physically effective fiber on chewing activity, ruminal fermentation, and digestibility in goats. *J Anim Sci.* 2011;89(2):501–509. doi: 10.2527/jas.2010-3013.
3. Galyean ML, Goetsch AL. Utilization of forage fiber by ruminants. *Forage Cell Wall Struct Dig.* 2015;33–71. doi: 10.2134/1993.foragecellwall.c2.
4. Fustini M, Palmonari A, Canestrari G, Bonfante E, Mammi LU, Pacchioli MT, Sniffen GC, Grant RJ, Cotanch KW, Formigoni A. Effect of undigested neutral detergent fiber content of alfalfa hay on lactating dairy cows: Feeding behavior, fiber digestibility, and lactation performance. *Journal of dairy science.* 2017 Jun 1;100(6):4475–83.
5. Shi R, Dong S, Mao J, Wang J, Cao Z, Wang Y, Li S, Zhao G. Dietary neutral detergent fiber levels impacting dairy cows' feeding behavior, rumen fermentation, and production performance during the period of peak-lactation. *Animals.* 2023 Sep 10;13(18):2876.
6. Beane KE, Redding MC, Wang X, Pan JH, Le B, Cicalo C, Jeon S, Kim YJ, Lee JH, Shin EC, Li Y. Effects of dietary fibers, micronutrients, and phytonutrients on gut microbiome: a review. *Applied Biological Chemistry.* 2021 Dec;64(1):36.
7. Cavallini D, Concolino A, Mammi L, et al. Factors influencing fiber digestibility in dairy cows. *J Dairy.* 2020. [Online]. Available: <https://cris.unibo.it/handle/11585/804412%0Ahttps://cris.unibo.it/bitstream/11585/804412/2/PIIS002203022030816X.pdf>
8. Aschalew ND, Wang T, Qin GX, Zhen YG, Zhang XF, Chen X, et al. Effects of physically effective fiber on rumen and milk parameters in dairy cows: a review. 2020:1317–1323.
9. Xie B, Gao J, Zhao G. Dietary supplementation with sodium sulfate improves rumen fermentation, fiber digestibility, and the plasma metabolome through modulation of rumen bacterial communities in steers. *Appl Environ Microbiol.* 2020;86(22):1–18. doi: 10.1128/AEM.01412-20.
10. Makki K, Deehan EC, Walter J, Bäckhed F. The impact of dietary fiber on gut microbiota in host health and disease. *Cell host & microbe.* 2018 Jun 13;23(6):705–15.
11. Korcz E, Kerényi Z, Varga L. Dietary fibers, prebiotics, and exopolysaccharides produced by lactic acid bacteria: Potential health benefits with special regard to cholesterol-lowering effects. *Food Funct.* 2018;9(6):3057–3068. doi: 10.1039/c8fo00118a.
12. Oldick BS, Firkins JL. Effects of degree of fat saturation on fiber digestion and microbial protein synthesis when diets are fed twelve times daily. *J Anim Sci.* 2000;78(9):2412–2420. doi: 10.2527/2000.7892412x.
13. Amminga ST. Influence of feeding management on ruminant fiber digestibility. *Forage Cell Wall Struct Dig.* 2015:571–602. doi: 10.2134/1993.foragecellwall.c22.

14. Chen H, Wang C, Huasai S, Chen A. Effects of dietary forage to concentrate ratio on nutrient digestibility, ruminal fermentation and rumen bacterial composition in Angus cows. *Sci Rep*. 2021;11(1):17023. doi: 10.1038/s41598-021-96580-5.
15. Yang K, Wei C, Zhao GY, Xu ZW, Lin SX. Effects of dietary supplementing tannic acid in the ration of beef cattle on rumen fermentation, methane emission, microbial flora and nutrient digestibility. *J Anim Physiol Anim Nutr (Berl)*. 2017;101(2):302–310. doi: 10.1111/jpn.12531.
16. Huhtanen P, Rinne M, Nousiainen J. A meta-analysis of feed digestion in dairy cows. 2. The effects of feeding level and diet composition on digestibility. *J Dairy Sci*. 2009;92(10):5031–5042. doi: 10.3168/jds.2008-1834.
17. Oba M, Allen MS. Evaluation of the importance of the digestibility of neutral detergent fiber from forage: Effects on dry matter intake and milk yield of dairy cows. *J Dairy Sci*. 1999;82(3):589–596. doi: 10.3168/jds.S0022-0302(99)75271-9.
18. Hendawy AO, Sugimura S, Sato K, Mansour MM, Abd El-Aziz AH, Samir H, Islam MA, Bostami AR, Mandour AS, Elfadadny A, Ragab RF. Effects of selenium supplementation on rumen microbiota, rumen fermentation, and apparent nutrient digestibility of ruminant animals: A review. *Fermentation*. 2021 Dec 24;8(1):4.
19. Brask M, Lund P, Hellwing ALF, Poulsen M, Weisbjerg MR. Enteric methane production, digestibility and rumen fermentation in dairy cows fed different forages with and without rapeseed fat supplementation. *Anim Feed Sci Technol*. 2013;184(1–4):67–79. doi: 10.1016/j.anifeedsci.2013.06.006.
20. Wang Y, Xia K, Wang XN, Lin X, Liu J, Li YJ, Liu XL, Zhao WJ, Zhang YG, Guo JH. Improvement of feed intake, digestibility, plasma metabolites, and lactation performance of dairy cows fed mixed silage of sugar beet pulp and rice straw inoculated with lactic acid bacteria. *Journal of dairy science*. 2022 Jan 1;105(1):269-80.
21. Yang L, et al. Alterations in nutrient digestibility and performance of heat-stressed dairy cows by dietary L-theanine supplementation. 2022. Elsevier. doi: 10.1016/j.aninu.2022.08.002.
22. Mousa GA, Allak MA, Shehata MG, Hashem NM, Hassan OG. Dietary supplementation with a combination of fibrolytic enzymes and probiotics improves digestibility, growth performance, blood metabolites, and economics of fattening lambs. *Animals*. 2022 Feb 15;12(4):476.
23. Grossi S, Compiani R, Rossi L, Dell'Anno M, Castillo I, Sgoifo Rossi CA. Effect of slow-release urea administration on production performance, health status, diet digestibility, and environmental sustainability in lactating dairy cows. *Animals*. 2021 Aug 14;11(8):2405.
24. Hassanat F, Benchaar C. Corn silage-based diet supplemented with increasing amounts of linseed oil: Effects on methane production, rumen fermentation, nutrient digestibility, nitrogen utilization, and milk production of dairy cows. *Journal of Dairy Science*. 2021 May 1;104(5):5375-90.
25. Rehman A, et al. Nutrient digestibility, nitrogen excretion, and milk production of mid-lactation Jersey × Friesian cows fed diets containing different proportions of rumen-undegradable protein. 2020. *SciELO Brasil*. doi: 10.1590/0001-3765202020180787.
26. Woodford JA, Jorgensen NA, Barrington GP. Impact of dietary fiber and physical form on performance of lactating dairy cows. *Journal of dairy science*. 1986 Apr 1;69(4):1035-47.
27. Shakira G, Qubtia M, Ahmed I, Hasan F, Anjum MI, Imran M. Effect of indigenously isolated *saccharomyces cerevisiae* probiotics on milk production, nutrient digestibility, blood chemistry and fecal microbiota in lactating dairy cows. 2018. *thejaps.org.pk*. [Online]. Available: <http://www.thejaps.org.pk/docs/Accepted/2007/28-2/10.pdf>
28. Huhtanen P, Rinne M, Nousiainen J. A meta-analysis of feed digestion in dairy cows. 2. The effects of feeding level and diet composition on digestibility. *Journal of dairy science*. 2009 Oct 1;92(10):5031-42.
29. Bernabucci U, Lacetera N, Danieli PP, Bani P, Nardone A, Ronchi B. Influence of different periods of exposure to hot environment on rumen function and diet digestibility in sheep. *Int J Biometeorol*. 2009;53(5):387–395. doi: 10.1007/s00484-009-0223-6.
30. Moallem U, Lehrer H, Livshitz L, Zachut M, Yakoby S. The effects of live yeast supplementation to dairy cows during the hot season on production, feed efficiency, and digestibility. *Journal of dairy science*. 2009 Jan 1;92(1):343-51.

31. Sánchez NR, Ledin I. Effect of feeding different levels of foliage from *Cratylia argentea* to creole dairy cows on intake, digestibility, milk production and milk composition. *Trop Anim Health Prod.* 2006;38(4):343–351. doi: 10.1007/s11250-006-4314-7.
32. Sales J, Janssens GPJ. Acid-insoluble ash as a marker in digestibility studies: A review. 2003. *jafs.com.pl*. doi: 10.22358/jafs/67718/2003.
33. Chow LO, Baron VS, Corbett R, Oba M. Effects of planting date on fiber digestibility of whole-crop barley and productivity of lactating dairy cows. 2008;91(4). *library-archives.canada.ca*. doi: 10.3168/jds.2007-0854.
34. Morsy AS, Soltan YA, El-Zaiat HM, Alencar SM, Abdalla AL. Bee propolis extract as a phytogetic feed additive to enhance diet digestibility, rumen microbial biosynthesis, mitigating methane formation and health status of late pregnant ewes. *Anim Feed Sci Technol.* 2021;273:114834. doi: 10.1016/j.anifeedsci.2021.114834.
35. Zebeli Q, Tafaj M, Weber I, Dijkstra J, Steingass H, Drochner W. Effects of varying dietary forage particle size in two concentrate levels on chewing activity, ruminal mat characteristics, and passage in dairy cows. *J Dairy Sci.* 2007;90(4):1929–1942. doi: 10.3168/jds.2006-354.
36. Maulfair DD, Fustini M, Heinrichs AJ. Effect of varying total mixed ration particle size on rumen digesta and fecal particle size and digestibility in lactating dairy cows. *J Dairy Sci.* 2011 Jul;94(7):3527–3536. doi: 10.3168/jds.2010-3718.
37. Evangelista C, Petrocchi Jasinski F, Basiricò L, Turriziani G, Bernabucci U. Particle size distribution of total mixed rations fed to Italian Mediterranean buffaloes measured by the PSPS: Impact on milk quality and digestibility. *Ital J Anim Sci.* 2024;23(1):1535–1550. doi: 10.1080/1828051X.2024.2417714.
38. Miller-Cushon EK, DeVries TJ. Effect of dietary dry matter concentration on the sorting behavior of lactating dairy cows fed a total mixed ration. *J Dairy Sci.* 2009;92(7):3292–3298. doi: 10.3168/jds.2008-1772.
39. Abulaiti, A., Ahsan, U., Naseer, Z., Ahmed, Z., Liu, W., Ruan, C., Pang, X. and Wang, S., 2024. Effect of dietary Chinese herbal preparation on dry matter intake, milk yield and milk composition, serum biochemistry, hematological profile, and reproductive efficiency of Holstein dairy cows in early postpartum period. *Frontiers in Veterinary Science*, 11, p.1434548.
40. Ncisana L, Mabhaudhi T, Mkhize NR, Ravhuhali K, Tjelele TJ, Nyathi MK, Mbambalala L, Msiza NH, Nzeru MS, Modi AT. Water regimes in selected fodder radish (*Raphanus sativus*) genotypes: Effects on nutritional value and in vitro ruminal dry matter degradability. *Heliyon.* 2024 Apr 30;10(8).
41. Ahmad M, Jamil M, Ullah S, Ali M. Examine the effects of climate change on forage quality and its subsequent impact on milk yield and composition. *Indus J.* 2024. [Online]. Available: <http://induspublishers.com/IJBR/article/view/431>
42. Zhou L, et al. Effects of dietary concentrate: Forage ratio on development of gastrointestinal tract in black Tibetan sheep. *J Appl Anim Res.* 2022;50(1):192–197. doi: 10.1080/09712119.2022.2053131.
43. Yang WZ, Beauchemin KA. Increasing physically effective fiber content of dairy cow diets through forage proportion versus forage chop length: Chewing and ruminal pH. *J Dairy Sci.* 2009;92(4):1603–1615. doi: 10.3168/jds.2008-1379.
44. AlZahal O, Dionissopoulos L, Laarman AH, Walker N, McBride BW. Active dry *Saccharomyces cerevisiae* can alleviate the effect of subacute ruminal acidosis in lactating dairy cows. 2014. Elsevier. doi: 10.3168/jds.2014-8212.
45. Xu Q, Qiao Q, Gao Y, Hou J, Hu M, Du Y, Zhao K, Li X. Gut microbiota and their role in health and metabolic disease of dairy cow. *Frontiers in nutrition.* 2021 Aug 4;8:701511.
46. Chaucheyras-Durand F, Walker ND, Bach A. Effects of active dry yeasts on the rumen microbial ecosystem: Past, present and future. *Anim Feed Sci Technol.* 2008;145(1–4):5–26. doi: 10.1016/j.anifeedsci.2007.04.019.
47. Anadón A, Ares I, Martínez-Larrañaga MR, Martínez MA. Prebiotics and probiotics in feed and animal health. *Nutraceuticals Vet Med.* 2019;261–285. doi: 10.1007/978-3-030-04624-8_19.

-
48. Nova E, Gómez-Martínez S, González-Soltero R. The influence of dietary factors on the gut microbiota. 2022. mdpi.com. doi: 10.3390/microorganisms10071368.
 49. Wu S, Bhat ZF, Gounder RS, Mohamed Ahmed IA, Al-Juhaimi FY, Ding Y, Bekhit AE. Effect of dietary protein and processing on gut microbiota—A systematic review. *Nutrients*. 2022 Jan 20;14(3):453.
 50. Kumar Singh A, Cabral C, Kumar R, Ganguly R, Kumar Rana H, Gupta A, Rosaria Lauro M, Carbone C, Reis F, Pandey AK. Beneficial effects of dietary polyphenols on gut microbiota and strategies to improve delivery efficiency. *Nutrients*. 2019 Sep 13;11(9):2216.
 51. Uyeno Y, Shigemori S, Shimosato T. Effect of probiotics/prebiotics on cattle health and productivity. *Microbes Environ*. 2015;30(2):126–132. doi: 10.1264/jsme2.ME14176.
 52. Rojas OJ, Stein HH. Processing of ingredients and diets and effects on nutritional value for pigs. 2017. Springer. doi: 10.1186/s40104-017-0177-1.
 53. Wanapat M, Polyorach S, Boonnop K, Mapato C, Cherdthong A. Effects of treating rice straw with urea or urea and calcium hydroxide upon intake, digestibility, rumen fermentation and milk yield of dairy cows. *Livest Sci*. 2009;125(2–3):238–243. doi: 10.1016/j.livsci.2009.05.001.
 54. Dehghan-banadaky M, Corbett R, Oba M. Effects of barley grain processing on productivity of cattle. *Anim Feed Sci Technol*. 2007;137(1–2):1–24. doi: 10.1016/j.anifeedsci.2006.11.021.
 55. Zoabi H, Ammar H, Ghzayel S, Aziz BA, Kholif AE, Díaz A, de Haro-Martí M, Chahine M, López S. Feeding sodium hydroxide-treated almond hulls to assaf sheep: effects on chemical composition, nutrient digestibility, and zootechnical performance. *Agriculture*. 2025 May 5;15(9):1000.
 56. Giallongo F, Hristov AN, Oh J, Frederick T, Weeks H, Werner J, Lapierre H, Patton RA, Gehman A, Parys C. Effects of slow-release urea and rumen-protected methionine and histidine on performance of dairy cows. *Journal of dairy science*. 2015 May 1;98(5):3292–308.
 57. Varga GA, Kolver ES. Microbial and animal limitations to fiber digestion and utilization. 1997. Elsevier. doi: 10.1093/jn/127.5.819s.
 58. Yan T, Mayne CS, Porter MG. Effects of dietary and animal factors on methane production in dairy cows offered grass silage-based diets. *Int Congr Ser*. 2006;1293:123–126. doi: 10.1016/j.ics.2006.02.024.
 59. Piluzza G, Sulas L, Bullitta S. Tannins in forage plants and their role in animal husbandry and environmental sustainability: A review. *Grass Forage Sci*. 2014;69(1):32–48. doi: 10.1111/gfs.12053.
 60. Adesogan AT, Ma ZX, Romero JJ, Arriola KG. Ruminant Nutrition Symposium: Improving cell wall digestion and animal performance with fibrolytic enzymes. *J Anim Sci*. 2014;92(4):1317–1330. doi: 10.2527/jas.2013-7273.
 61. El-Zaiat HM, Al-Marzooqi W, Al-Kharousi K. Effects of chitosan-based additive on rumen fermentation and microbial community, nutrients digestibility and lactation performance in goats. *J Anim Physiol Anim Nutr (Berl)*. 2024. doi: 10.1111/jpn.14057.
 62. Ferraretto LF, Shaver RD, Bertics SJ. Effect of dietary supplementation with live-cell yeast at two dosages on lactation performance, ruminal fermentation, and total-tract nutrient digestibility in dairy cows. 2012. Elsevier. doi: 10.3168/jds.2011-5190.
 63. Bernabucci U, Bani P, Ronchi B, Lacetera N, Nardone A. Influence of short- and long-term exposure to a hot environment on rumen passage rate and diet digestibility by Friesian heifers. *J Dairy Sci*. 1999;82(5):967–973. doi: 10.3168/jds.S0022-0302(99)75316-6.