

Adverse Drug Reactions: A Global Health Challenge

Swarnima Pandey^{1,*}, Shivam Pathak², Ankit Verma², Sanjog Pal Singh²

Abstract

Adverse drug reaction (ADR) is an unfavorable event with a causal connection to the medication. It is an undesirable event occurred by a pharmacological substance. They are a major cause of Mortality all over, they are classified into augmented, chronic, bizarre, Delayed based on the severity of the damage they caused. The primary causes of adverse drug reactions (ADRs) are drug overdose, drug–drug interactions, genetic variations, and individual characteristics such as age, sex, and the presence of disease. Pharmacovigilance plays a key role in determination and prevention of ADRs. ADR monitoring and reporting is crucial because of the increasing number of ADR cases in our country. Timely reporting of ADR to regulatory authorities is important for drug monitoring, patient safety, and to identify the therapeutic range of medicines. In this review authors have covered different aspects of ADRs.

Keywords: Bizarre reactions, drug–drug interactions, primary care, patient monitoring, adverse drug reactions

INTRODUCTION

The following definitions are used in this article for clarity due to the variety of definitions found in world literature. “The word ‘adverse effect’ embraces all undesirable outcomes (including test findings suggesting harm without any clear clinical symptoms/signs); it removes the danger of misclassification, makes no assumptions about mechanism, and invokes no ambiguity” [1]. An undesirable outcome that has been linked to a drug, its usage, or its misuse is referred to as a “adverse drug effect” (ADE) [1].

A patient’s clinical reaction to a medication is known as a “adverse drug reaction” (ADR). “An substantially damaging or unpleasant reaction, arising from an intervention associated to the use of a medicinal product, which predicts hazard from future administration and necessitates prevention or specialized therapy, or change of the drug regimen, or withdrawal of the product” is the definition given here [2, 3]. Although considerable morbidity and occasional mortality are caused by adverse medication responses and bad effects, there is little focus on how to diagnose and treat adverse drug effects in an individual to prevent or lessen catastrophic harm.

A precise diagnosis is necessary for effective patient care. The body can react pathophysiologically

*Author for Correspondence

Swarnima Pandey
E-mail: Yesgoldi@gmail.com

¹Principal, Apex College of Pharmacy, Rampur, Uttar Pradesh, India

²Student, Department of Pharmacy, Apex College of Pharmacy, Rampur, Uttar Pradesh, India

Received Date: March 28, 2026

Accepted Date: April 02, 2026

Published Date: April 24, 2026

Citation: Swarnima Pandey, Shivam Pathak, Ankit Verma, Sanjog Pal Singh. Adverse Drug Reactions: A Global Health Challenge. Research & Reviews: A Journal of Pharmacology. 2026; 16(2): 1–6p.

to medications and other illness attacks in a restricted number of ways. This implies that negative consequences frequently pass for other illnesses. In some more uncommon cases, adverse drug reactions may be more frequently linked to exposure to drugs or chemicals than to other potential causes (such agranulocytosis or Stevens–Johnson syndrome), which partly simplifies the diagnostic process. When alternative causative probabilities are more likely than a drug and when it may be challenging to determine whether an individual has actually been exposed to a certain drug, at what dose, and when, it is considerably more difficult to make a diagnosis [4].

Adverse drug effects that the astute clinician must consider include therapeutic mistake, unanticipated failure of effect (perhaps due to a subpar or counterfeit product), drug misuse, accidental or suicidal self-administration, and homicidal drug use. There is growing evidence that we have overlooked the negative consequences associated with drug usage. They require a diagnosis for both the negative effects and the reasons behind them. To help identify strategies to stop harm in the future, we must look at the systemic sources of negative impacts in each unique case [5].

Approximately 70% of adverse drug reactions that result in hospital admissions are avoidable side effects of treatment, which are essentially medical blunders. Medical errors can be caused by a variety of factors, from blatant carelessness to blunders made by diligent medical personnel who are overworked, under pressure, or preoccupied. Tracing back to the ultimate causative variables is the only way to properly examine the complex ultimate causes. This is a lengthy procedure, but it may be crucial for identifying system faults that can be fixed and preventing future issues [6, 7]. Improved working hours to prevent fatigue and stress or double-checking prescription interpretation before dispensing to prevent errors due to misread handwriting are two simple examples of such a strategy [2].

Since the research is time-consuming and the underlying cause or causes of side effects may be complex, it does not play a significant role in clinical evaluation. However, when a pharmaceutical error is likely, the clinical personnel overseeing a patient is typically in the greatest position to at least initiate an assessment of the circumstances surrounding a major adverse impact. The reasons for systemic issues in the provision of healthcare, among other things, will be the subject of the subsequent study. 8,9 Instead of placing the blame on specific people, it is crucial to comprehend such systematic errors to identify strategies to prevent them.

Types of Adverse Drug Reactions

Drug reactions can be divided into a few categories based on how they happen and what they do. The common adverse drug reactions are Type A and Type B reactions. Type A reactions are predictable. Depending on the dose of the drug. Type B reactions are unusual. Do not depend on the dose of the drug [7, 8].

Both Type A and Type B reactions are influenced by genetics. This means that some people are more likely to have drug reactions because of their genes. For example, genetic testing can help prevent drug reactions when people take certain drugs like thiopurines and warfarin.

Adverse drug reactions can also be categorized by what they do to the body. They can affect how drugs are absorbed how they work and how they interact with drugs. Understanding the kinds of adverse drug reactions is important for keeping people safe when they take drugs.

With computers and algorithms to help identify and categorize adverse drug reactions we still need experts to study them and make sure we understand what is happening. Many things can contribute to drug reactions, including genetics, how drugs are broken down in the body and how they interact with other drugs [9, 10].

Some common drugs that can cause drug reactions are narcotics, diuretics, and anticoagulants. These are drugs that people take every day and they can still cause problems. The idea that new or poorly tested drugs can cause serious problems is not true.

What is also interesting is that some drug combinations can increase the risk of drug reactions. This is because some drugs do not work together. To make drugs safer we need to understand what causes drug reactions. This includes things like genetics how drugs interact with each other and how they are broken down in the body.

Adverse drug reactions can be serious and even life-threatening. They can cause problems with the system, stomach, skin and other parts of the body. Sometimes people must stay in the hospital for a time because of adverse drug reactions.

Signs and symptoms of drug reactions can be different for different people. Some common symptoms include changes to the system and stomach problems. Skin problems are also common. In some cases, adverse drug reactions can be fatal.

SIGNS OF DRUG ALLERGY

- Skin that is red, flaky, itching or swollen.
- A red flat spot on your skin with tiny pimples.
- Bumps, on the skin that look like bees.
- Peeling skin.
- Vision problems.
- Itching.
- Eye watering.
- Skin rash.
- Hives.
- Runny nose.
- Swelling of the skin.

Anaphylaxis is a sign of a medication allergy. It can be fatal if not treated away. Some severe symptoms include:

- Trouble swallowing.
- Trouble breathing.
- Feeling dizzy or lightheaded.
- Low blood pressure.
- Fast heart rate.
- Feeling anxious or confused.

We need to be careful when taking drugs and watch for signs of drug reactions. This includes understanding what causes them and how to prevent them. By doing this we can make drugs safer. Help people stay healthy, exceptional AUROC score of 0.917. It embeds items, such as medicines, ADRs, and proteins, using the Node2Vec algorithm. An ensemble model that combines decision trees, random forests, support vector machines, and adaptive boosting is another successful strategy; on benchmark datasets, it obtained an F measure of 89%. Successful use of deep learning techniques has also been documented; one study found that 14 models that predicted for different ADRs had an average validation precision of 89.4%. Furthermore, even before they are formally published, web search log analysis has demonstrated promise for early ADR detection. Interestingly, 65.8% of ADR-positive admissions have been found by automated laboratory data analysis employing automatic laboratory signals (ALS), which may increase the probability of identification of undiscovered ADRs in clinical settings. To sum up, AI solutions for ADR detection cover a broad spectrum of methods, including data mining, deep neural networks, and machine learning algorithms.

WHY MONITOR ADVERSE DRUG REACTIONS IN INDIA

It is seen that medicine which works well for patients may not work the same way for people from different backgrounds. This is because the genes of patients play a role in how they respond to medicines. Genes can affect how the body processes medicines. This can change how well the medicine works or how safe it is. As a result, people may have different reactions to the same medicine, which can make it hard to develop new medicines. For example, a medicine called pioglitazone is not allowed in some countries because it may increase the risk of bladder cancer. It is still available in India, where bladder cancer is less common. So even if a medicine is approved in one country it is crucial to test it on populations and monitor its safety closely.

Pharmacovigilance Program in India

The All India Institute of Medical Sciences in New Delhi was chosen to lead the Pharmacovigilance Program of India in 2010. The goal of this program is to keep people in India safe by making sure the

benefits of medicines outweigh the risks. In Delhi, in 2024, a training program was organized to teach people about pharmacovigilance and patient safety.

Aims of Pharmacovigilance Program in India

- To train people to develop skills in monitoring medicine safety.
- To build capacity in medicine safety.
- The program wants to improve safety and welfare in India by monitoring drug safety and reducing risks associated with medicines.
- The Indian government started this program on July 14, 2010, to protect health.
- The India Pharmacopoeia Commission also works to make sure that the benefits of using medicines are greater, than the risks.
- To make people aware and understand the importance of reporting Adverse Events and Adverse Drug Reactions.
- To encourage research in Pharmacovigilance.

Methods to Monitor Drug Safety

Pharmacovigilance Methods are used to monitor drug safety. There are ways to do this.

One way is called Passive Surveillance. This is when people report problems with drugs on their own. This can be done by doctors, hospitals or patients. The reports go from areas to state, then to the country and finally to the world.

Another way is called Active Surveillance. This is when we carefully watch people who are taking drugs to see if they have any problems.

We use this method to learn more about the problems that can happen with drugs how often they happen. What causes them? However, this method can be difficult to use because it needs a lot of resources. So, we only use it for drugs and in certain places.

Active surveillance can also be done in the community like when we watch a group of people who are taking a drug. We also do studies to learn more about drug safety. These studies can be done to see if a certain problem is caused by a drug. We can do these studies by looking at what happened to people in the past or by watching what happened to people in the future.

ENVIRONMENTAL PHARMACOVIGILANCE

An Indian scientist suggested that we need to keep an eye on the effects of medicines on plants and animals after people or animals take them. This is different from keeping an eye on the effects of medicines on people and animals. When we take medicines, our bodies get rid of them. They can end up in the environment. If we do a job of checking the risks of medicines before we start testing them on people, we can reduce the harm they can cause to the environment.

At first this field of study was called Pharmacoenvironmentology. Now it is also known as EcoPharmacovigilance, PharmEcovigilance and PharmaEcostewardship.

The Journal of Pharmacovigilance and Drug Safety wrote about Pharmacovigilance in India. They said that different animals can react differently to the same medicine. Within the same species it can be hard to understand how different amounts of a drug affect different animals.

This is because animals of ages and genders can react differently to the same drug. This makes it challenging to keep animals safe when we give them medicine. There is also something called Environmental Pharmacovigilance. This is when we watch to see if drugs hurt the environment. We do this because when we take medicine our bodies get rid of the drugs and they can go into the air, water and soil. If we are careful and check the drugs before we use them, we can reduce the harm they can cause to the environment. This is an area of study that is important for keeping our planet safe.

CHALLENGES WITH PHARMACOVIGILANCE IN INDIA

The Journal of Pharmacovigilance and Drug Safety wrote about these challenges. They said that different animals can react differently to the same medicine. Within the same species it can be hard to understand how different amounts of a drug affect different animals. This is because animals of ages and genders can react differently to the same drug. This makes it challenging to keep animals safe when we give them medicine. It is also a challenge for human Pharmacovigilance [8, 11].

The problem with Pharmacovigilance and Post-Marketing Surveillance in India is that many bad reactions to medicines are not reported. This happens because doctors and other medical workers do not know enough about medicines and there are not people trained to keep an eye on the bad effects of medicines. Also, many people in India do not know much about Pharmacovigilance. Other challenges include equipment, a long time between when rules are made and when they are put into action and not many checks to make sure medicines are safe. Sometimes doctors do not even realize that a patient's problems are caused by a reaction to a medicine and this can be very serious. Patients may also not tell their doctors about all the medicines they are taking including those from types of medicine.

The cost of reactions to medicines is very high. In India new medicines are often sold without being tested for safety and this can lead to more people having bad reactions. In the past companies, in India just looked at how people reacted to medicines over a time to see if they were safe. It is also hard to keep an eye on the effects of traditional medicines because each type of traditional medicine is different and uses different methods [6, 7].

Multiple Evidence Fusion

Multiple evidence fusion is a good way to predict bad reactions to drugs. This method can handle both new drugs. It works by combining lots of information about drugs and bad reactions. This includes information about how drugs connected to bad reactions. Multiple evidence fusion is better than methods that only use one or a few types of information. It is very good at finding bad reactions. For example, people have used information on the internet to find reactions. They looked at what people were searching for online. Used that information to find patterns. This can help us find reactions before they happen [8, 11].

Some other methods are also very good at finding reactions. One method uses a combination of techniques like decision trees and random forests. This method is very good at finding reactions with an accuracy of 89%. So, we are getting better at using computers to find reactions. We are using lots of information and methods to do this. Multiple evidence fusion is a way to do this, but other methods are also useful. These methods can help us find reactions early which can keep patients safe and reduce medical costs.

MANAGEMENT OF ADR

Reactions to drugs are a big problem worldwide. We need to be good at managing them to keep patients safe. Pharmacists play a role in tracking and reporting bad reactions. Lots of bad reactions are not reported. In fact, up to 98% of reactions may not be reported.

This is a problem because we need to know about reactions to keep patients safe. Different countries have ways of reporting bad reactions. For example, in Saudi Arabia not many pharmacists report reactions. In Australia pharmacists are very good, at reporting bad reactions.

We need to find ways to get more people to report reactions. One way to do this is to make it easier for people to report them. We can also use computers to help us track reactions. Some hospitals have used education and reminders to get more people to report reactions.

In the end we need to work to get better at reporting and managing bad reactions. This includes pharmacists, doctors and other healthcare professionals. We all need to be aware of the importance of reporting reactions and work together to keep patients safe [10, 12].

CONCLUSION

Numerous clinical and demographic characteristics have a substantial impact on the incidence of adverse drug reactions (ADRs); middle-aged adults, women, people receiving intravenous medication, and people taking anti-tumor medications are at higher risk. Enhancing the effectiveness of monitoring procedures and creating individualized treatment plans require an understanding of the intricate interactions between these variables in severe adverse drug reactions. In the long run, this strategy may enhance clinical results and patient safety.

In summary, safe and excellent healthcare depends on the efficient treatment and prevention of adverse drug reactions (ADRs). Understanding the categorization and identification procedures as well as the variety of ADRs, from mild to severe, are crucial for prompt intervention. To lessen the effects of adverse reactions, early identification, timely treatment, and preventative measures including desensitization and premedication regimens are essential. Additionally, a proactive approach to prevention includes patient education, effective communication between healthcare practitioners, comprehensive patient evaluation, and the use of technology like electronic health records. The categorization of adverse drug reactions (ADRs) into discrete kinds facilitates comprehension of their attributes and guides suitable therapies. Participating actively in pharmacovigilance programs and reporting systems also encourages a team effort to continuously evaluate medication safety and spot new concerns. To promote a culture of safety and provide the best possible patient care, cooperation between medical experts, regulatory bodies, and patients is essential.

Acknowledgment

Authors are sincerely grateful To Dr. Mandeep Singh Managing Director Apex Group of Institutions Rampur for his motivation to publish this article.

REFERENCES

1. Jeffrey KA, Robin EF. Clarification of terminology in drug safety. *Drug Saf.* 2005;28:851–870.
2. U.S. Food and Drug Administration (FDA). Guidance for industry: Good pharmacovigilance practices and pharmacoepidemiologic assessment. Rockville (MD): Center for Drug Evaluation and Research and Center for Biologics Evaluation and Research; 2005 Mar.
3. Abubakar AR, Simbak NB, Haque M. Adverse drug reactions: Predisposing factors, modern classifications and causality assessment. *Res J Pharm.* 2014;7(9):1091–1098.
4. Edwards IR, Aronson JK. Adverse drug reactions: Definitions, diagnosis, and management. *Lancet.* 2000;255–259.
5. Dhikav V, Singh S, Anand KS. Adverse drug reaction monitoring in India. *JIACM.* 2004;6(1):27–33.
6. Pallasch TJ. Principles of pharmacotherapy V: Toxicology and adverse drug reactions. *Anesth Prog.* 1989;36:41–45.
7. Phillips KA, Veenstra DL, Oren E. Potential role of pharmacogenomics in reducing adverse drug reactions: A systematic review. *JAMA.* 2001;4:2270–2279.
8. Lavan AH, Gallagher P. Predicting risk of adverse drug reactions in older adults. *Ther Adv Drug Saf.* 2016;7(1):11.
9. Whittle C, Rogers W. Clinical pharmacy and therapeutics. Edinburgh: Churchill Livingstone Elsevier; 2007:51–60.
10. Ragoonanan D, Khazal S, Abdel-Azim H, McCall D, Cuglievan B, Tambaro FP, et al. Diagnosis, grading and management of toxicities from immunotherapies in children, adolescents and young adults with cancer. *Nat Rev Clin Oncol.* 2021;18(7):435–453. doi: 10.1038/s41571-021-00474-4.
11. Tripathi KD. Textbook of pharmacology. New Delhi: Jaypee Brothers Medical Publishers; 2013:218–222.
12. Pitts P, Louet HL, Moride Y, Conti RM. 21st century pharmacovigilance: Efforts, roles, and responsibilities. *Lancet Oncol.* 2016;17(11):e486–e492. doi: 10.1016/S1470-2045(16)30312-6.