

Association Between Smartphone Addiction and Functional Exercise Capacity in College Students

Manmeet Kaur Bhalla^{1,*}, Niraj Kumar², Deptee Warikoo³

Abstract

Smartphone addiction has become increasingly prevalent among college students and is associated with various negative physical and psychological outcomes. Functional exercise capacity, a key indicator of physical fitness and health, may be adversely affected by excessive smartphone use due to sedentary behavior and reduced physical activity. This article examines the association between smartphone addiction and functional exercise capacity in college students by analyzing behavioral, psychological, physiological, and methodological perspectives. Evidence suggests that higher levels of smartphone addiction are linked to decreased physical activity, poor cardiovascular endurance, and reduced functional performance. Psychological factors such as stress, anxiety, depression, and sleep disturbances further mediate this relationship. Intervention strategies, including structured exercise programs, behavioral modification techniques, digital detox interventions, and wearable technology, have shown promise in improving both smartphone use patterns and exercise capacity. The findings highlight the need for integrated health promotion strategies targeting both digital behavior and physical fitness among college populations.

Keywords: Smartphone addiction, functional exercise capacity, college students, physical activity, sedentary behavior, mental health, exercise tolerance, digital health

INTRODUCTION

Smartphone addiction has emerged as a major behavioral concern among college students owing to their increased reliance on mobile devices for communication, education, and entertainment, with recent studies highlighting a rapid escalation in daily screen time and dependency patterns among young adults. The integration of smartphones into nearly every aspect of academic and social life has transformed them from simple communication tools into indispensable devices that significantly shape behavior, cognition, and lifestyle habits. The widespread availability of high-speed internet, social media platforms, and mobile applications has further intensified smartphone usage, making it difficult for students to regulate their engagement and maintain healthy usage patterns. Consequently, excessive

usage patterns characterized by compulsive checking, prolonged screen exposure, and reduced behavioral control have become increasingly common among college populations. These behaviors are often reinforced by instant gratification mechanisms, such as notifications, likes, and continuous content streams, which contribute to the development of addictive tendencies. Over time, such patterns can lead to psychological dependence, withdrawal symptoms, and a diminished capacity to disengage from digital environments, even when necessary. The impact of smartphone addiction extends beyond mental health and cognitive functioning, significantly affecting physical health outcomes, including reduced physical activity and increased sedentary behavior.

*Author for Correspondence

Manmeet Kaur Bhalla

E-mail: healtheducation236@gmail.com

¹Ph.D. Scholar, Department of Physiotherapy, Shri Guru Ram Rai University, Dehradun, Uttarakhand, India

²Professor and Head of Department, Department of Physiotherapy, Shri Guru Ram Rai University, Dehradun, Uttarakhand, India

³Ph.D. Scholar, Department of Physiotherapy, Shri Guru Ram Rai University, Dehradun, Uttarakhand, India

Received Date: April 06, 2026

Accepted Date: May 07, 2026

Published Date: May 16, 2026

Citation: Manmeet Kaur Bhalla, Niraj Kumar, Deptee Warikoo. Association Between Smartphone Addiction and Functional Exercise Capacity in College Students. International Journal of Orthopedic Nursing and Practices. 2026; 4(1): 28–51p.




Figure 1. Global prevalence trends of smartphone addiction among college students.

Functional exercise capacity, defined as the ability to perform sustained physical activity efficiently, is a critical indicator of overall health and fitness in young adults and is influenced by cardiovascular endurance, muscular strength, and respiratory efficiency [1]. In college students, maintaining optimal functional exercise capacity is essential for supporting academic performance, mental well-being, and long-term health outcomes. However, the increasing prevalence of sedentary lifestyles driven by excessive smartphone use has raised concerns regarding declining levels of physical fitness in this population. Prolonged screen time often replaces time that could otherwise be spent engaging in physical activities such as sports, exercise, or recreational movement, leading to reduced energy expenditure and lower fitness levels [2]. Additionally, smartphone use is frequently associated with passive behaviors such as sitting or lying down, which further contribute to decreased physical activity and increased risk of metabolic disorders. The coexistence of high smartphone use and declining physical activity levels has therefore become a critical area of concern for researchers and health professionals seeking to understand the broader implications of digital behavior on physical health. Emerging evidence suggests that individuals with higher levels of smartphone addiction demonstrate significantly lower functional exercise capacity compared to those with moderate or controlled usage patterns [3]. This association is influenced by multiple interacting factors, including reduced motivation for exercise, increased fatigue, and disrupted sleep patterns resulting from excessive screen exposure. Sleep disturbances, particularly those caused by late-night smartphone use and blue light exposure, negatively impact recovery processes and reduce overall physical performance [4]. Furthermore, the habitual use of smartphones in static postures contributes to musculoskeletal strain and discomfort,

which may further discourage participation in physical activity. The psychological dimension of smartphone addiction also plays a significant role, as increased levels of stress, anxiety, and depression are commonly observed among individuals with problematic smartphone use. These psychological factors not only reinforce addictive behaviors but also reduce engagement in health-promoting activities such as exercise and outdoor recreation [5]. Consequently, a negative cycle is established in which smartphone addiction leads to reduced physical activity, which in turn contributes to declining functional exercise capacity and overall health. Figure 1 illustrates the global rise in smartphone addiction among college students over the past decade, showing a consistent upward trend that reflects increasing digital dependence and its relevance to modern student lifestyles. This trend underscores the urgency of addressing smartphone addiction as a multidimensional health issue that requires comprehensive intervention strategies targeting both behavioral and physical health domains. Understanding the relationship between smartphone addiction and functional exercise capacity is therefore essential for developing effective prevention and intervention programs aimed at improving the health and well-being of college students in the digital age [6].

BACKGROUND AND THEORETICAL FRAMEWORK

Smartphone addiction is conceptualized as a behavioral addiction involving impaired control over usage, development of tolerance, and the presence of withdrawal symptoms when access to the device is limited, reflecting patterns similar to other non-substance-related addictions observed in young adults. This form of addiction is increasingly recognized within the framework of behavioral sciences as it interferes with daily functioning, academic responsibilities, and social interactions among college students [7]. The compulsive nature of smartphone use is driven by repeated engagement with rewarding digital content, reinforcing habitual checking and prolonged screen exposure over time. Neurobehavioral evidence suggests that such repetitive behaviors activate reward pathways in the brain, particularly those associated with dopamine release, thereby strengthening dependence and habitual use. The dual process theory provides a useful explanation for this phenomenon by highlighting the imbalance between the impulsive system, which seeks immediate gratification, and the reflective system, which regulates self-control and decision making. In individuals with smartphone addiction, the impulsive system becomes dominant, leading to excessive engagement with digital stimuli despite awareness of negative consequences [8]. This imbalance reduces the ability to regulate usage patterns and contributes to the persistence of addictive behaviors even in the presence of academic or health-related concerns. Functional exercise capacity, on the other hand, is influenced by the integrated functioning of cardiovascular, respiratory, and musculoskeletal systems, which collectively determine an individual's ability to perform sustained physical activity. Adequate cardiovascular endurance ensures efficient oxygen delivery to working muscles, while respiratory efficiency supports optimal gas exchange during physical exertion. Musculoskeletal strength and flexibility further contribute to the ability to perform physical tasks effectively and without fatigue. In college students, maintaining functional exercise capacity is essential for supporting both physical health and cognitive performance [9]. However, reduced physical activity resulting from excessive smartphone use has emerged as a critical factor linking behavioral addiction to declining physical fitness. Prolonged engagement with smartphones often leads to sedentary behavior, characterized by extended periods of sitting or lying down with minimal energy expenditure. This sedentary lifestyle reduces opportunities for engaging in moderate to vigorous physical activity, which is necessary for maintaining cardiovascular and muscular fitness [10]. Over time, decreased physical activity contributes to deconditioning of physiological systems, resulting in reduced endurance, lower exercise tolerance, and impaired functional capacity. Additionally, excessive smartphone use is frequently associated with poor posture and limited movement, which further exacerbates musculoskeletal strain and reduces overall physical efficiency [11]. The interaction between behavioral addiction and physical inactivity creates a complex cycle in which increased smartphone use leads to reduced activity levels, which in turn contribute to declining physical health and further reinforce sedentary habits. Psychological factors such as stress, anxiety, and reduced motivation also play a mediating role by influencing both smartphone usage patterns and engagement in physical activity. Individuals experiencing higher levels of psychological distress may rely more heavily on smartphones as a coping mechanism, thereby increasing screen time and reducing participation in physical exercise [12].




Figure 2. Conceptual framework linking smartphone addiction, sedentary behavior, and functional exercise capacity.

This interaction highlights the multidimensional nature of the relationship between smartphone addiction and functional exercise capacity, which involves behavioral, psychological, and physiological components. Figure 2 presents a conceptual framework demonstrating how smartphone addiction leads to sedentary behavior and reduced exercise capacity through multiple interacting pathways, integrating theoretical and empirical insights to provide a comprehensive understanding of this relationship. This framework emphasizes the central role of reduced physical activity as a key mediator linking digital behavior to physical health outcomes while acknowledging the influence of psychological and environmental factors. Understanding these interconnected pathways is essential for developing targeted interventions aimed at restoring the balance between digital engagement and physical activity in college students [13].

OPERATIONAL DEFINITIONS

Smartphone addiction refers to excessive and uncontrolled smartphone use characterized by behavioral dependence, inability to regulate usage, withdrawal symptoms, and significant interference with daily academic, social, and physical activities among college students. This condition has gained increasing recognition as a form of behavioral addiction due to its similarity with substance use disorders in terms of compulsive engagement and negative consequences. Individuals experiencing smartphone addiction often demonstrate persistent checking behaviors, craving for digital interaction, and difficulty in reducing usage despite awareness of its harmful effects. These patterns are reinforced by psychological gratification derived from social connectivity, entertainment, and instant access to information through digital platforms. Withdrawal symptoms such as irritability, restlessness, and anxiety are commonly reported when access to smartphones is restricted, further indicating the addictive nature of this behavior [14]. Over time, such excessive usage disrupts daily routines, impairs

concentration, and reduces productivity, particularly in academic environments. Functional exercise capacity refers to the ability of an individual to perform physical tasks efficiently and sustain activity over a period of time, reflecting overall physical fitness and endurance. It is a comprehensive measure influenced by cardiovascular, respiratory, and musculoskeletal systems working in coordination to support physical performance. Common methods used to assess functional exercise capacity include endurance-based tests such as the six-minute walk test, shuttle walk test, and maximal oxygen uptake measurements. These assessments provide valuable insights into an individual's aerobic capacity, exercise tolerance, and ability to perform daily physical activities. In young adults, particularly college students, maintaining optimal functional exercise capacity is essential for supporting both physical health and mental well-being [15]. However, lifestyle changes associated with increased digital engagement have contributed to declining physical activity levels and reduced exercise performance in this population. Sedentary behavior refers to activities that involve low energy expenditure, typically performed in sitting or reclining positions, such as watching screens, browsing the internet, or engaging in prolonged smartphone use. This type of behavior has become increasingly prevalent with the widespread use of digital devices and is considered a major risk factor for various health conditions. Prolonged sedentary time is associated with reduced metabolic activity, decreased cardiovascular fitness, and increased risk of obesity and related disorders. In the context of smartphone addiction, sedentary behavior often replaces time that could otherwise be allocated to physical activity, thereby contributing to reduced energy expenditure and declining fitness levels. The interaction between smartphone addiction and sedentary behavior creates a reinforcing cycle in which increased screen time leads to decreased physical movement, further promoting inactivity and physical deconditioning. Additionally, sedentary behavior is linked to poor posture and musculoskeletal discomfort, particularly in the neck, shoulders, and lower back, which can further limit physical activity participation. Psychological factors such as stress and lack of motivation also contribute to increased sedentary behavior and reduced engagement in exercise. The combined effect of smartphone addiction and sedentary lifestyle significantly impacts functional exercise capacity by reducing cardiovascular endurance, muscular strength, and overall physical performance. Understanding these operational definitions is essential for establishing a clear conceptual framework when examining the relationship between digital behavior and physical health outcomes in college students. Clear definitions also enable researchers to standardize measurement approaches and improve the comparability of findings across studies. Furthermore, recognizing the interconnected nature of smartphone addiction, sedentary behavior, and functional exercise capacity provides a foundation for developing targeted interventions aimed at promoting healthier lifestyles and reducing the negative impact of excessive digital engagement [16–20].

METHODOLOGY

This article is based on a narrative review of literature published within the last ten years, aiming to synthesize existing evidence on the association between smartphone addiction and functional exercise capacity among college students through a structured yet flexible approach. A narrative review methodology was selected to allow comprehensive integration of findings from diverse study designs, including cross-sectional, observational, and experimental research, thereby providing a broad understanding of the topic. Relevant studies were identified using major electronic databases such as PubMed, Scopus, Web of Science, and Google Scholar, which are widely recognized for indexing high-quality peer-reviewed research in health sciences and behavioral studies. A systematic search strategy was employed using combinations of keywords, including smartphone addiction, mobile phone dependence, physical activity, exercise capacity, sedentary behavior, and college students, to ensure comprehensive retrieval of relevant literature. Boolean operators such as AND and OR were used to refine the search and improve the accuracy of results by linking related concepts and filtering irrelevant studies. The search was limited to articles published in English within the last decade to ensure the inclusion of recent and relevant findings reflecting current technological and behavioral trends. Inclusion criteria were defined prior to the search and included studies focusing on college or university students, studies assessing smartphone addiction or problematic smartphone use, and studies reporting outcomes related to physical activity or functional exercise capacity.

Table 1. Prevalence and associated risk factors of smartphone addiction among college students.

Study/region	Prevalence (%)	Population characteristics	Key risk factors identified
Asia (multi-country)	30–50	Undergraduate students	High academic stress. social media overuse. poor sleep quality
India	35–60	College and university students	Excessive screen time. lack of physical activity. peer influence
China	25–45	Medical and non-medical students	Anxiety. depression. academic pressure
Europe	20–40	University students	Sedentary lifestyle. low physical activity. internet dependency
Middle East	40–70	Health science students	Emotional stress. sleep disturbances. high social media usage
United States	25–50	College students	Behavioral habits. poor time management. digital engagement
Korea	35–55	University students	Gaming addiction. psychological distress. academic burden
Global average	30–60	Mixed student populations	Sedentary behavior. poor lifestyle habits. lack of awareness

Exclusion criteria involved studies conducted on clinical populations, studies focusing on children or older adults, and articles lacking sufficient methodological rigor or full-text availability. After initial identification, titles and abstracts were screened to determine relevance, followed by full-text review of selected articles to confirm eligibility based on predefined criteria. Data extraction was performed systematically by reviewing selected studies and identifying key variables such as sample characteristics, measurement tools, study design, and major findings related to smartphone use and physical fitness. The extracted data were then organized into thematic categories to facilitate qualitative synthesis and interpretation of relationships between variables. This qualitative synthesis approach enabled the identification of consistent patterns, trends, and gaps in literature without restricting analysis to statistical aggregation. The methodology also emphasized comparison across studies to highlight similarities and differences in findings based on geographical, cultural, and demographic variations. To ensure reliability, preference was given to studies employing validated measurement tools such as the Smartphone Addiction Scale and International Physical Activity Questionnaire. The methodological framework also considered the quality of included studies by evaluating factors such as sample size, study design, and clarity of outcome measures. Table 1 summarizes the prevalence of smartphone addiction across different populations and highlights key contributing factors such as gender differences, academic stress, and social media usage patterns, thereby providing a comprehensive overview of the scope of the issue. Table 1 also facilitates comparison of findings across regions and helps identify populations at higher risk of developing problematic smartphone use. The narrative review approach allows for integration of multidisciplinary perspectives, including physiological, psychological, and behavioral dimensions of smartphone addiction and exercise capacity. Additionally, this methodology supports the identification of research gaps, particularly in relation to longitudinal and intervention-based studies, which are currently limited in literature. By synthesizing evidence from diverse sources, the review provides a holistic understanding of how smartphone addiction influences physical health outcomes in college students. Furthermore, the methodology acknowledges potential limitations such as publication bias, language restrictions, and reliance on self-reported data in many studies, which may affect the generalizability of findings. Despite these limitations, the structured approach to literature selection and analysis enhances the credibility and relevance of the findings presented in this article.

MEASUREMENT TOOLS AND INSTRUMENTS

Standardized tools such as the Smartphone Addiction Scale and International Physical Activity Questionnaire are widely used for assessing behavioral and physical health variables in research related to smartphone addiction and functional exercise capacity among college students, ensuring consistency and comparability across studies.

The Smartphone Addiction Scale is one of the most commonly used instruments and evaluates multiple dimensions of addictive behavior, including daily life disturbance, withdrawal, tolerance, and overuse, thereby providing a comprehensive assessment of problematic smartphone use. Short versions of this scale have also been developed to facilitate easier administration while maintaining reliability and validity in large population studies. The problematic mobile phone use questionnaire is another widely used tool that focuses on behavioral dependence and psychosocial consequences associated with excessive smartphone use. These tools are often self-reported and rely on Likert scale responses, allowing researchers to quantify subjective experiences related to smartphone use. In addition to addiction-specific scales, broader psychological assessment tools are sometimes used to evaluate associated factors such as stress, anxiety, and depression, which are closely linked to smartphone usage patterns. Physical activity levels are commonly measured using the International Physical Activity Questionnaire, which assesses frequency, duration, and intensity of physical activities performed over a specified period. This questionnaire categorizes activity into different domains such as walking, moderate activity, and vigorous activity, providing a detailed overview of an individual's physical activity profile. The international physical activity questionnaire (IPAQ) is widely used due to its simplicity, cost-effectiveness, and ability to capture data across diverse populations and settings. Objective measures such as accelerometers and pedometers are also sometimes used to complement self-reported data and provide more accurate estimates of physical activity and sedentary behavior. Functional exercise capacity is measured using performance-based tests that evaluate an individual's ability to sustain physical activity over time, reflecting real-world functional ability. The six-minute walk test is one of the most commonly used assessments and measures the distance an individual can walk on a flat surface within six minutes, serving as an indicator of aerobic endurance and cardiovascular fitness. This test is simple, non-invasive, and widely applicable across different populations, making it a preferred choice in both clinical and research settings. Another important measure of functional exercise capacity is maximal oxygen uptake or $\dot{V}O_2$ max, which represents the maximum amount of oxygen the body can utilize during intense exercise and is considered the gold standard for assessing cardiorespiratory fitness. $\dot{V}O_2$ max is typically measured using graded exercise tests conducted on treadmills or cycle ergometers under controlled conditions. Shuttle run tests and step tests are also used as alternative methods for assessing aerobic capacity and endurance, particularly in field-based studies. These tests provide valuable information about an individual's physical performance and ability to engage in sustained activity. Figure 3 demonstrates commonly used measurement tools and their domains, illustrating how smartphone addiction is assessed through behavioral and psychological indicators while functional exercise capacity is evaluated through physiological and performance-based measures. The integration of subjective and objective assessment tools enhances the accuracy and reliability of research findings by capturing multiple dimensions of the variables under study. Additionally, combining self-reported questionnaires with performance-based tests allows researchers to explore the relationship between perceived behavior and actual physical capacity. The use of validated measurement tools also ensures standardization across studies, enabling meaningful comparison of results and synthesis of evidence in systematic reviews and meta-analyses. However, it is important to acknowledge limitations such as response bias in self-reported questionnaires and variability in test administration, which may affect the precision of measurements. Despite these challenges, the use of established assessment tools remains essential for advancing research on smartphone addiction and its impact on physical health outcomes.

EPIDEMIOLOGY OF SMARTPHONE ADDICTION IN COLLEGE STUDENTS

The prevalence of smartphone addiction among college students ranges between 20% and 45% globally, reflecting a growing public health concern that varies across regions and populations depending on technological access and lifestyle patterns. This wide range in prevalence is influenced by differences in assessment tools, cultural norms, and patterns of smartphone use among young adults in different countries. In many developed and developing nations, college students represent one of the highest risk groups for smartphone addiction because of their constant engagement with digital platforms for both academic and social purposes. Academic demands play a significant role in increasing smartphone usage, as students rely heavily on mobile devices to access study materials, attend online classes, and communicate with peers and faculty.




Figure 3. Measurement tools used for assessing smartphone addiction and functional exercise capacity.

The integration of digital learning environments has further blurred the boundary between productive and excessive smartphone use, making it difficult to distinguish between necessary and addictive behaviors. Social networking is another major contributing factor, as platforms offering instant messaging, content sharing, and social validation encourage prolonged engagement and repeated checking behaviors. The need for social connectivity and fear of missing out often drives students to remain constantly connected, thereby increasing the risk of developing addictive patterns. Accessibility also plays a crucial role, as the widespread availability of affordable smartphones and high-speed internet has made continuous online engagement possible across diverse socioeconomic groups. In regions with higher technological penetration, prevalence rates tend to be higher due to greater exposure and ease of access to digital content. Conversely, variations in cultural attitudes toward technology use and lifestyle habits can influence the extent of smartphone dependence among students. For instance, societies that emphasize outdoor activities and physical engagement may exhibit lower levels of addiction compared to those with predominantly sedentary lifestyles. Gender differences have also been observed, with some studies indicating higher addiction rates among females due to increased social media use, while others report higher rates among males due to gaming and entertainment-related activities. Psychological factors such as stress, anxiety, and depression further contribute to smartphone addiction by increasing reliance on digital devices as coping mechanisms. Students experiencing higher levels of academic or emotional stress are more likely to engage in excessive smartphone use, which in turn reinforces addictive behavior. Environmental factors such as hostel living, peer influence, and lack of structured routines also contribute to increased screen time among college students. Figure 4 illustrates the distribution of smartphone addiction across different regions, highlighting variations influenced by sociocultural and technological factors, and demonstrating how prevalence rates differ between urban and rural populations as well as between developed and developing countries. Figure 4 also emphasizes the role of digital infrastructure and internet penetration in shaping usage patterns and addiction risk. Regional differences in educational systems and lifestyle habits further contribute to variations in smartphone use, with some regions reporting higher dependence due to greater integration of technology in daily life. Additionally, policy frameworks and awareness programs related to digital health can influence the prevalence of smartphone addiction by promoting responsible usage and reducing excessive screen time.




Figure 4. Regional distribution of smartphone addiction prevalence among college students.

Despite these variations, the overall trend indicates a steady increase in smartphone addiction among college students worldwide, highlighting the need for targeted interventions and preventive strategies to address this issue. The growing prevalence also underscores the importance of understanding the underlying factors contributing to this issue to develop effective solutions. Addressing smartphone addiction requires a multifaceted approach that considers behavioral, psychological, and environmental determinants, as well as regional differences in technology use.

FUNCTIONAL EXERCISE CAPACITY IN YOUNG ADULTS

Functional exercise capacity is influenced by a wide range of lifestyle and physiological factors that collectively determine an individual's ability to perform sustained physical activity efficiently, particularly among college students who are undergoing significant behavioral and environmental transitions. Lifestyle factors such as physical activity levels, dietary habits, sleep patterns, and daily routines play a crucial role in maintaining optimal functional capacity and overall physical fitness. Regular engagement in moderate to vigorous physical activity enhances cardiovascular endurance, muscular strength, and respiratory efficiency, which are essential components of functional exercise capacity. However, modern lifestyle changes characterized by increased reliance on digital devices have significantly altered activity patterns among students, leading to reduced engagement in physical exercise. Reduced physical activity is one of the primary contributors to declining fitness levels, as it results in decreased energy expenditure and gradual deconditioning of physiological systems. The increasing prevalence of screen-based activities, particularly smartphone use, has further exacerbated this issue by promoting sedentary behavior and limiting opportunities for movement. Increased screen time is associated with prolonged sitting, which negatively affects cardiovascular health, metabolic function, and musculoskeletal integrity. Over time, these effects lead to reduced aerobic capacity, diminished endurance, and impaired ability to perform physical tasks efficiently. Physiological factors such as age, gender, body composition, and genetic predisposition also influence functional exercise capacity, although lifestyle factors often have a more immediate and modifiable impact. For instance, individuals with higher body fat percentage and lower muscle mass may experience reduced exercise tolerance and increased fatigue during physical activity. Cardiovascular fitness, measured through parameters such as heart rate response and oxygen uptake, is a key determinant of exercise capacity and is directly influenced by physical activity levels.




Figure 5. Determinants of functional exercise capacity in college students.

Respiratory efficiency, including lung capacity and ventilation rate, plays a significant role in sustaining physical performance during exercise. Behavioral factors, such as motivation, self-discipline, and health awareness, further affect an individual's engagement in physical activity and, consequently, their functional capacity. Students who lack motivation or perceive exercise as a low priority are more likely to adopt sedentary lifestyles, thereby reducing their physical fitness. Environmental influences, including access to recreational facilities, availability of safe outdoor spaces, and institutional support for physical activity, also contribute significantly to exercise behavior. In college settings, factors such as academic workload, time constraints, and social environment can facilitate or hinder participation in physical activity.

For example, students with demanding academic schedules may find it difficult to allocate time for regular exercise, leading to decreased activity levels and reduced fitness levels. Additionally, peer influence and social norms can shape attitudes toward physical activity, either encouraging active lifestyles or reinforcing sedentary behavior. Psychological factors such as stress, anxiety, and depression also impact functional exercise capacity by affecting energy levels, motivation, and overall well-being. Students experiencing high levels of stress may be less likely to engage in physical activity and more likely to engage in passive behaviors, such as smartphone use. Figure 5 shows the determinants of functional exercise capacity, including physical, behavioral, and environmental influences, providing a comprehensive overview of the multiple factors that interact to influence physical fitness in college students. Figure 5 highlights the interconnected nature of these determinants and emphasizes the importance of addressing multiple domains when designing interventions to improve exercise capacity. Understanding these determinants is essential for developing targeted strategies to promote active lifestyles and reduce the negative impact of sedentary behavior on physical health.

ASSOCIATION BETWEEN SMARTPHONE ADDICTION AND PHYSICAL ACTIVITY

A negative association exists between smartphone addiction and physical activity levels, with numerous studies demonstrating that increased reliance on smartphones is linked to a decline in engagement in regular exercise among college students. This relationship is primarily driven by the displacement effect, wherein time spent on smartphones replaces time that could otherwise be used for physical activities such as sports, walking, or structured exercise. Increased screen time contributes to prolonged sedentary behavior, which reduces overall energy expenditure and leads to lower physical fitness.




Figure 6. Inverse relationship between smartphone addiction and physical activity levels.

College students who exhibit higher levels of smartphone addiction are more likely to spend extended periods sitting or lying down while interacting with their devices, further limiting opportunities for physical movement. The convenience and accessibility of digital entertainment, including social media, streaming platforms, and mobile gaming, make smartphones a preferred source of leisure activity, thereby reducing motivation to engage in physically demanding tasks. Behavioral patterns associated with smartphone addiction, such as compulsive checking and multitasking, also interfere with structured exercise routines and reduce adherence to fitness programs. In addition, excessive smartphone use is often associated with irregular daily schedules and poor time management, which can further hinder participation in planned physical activities. Psychological factors such as decreased motivation, increased fatigue, and reduced self-regulation play a significant role in mediating the relationship between smartphone addiction and physical inactivity. Individuals with high smartphone dependence may experience lower intrinsic motivation for exercise, preferring passive activities that require minimal physical effort. Moreover, the overstimulation associated with continuous digital engagement can lead to mental fatigue, which reduces the willingness to participate in physical activity. Sleep disturbances resulting from excessive screen time, particularly during late-night hours, also contribute to reduced energy levels and decreased physical performance during the day. Social and environmental factors further influence this relationship, as peer behavior and cultural norms surrounding technology use can either encourage or discourage active lifestyles. Students who are part of social groups that prioritize digital interaction over physical engagement are more likely to adopt sedentary behaviors and neglect exercise. Conversely, supportive environments that promote physical activity can help mitigate the negative impact of smartphone addiction on fitness levels. The physiological consequences of reduced physical activity include decreased cardiovascular endurance, reduced muscular strength, and impaired metabolic function, all of which contribute to lower functional exercise capacity. Over time, these changes can increase the risk of chronic health conditions such as obesity, cardiovascular disease, and metabolic syndrome. The inverse relationship between smartphone use and physical activity is further explained in Figure 6, which depicts how increasing smartphone use correlates with decreasing physical activity intensity, illustrating a clear downward trend in activity levels as screen time increases.

This graphical representation highlights the dose-dependent nature of the relationship, indicating that higher levels of smartphone engagement are associated with progressively lower levels of physical exertion. Figure 6 also emphasizes the cumulative impact of prolonged screen exposure on physical health, demonstrating how small reductions in daily activity can lead to significant declines in overall fitness over time. Understanding this inverse relationship is essential for developing targeted interventions aimed at reducing smartphone addiction and promoting physical activity among college students. Strategies such as time management training, digital detox programs, and structured exercise interventions have been shown to improve both behavioral and physical outcomes. Additionally, integrating technology-based solutions such as fitness tracking applications and activity reminders can help encourage movement and reduce sedentary behavior.

DOSE RESPONSE RELATIONSHIP

Higher duration of smartphone use is consistently associated with greater declines in physical activity and exercise capacity among college students, indicating a significant behavioral and physiological impact of prolonged screen exposure. As daily smartphone usage increases, the amount of time available for engaging in physical activities correspondingly decreases, leading to reduced overall movement and energy expenditure. This inverse relationship is particularly concerning for young adults, as consistent physical activity is essential for maintaining cardiovascular fitness, muscular strength, and metabolic health. Prolonged screen exposure often results in extended periods of sedentary behavior, which contributes to physical deconditioning and reduced functional exercise capacity over time. The cumulative effect of such inactivity is reflected in diminished aerobic endurance and lower tolerance for physical exertion, even during routine activities. Additionally, increased smartphone use is frequently associated with behavioral patterns such as late-night usage and irregular daily routines, which can further disrupt opportunities for exercise. Sleep disturbances caused by excessive screen time negatively impact recovery and energy levels, thereby reducing the ability to engage in physical activity during waking hours. Psychological factors also contribute to this relationship, as individuals with higher smartphone use may experience reduced motivation, increased fatigue, and a preference for passive forms of leisure. Over time, these behavioral tendencies reinforce sedentary habits and further limit participation in physical exercise. The dose-dependent nature of this relationship suggests that even small increases in daily screen time can lead to measurable declines in physical activity levels and exercise performance. For example, students who spend several hours per day on their smartphones are less likely to meet recommended physical activity guidelines compared to those with moderate usage patterns. This gradual reduction in activity contributes to long-term physiological changes such as decreased cardiovascular efficiency, reduced muscle endurance, and impaired metabolic function. Furthermore, prolonged smartphone use often occurs in static postures, which not only limits movement but also contributes to musculoskeletal discomfort and reduced physical efficiency. Environmental and social factors may further amplify this effect, as increased digital engagement is often normalized within peer groups, reducing the perceived importance of physical activity. The cumulative impact of prolonged screen exposure is therefore not limited to immediate behavioral changes but extends to long-term health outcomes, including reduced functional exercise capacity and increased risk of chronic diseases. Understanding this relationship is essential for developing targeted interventions aimed at reducing excessive smartphone use and promoting active lifestyles among college students. Strategies such as limiting screen time, encouraging regular physical activity breaks, and integrating movement into daily routines can help mitigate the negative effects of prolonged smartphone use.

IMPACT ON FUNCTIONAL EXERCISE CAPACITY

Smartphone addiction negatively impacts exercise capacity through multiple interrelated mechanisms, including reduced physical activity, poor sleep quality, and progressive physiological deconditioning, all of which contribute to declining fitness levels among college students. Reduced physical activity is one of the most direct consequences of excessive smartphone use, as prolonged screen engagement limits time available for exercise and promotes sedentary behavior. Students with high levels of smartphone addiction often spend several hours daily in inactive postures, leading to decreased energy expenditure and reduced stimulation of cardiovascular and muscular systems. Over

time, this inactivity results in diminished aerobic capacity, reduced muscular endurance, and impaired overall physical performance. Poor sleep is another critical factor linking smartphone addiction to reduced exercise capacity, as excessive screen exposure, particularly during nighttime, disrupts circadian rhythms and delays sleep onset. The blue light emitted by smartphone screens suppresses melatonin production, leading to shorter sleep duration and poorer sleep quality. Inadequate sleep negatively affects recovery processes, reduces energy levels, and impairs cognitive and physical performance during the day. As a result, individuals with high smartphone usage often experience fatigue and decreased motivation for engaging in physical activity. Physiological deconditioning further exacerbates this issue, as prolonged inactivity leads to a gradual decline in cardiovascular efficiency, muscle strength, and metabolic function. This deconditioning process reduces the body's ability to perform sustained physical activity, thereby lowering functional exercise capacity. Additionally, excessive smartphone use is associated with poor posture and musculoskeletal strain, particularly in the neck, shoulders, and lower back, which can further limit physical performance and increase discomfort during exercise. Psychological factors such as stress, anxiety, and reduced self-regulation also play a mediating role by influencing both smartphone usage patterns and participation in physical activity. Individuals experiencing higher levels of psychological distress may rely more heavily on smartphones as a coping mechanism, thereby increasing screen time and reducing engagement in health-promoting behaviors. This creates a reinforcing cycle in which smartphone addiction leads to reduced physical activity and poor health outcomes, which in turn further increase dependence on digital devices. Table 2 presents a comparison of functional exercise capacity parameters between high and low smartphone usage groups, clearly demonstrating reduced endurance and performance in highly addicted individuals, with lower values observed in measures such as six-minute walk distance, VO_2 max, and exercise tolerance. Table 2 highlights significant differences in cardiovascular and respiratory efficiency, indicating that individuals with excessive smartphone use exhibit compromised physiological functions. These findings are consistent across multiple studies, reinforcing the negative impact of smartphone addiction on physical fitness and performance. Furthermore, the data suggest that even moderate increases in smartphone usage can lead to measurable declines in exercise capacity, emphasizing the importance of early intervention. Environmental and lifestyle factors such as academic workload, lack of access to recreational facilities, and social norms surrounding technology use may further influence this relationship. Addressing smartphone addiction, therefore, requires a comprehensive approach that targets behavioral, psychological, and environmental determinants of health.

Interventions such as structured exercise programs, sleep hygiene education, and digital detox strategies have shown potential for improving both smartphone usage patterns and physical fitness outcomes. Encouraging regular physical activity and reducing sedentary behavior can help reverse the effects of physiological deconditioning and restore the functional exercise capacity.

Table 2. Comparison of functional exercise capacity between high and low smartphone addiction groups.

Parameter	High smartphone addiction group	Low smartphone addiction group	Interpretation
Six-minute walk distance (meters)	Lower	Higher	Reduced endurance in the high addiction group
VO_2 max. (ml/kg/min)	Decreased	Normal or higher	Lower aerobic capacity with higher smartphone use
Resting heart rate (beats/min)	Higher	Lower	Indicates poorer cardiovascular fitness
Heart rate recovery	Slower	Faster	Delayed recovery suggests reduced fitness
Physical activity level	Low	Moderate to high	Sedentary behavior linked to addiction
Muscular endurance	Reduced	Better	Lower stamina in high usage individuals
Flexibility	Decreased	Normal	Poor posture and inactivity affect flexibility
Perceived exertion (RPE)	Higher	Lower	Higher fatigue during exercise




Figure 7. Psychological mediation model linking smartphone addiction and functional exercise capacity.

PSYCHOLOGICAL AND BEHAVIORAL MEDIATORS

Psychological factors such as stress, anxiety, and depression play a crucial role in influencing both smartphone use and physical activity levels among college students, thereby shaping the relationship between digital behavior and functional exercise capacity. High levels of academic stress and social pressure often lead students to seek immediate relief through smartphone use, particularly through social media and entertainment applications that provide temporary distraction and emotional comfort. This coping mechanism, although initially effective in reducing stress, can gradually develop into habitual and excessive use, increasing the risk of smartphone addiction. Anxiety further contributes to this pattern, as individuals experiencing heightened worry or social apprehension may rely on digital communication as a safer alternative to face-to-face interaction. This increased reliance on smartphones reduces opportunities for physical engagement and promotes sedentary behavior, thereby negatively affecting physical activity levels. Depression is also strongly associated with both increased smartphone use and reduced physical activity, as individuals with depressive symptoms often exhibit low motivation, fatigue, and decreased interest in physical exercise. These psychological states create a barrier to engaging in health-promoting behaviors, reinforcing inactivity, and further worsening physical fitness. The interaction between these psychological factors and behavioral patterns forms a self-reinforcing cycle in which increased stress, anxiety, and depression lead to greater smartphone use, which in turn reduces physical activity and contributes to further psychological distress. This cyclical relationship highlights the complex interplay between mental health and lifestyle behaviors in determining overall well-being. Reduced physical activity, in particular, has been shown to exacerbate symptoms of stress and depression, as regular exercise is known to have positive effects on mood regulation and psychological resilience. Therefore, the decline in physical activity associated with smartphone addiction not only affects physical health but also contributes to worsening mental health outcomes. Additionally, excessive smartphone use can lead to cognitive overload and mental fatigue, which further reduces motivation to engage in physical activities and increases preference for passive behaviors. Social comparison and exposure to unrealistic standards on digital platforms may also contribute to feelings of inadequacy and anxiety, reinforcing dependence on smartphones for validation and escape. Environmental and contextual factors such as academic workload, peer influence, and lack of social support can further intensify psychological stress and influence smartphone usage patterns. Figure 7 illustrates the mediating role of psychological variables in the relationship between smartphone

addiction and exercise capacity, demonstrating how stress, anxiety, and depression act as intermediate factors that influence both digital behavior and physical fitness outcomes. Figure 7 highlights the bidirectional nature of these relationships, showing that psychological distress not only leads to increased smartphone use but is also exacerbated by reduced physical activity and poor health. This mediation model emphasizes the importance of addressing psychological factors when designing interventions aimed at reducing smartphone addiction and improving exercise capacity. Interventions that focus on stress management, emotional regulation, and mental health support can help break this cycle by reducing reliance on smartphones and encouraging active lifestyles. Mindfulness-based approaches, cognitive behavioral strategies, and counseling services are effective in improving psychological well-being and promoting healthier behavioral patterns. Encouraging participation in physical activity not only improves physical fitness but also enhances mood and reduces symptoms of anxiety and depression, thereby creating a positive feedback loop that supports overall health. Integrating mental health and physical activity interventions within college settings can therefore provide a comprehensive approach to addressing the interconnected challenges of smartphone addiction and reduced exercise capacity.

SLEEP AND CIRCADIAN RHYTHM DISTURBANCES

Excessive smartphone use has been widely associated with significant disruptions in sleep patterns among college students, primarily due to prolonged screen exposure and irregular usage habits that interfere with normal circadian rhythms. The use of smartphones late at night delays sleep onset, as engaging content and continuous notifications keep the brain in an alert state, making it difficult for individuals to transition into restful sleep. One of the key mechanisms underlying this disruption is the emission of blue light from smartphone screens, which suppresses melatonin production and alters the body's natural sleep-wake cycle. As a result, individuals who frequently use smartphones before bedtime often experience shorter sleep duration and poorer sleep quality compared to those with limited nighttime use. Sleep deprivation resulting from these patterns has a direct impact on physical health, particularly in terms of reduced energy levels and increased daytime fatigue. Fatigue negatively affects motivation and willingness to engage in physical activity, leading to decreased participation in exercise and reduced overall physical fitness. Furthermore, inadequate sleep impairs cognitive and motor functions, which are essential for performing physical tasks efficiently and safely. Reduced alertness and slower reaction times can compromise exercise performance and increase the risk of injury during physical activity. Sleep deprivation also affects physiological recovery processes, as the body relies on sleep for muscle repair, energy restoration, and hormonal regulation. Disruption in these processes leads to decreased endurance and reduced ability to sustain prolonged physical activity. Additionally, poor sleep is associated with increased levels of stress hormones such as cortisol, which can further impair physical performance and contribute to fatigue. The cumulative effect of these factors is a decline in functional exercise capacity, as individuals become less capable of performing activities that require sustained effort. Over time, chronic sleep deprivation linked to excessive smartphone use can lead to long-term health consequences, including reduced cardiovascular fitness and increased risk of metabolic disorders. Psychological effects such as irritability, reduced concentration, and mood disturbances further compound the impact of poor sleep on physical activity and overall well-being. In college students, where academic demands and lifestyle factors already challenge healthy sleep patterns, excessive smartphone use exacerbates these issues and creates a cycle of poor sleep and reduced physical performance. Addressing this problem requires interventions aimed at promoting healthy sleep hygiene, such as limiting screen time before bedtime and encouraging consistent sleep schedules.

POSTURAL AND MUSCULOSKELETAL IMPLICATIONS

Prolonged smartphone use has emerged as a significant contributor to musculoskeletal disorders, particularly affecting the cervical spine, shoulders, and upper back, due to sustained poor posture and repetitive strain, as individuals tend to adopt forward head posture and excessive neck flexion while using mobile devices, which increases mechanical load on the cervical spine and surrounding musculature, ultimately leading to discomfort, pain, and reduced physical efficiency during daily and

exercise-related activities. Continuous use of smartphones encourages static postures that alter normal spinal alignment, resulting in increased stress on ligaments and muscles, thereby contributing to fatigue and reduced endurance capacity during physical exertion. Sustained neck flexion during smartphone activities significantly elevates muscle activation in the neck extensors and upper trapezius, which accelerates muscle fatigue and limits the ability to perform prolonged physical tasks efficiently. Research indicates that prolonged smartphone usage is strongly associated with neck pain and decreased neck muscle endurance, which directly affects exercise tolerance and physical performance in young adults. Improper posture during smartphone use, such as slouching or forward head positioning, has been identified as a major risk factor for musculoskeletal discomfort in the neck, shoulders, and lower back regions. Extended periods of smartphone engagement led to cumulative biomechanical stress, which disrupts normal musculoskeletal function and contributes to chronic pain conditions that impair movement efficiency. Experimental findings demonstrate that even short durations of continuous smartphone use can result in noticeable increases in neck and shoulder discomfort, indicating the rapid onset of musculoskeletal strain. The repetitive nature of smartphone interactions, including scrolling and typing, further exacerbates strain on upper limb muscles and joints, compounding the overall musculoskeletal burden. Alterations in posture and gait associated with excessive smartphone use reduce overall physical stability and coordination, thereby negatively influencing exercise performance and functional mobility. Forward head posture induced by prolonged smartphone use increases spinal loading, which not only causes pain but also reduces respiratory efficiency and endurance during physical activity. Musculoskeletal discomfort caused by excessive smartphone use often leads to decreased participation in physical activity, further lowering exercise tolerance and contributing to sedentary behavior patterns. Over time, these combined effects of poor posture, muscle fatigue, and chronic discomfort significantly impair physical efficiency, highlighting the importance of ergonomic awareness and moderated smartphone use in maintaining musculoskeletal health and optimal exercise capacity.

PHYSIOLOGICAL MECHANISMS

Smartphone addiction has been increasingly associated with reduced cardiovascular efficiency and impaired oxygen utilization, primarily due to sedentary behavior and decreased engagement in physical activity, which are essential determinants of cardiorespiratory fitness and overall health status. Excessive smartphone use often replaces time that could be spent on moderate to vigorous physical activity, leading to diminished aerobic capacity and reduced efficiency of the cardiovascular system during exercise. Cardiorespiratory fitness, commonly measured through maximal oxygen uptake, is highly dependent on regular physical activity, and reduced activity linked to smartphone addiction contributes to lower oxygen consumption efficiency. A sedentary lifestyle induced by prolonged smartphone engagement negatively affects heart rate variability and circulation, which are critical for maintaining optimal cardiovascular endurance. Reduced physical activity levels associated with smartphone addiction have been shown to correlate with increased body mass index and decreased metabolic efficiency, further impairing cardiovascular performance. Additionally, poor posture and restricted thoracic expansion during prolonged device use can indirectly limit respiratory efficiency, thereby reducing oxygen intake and utilization during physical exertion. Lower engagement in exercise due to smartphone overuse contributes to diminished stroke volume and cardiac output over time, reducing the body's ability to meet oxygen demands during physical activity. Studies suggest that individuals with higher levels of smartphone addiction demonstrate poorer fitness indicators, including reduced endurance and lower exercise tolerance, highlighting its impact on cardiovascular health. The decline in cardiovascular efficiency is further compounded by associated factors such as poor sleep quality and increased stress, both of which are commonly linked with excessive smartphone use and negatively influence oxygen utilization. Moreover, reduced participation in physical training diminishes mitochondrial efficiency and aerobic metabolism, leading to decreased energy production and endurance capacity. Over time, these physiological alterations result in a measurable decline in overall fitness levels, as the cardiovascular and respiratory systems become less efficient in delivering and utilizing oxygen during physical activity. Therefore, smartphone addiction not only promotes sedentary behavior but also significantly impairs cardiovascular efficiency and oxygen utilization, ultimately reducing overall physical fitness and increasing the risk of long-term health complications.

GENDER DIFFERENCES AND SOCIODEMOGRAPHIC FACTORS

Differences in smartphone usage patterns are evident across gender and socioeconomic groups, significantly influencing both addiction levels and engagement in physical activity. Research indicates that males and females tend to use smartphones for different purposes, with males more inclined toward gaming and females more engaged in social networking, leading to varying addiction risk profiles. Studies have shown that sex plays a critical role in determining smartphone addiction severity, with some recent findings suggesting higher prevalence rates among females, particularly due to increased social media engagement and emotional attachment to digital communication. Additionally, gender differences extend to physical activity levels, where higher smartphone addiction is often associated with lower participation in exercise, especially among adolescents and young adults. Socioeconomic status further contributes to disparities in smartphone usage, as individuals from higher-income groups may have greater access to smartphones and digital resources, increasing their screen time and potential dependency, while those from lower socioeconomic backgrounds may experience different usage patterns influenced by accessibility and lifestyle factors. Research also highlights that urban populations, often associated with higher socioeconomic status, demonstrate increased smartphone dependency compared to rural populations, which can influence sedentary behavior and reduce physical activity. Furthermore, behavioral and psychological factors linked to socioeconomic conditions, such as stress, lifestyle habits, and access to recreational facilities, mediate the relationship between smartphone addiction and physical activity levels. Evidence suggests that individuals with greater screen exposure due to socioeconomic advantages may engage less in outdoor or physical activities, contributing to declining fitness levels and increased sedentary lifestyles. Conversely, limited access to digital devices in lower socioeconomic groups may reduce the risk of addiction but does not necessarily guarantee higher physical activity, as other environmental and social constraints may exist.

ROLE OF ACADEMIC ENVIRONMENT AND LIFESTYLE FACTORS

Academic stress and lifestyle habits play a crucial role in shaping smartphone use and physical activity levels among college students, as increased academic pressure often leads individuals to seek distraction and relief through excessive smartphone engagement, thereby increasing the risk of addiction. High levels of perceived stress among students have been identified as a significant predictor of problematic smartphone use, as students frequently turn to digital devices to cope with anxiety, academic workload, and emotional strain. This coping mechanism, although temporarily relieving stress, contributes to prolonged screen time and reduced participation in physical activities, ultimately affecting overall health and fitness levels. Lifestyle habits such as irregular sleep patterns, poor time management, and sedentary routines further exacerbate smartphone dependency, creating a cycle where increased usage leads to decreased physical activity and vice versa. Moreover, academic demands often limit the time available for exercise, and students with higher stress levels are less likely to engage in structured physical activity, further reducing their fitness levels. Research also indicates that physical activity can act as a protective factor by improving self-control and reducing stress perception, thereby decreasing smartphone addiction tendencies among college students. Additionally, emotional regulation and coping strategies influenced by lifestyle habits significantly mediate the relationship between academic stress and smartphone use, highlighting the behavioral complexity of this issue. The interaction between academic stress and unhealthy lifestyle patterns contributes to decreased motivation for physical activity and increased reliance on smartphones for entertainment and social interaction. Consequently, addressing academic stress and promoting healthier lifestyle habits are essential strategies to reduce smartphone addiction and improve physical activity levels among college students, thereby enhancing their overall well-being and academic performance.

DIGITAL HEALTH AND WEARABLE TECHNOLOGY

Wearable devices and health applications have emerged as effective tools for monitoring physical activity and reducing sedentary behavior by providing real-time feedback, goal tracking, and personalized health insights, which enhance user awareness and motivation to engage in regular exercise.

These technologies utilize sensors to track steps, heart rate, and energy expenditure, enabling individuals to monitor their daily activity levels and make informed decisions to improve their health. Health applications integrated with smartphones further support behavior change by offering reminders, progress reports, and tailored exercise programmes that encourage consistent physical activity. The use of wearable technology has been associated with increased physical activity levels and reduced sedentary time, particularly among young adults and college students. Additionally, gamification features and social connectivity embedded within these applications promote engagement and adherence to physical activity routines by fostering competition and providing peer support. Figure 8 presents an integrated intervention model combining digital health tools and physical activity strategies to improve outcomes, illustrating how wearable devices, mobile applications, and structured exercise programs can work synergistically to enhance physical fitness and reduce inactivity in older adults. Such models emphasize continuous monitoring, behavioral feedback, and goal setting as key components in promoting long-term adherence to healthy lifestyles. Furthermore, wearable devices can assist healthcare professionals in remotely tracking patient activity and providing timely interventions, thereby improving overall health outcomes and preventive care quality. The integration of artificial intelligence in modern health applications has further enhanced their capability to provide personalized recommendations based on user data, optimize exercise plans, and improve adherence. Evidence suggests that individuals who regularly use wearable devices demonstrate better physical activity patterns and improved cardiovascular health than non-users. Moreover, these technologies help identify periods of inactivity and prompt users to take breaks, thereby reducing prolonged sedentary behavior and associated health risks.

INTERVENTION STRATEGIES

Interventions aimed at reducing smartphone addiction and improving physical activity levels include structured exercise programs, behavioral therapy, and digital detox approaches, each addressing different aspects of behavior and lifestyle modification to enhance overall health outcomes. Structured exercise programs play a crucial role in increasing physical activity levels and improving cardiovascular and musculoskeletal fitness, while also reducing screen time by encouraging individuals to engage in scheduled physical routines. Behavioral therapy, particularly cognitive behavioral therapy, is effective in addressing problematic smartphone use by modifying thought patterns and promoting healthier coping strategies.




Figure 8. Integrated intervention model for reducing smartphone addiction and improving exercise capacity.

Table 3. Intervention strategies for smartphone addiction and improvement of functional exercise capacity.

Intervention type	Strategy description	Target outcome	Effectiveness
Structured exercise programs	Regular aerobic and strength training sessions integrated into the daily routine	Improved cardiovascular fitness. increased endurance	High
Behavioral therapy	Cognitive behavioral therapy to modify addictive patterns and improve self-control	Reduced smartphone dependency. better coping strategies	High
Digital detox programs	Scheduled reduction or restriction of smartphone usage	Decreased screen time. improved focus and activity levels	Moderate to high
Health education	Awareness programs on the risks of excessive smartphone use and the benefits of physical activity	Improved knowledge. behavior change	Moderate
Time management training	Planning daily schedules to balance academic work, screen time, and exercise	Better routine adherence. reduced overuse	Moderate
Use of wearable devices	Activity trackers and fitness apps to monitor movement and encourage activity	Increased physical activity. Reduced sedentary behavior	High
Mindfulness and stress management	Techniques such as meditation and relaxation exercises	Reduced stress. improved mental well-being	Moderate
Social support programs	Peer group activities and team-based exercise interventions	Increased motivation. sustained engagement	High
Institutional policies	Guidelines limiting smartphone use in academic settings and promoting physical activity	Controlled usage. improved lifestyle habits	Moderate to high

Digital detox interventions, which involve intentional reduction or temporary cessation of smartphone use, help individuals break dependency cycles and restore balance between digital engagement and real-world activities. Table 3 summarizes these strategies and their effectiveness, providing a practical framework for implementation, highlighting how combined approaches yield better outcomes than single interventions. Additionally, structured interventions that integrate physical exercise with behavioral counseling have demonstrated significant improvements in both physical fitness and a reduction in smartphone addiction levels. Educational programs that raise awareness about the negative impacts of excessive smartphone use further support behavior change and promote healthier lifestyle habits among students. Time management strategies and goal-setting techniques are also essential components of intervention programs, helping individuals allocate time effectively between academic responsibilities, physical activity, and digital usage. Furthermore, social support and peer-based interventions enhance adherence to these strategies by fostering accountability and motivation. Evidence suggests that multi-component interventions combining exercise, behavioral therapy, and digital detox approaches are more effective in sustaining long-term behavior change and improving overall well-being. These interventions not only reduce sedentary behavior but also enhance mental health, physical fitness, and productivity among individuals with high smartphone usage.

LONGITUDINAL AND CAUSAL EVIDENCE

Most studies examining the relationship between smartphone use, physical activity, and health outcomes are predominantly cross-sectional in design, which limits the ability to establish causal relationships between variables and restricts the depth of interpretation regarding long-term effects. Cross-sectional studies provide a snapshot of associations at a single point in time, making it difficult to determine whether smartphone addiction leads to reduced physical activity or if inactive individuals are more likely to engage in excessive smartphone use. This limitation reduces the strength of evidence available for developing effective interventions, as causal pathways remain unclear and potentially confounded by external variables. Additionally, cross-sectional designs are susceptible to biases, such

as self-reporting errors and recall bias, which may affect the accuracy of data related to smartphone usage and physical activity levels. The lack of temporal sequencing in such studies further complicates the understanding of how behavioral patterns evolve over time and interact with physiological and psychological outcomes. Therefore, longitudinal research designs are essential to track changes in smartphone use and physical activity over extended periods, allowing for a clearer understanding of cause-and-effect relationships. These studies provide insights into the progression of smartphone addiction and its long-term impact on physical fitness, mental health, and lifestyle behaviors. Furthermore, longitudinal approaches enable researchers to control confounding variables and observe trends, thereby improving the validity and reliability of the findings. Intervention-based longitudinal studies are particularly valuable for assessing the effectiveness of strategies aimed at reducing smartphone addiction and increasing physical activity levels over time. Such research can also help identify critical periods for intervention and inform policy development in educational and healthcare settings.

Limitations of Existing Studies

Research examining smartphone use, physical activity, and associated health outcomes is often limited by reliance on self-reported data, which can introduce biases such as social desirability and inaccurate recall, thereby affecting the validity of findings. Participants may overestimate their physical activity levels or underestimate their screen time, leading to discrepancies between reported and actual behaviors. Such inaccuracies reduce the reliability of conclusions drawn regarding the relationship between smartphone addiction and health outcomes. Additionally, subjective measures of smartphone usage and activity patterns lack the precision offered by objective monitoring tools such as accelerometers and digital tracking applications. Another significant limitation in many studies is the use of small sample sizes, which restricts statistical power and limits the ability to detect meaningful associations between variables. Small samples may not adequately represent the broader population, leading to findings that cannot be generalized across different demographic or geographic groups. This issue is particularly relevant in studies focusing on college students, where samples are often drawn from a single institution or region, reducing external validity. Furthermore, limited sample diversity in terms of age, gender, and socioeconomic status can further constrain the applicability of results to wider populations. The combination of self-reported data and small sample sizes increases the risk of bias and reduces confidence in the observed relationships between smartphone use and physical activity. To address these limitations, future research should incorporate larger and more diverse samples along with objective measurement tools to enhance data accuracy and generalizability. Longitudinal and multicenter studies can further strengthen evidence by providing more robust and representative findings.

Public Health and Policy Implications

Smartphone addiction has emerged as a significant public health concern due to its widespread prevalence and its negative impact on physical, mental, and social well-being, necessitating comprehensive institutional and policy-level interventions to address this growing issue. The increasing dependence on smartphones has been associated with sedentary behavior, reduced physical activity, and various health complications, highlighting the need for structured public health strategies. Educational institutions play a critical role in mitigating smartphone addiction by implementing awareness programs, promoting digital literacy, and encouraging balanced technology use among students. Policy-level interventions such as guidelines on screen time, integration of physical activity programs, and restrictions on excessive device use in academic settings can contribute to healthier behavioral patterns. Healthcare systems can also support intervention efforts by incorporating screening for smartphone addiction into routine health assessments and providing counseling services where necessary. Public health campaigns focusing on the risks associated with excessive smartphone use and the benefits of active lifestyles can further enhance community awareness and engagement. Additionally, collaboration between policymakers, educators, and healthcare professionals is essential to develop effective, evidence-based strategies for reducing smartphone dependency. Workplace

policies promoting digital well-being and encouraging regular breaks from screen use can also help reduce overall exposure and associated health risks. Technological solutions, such as usage tracking applications and digital well-being tools, can be integrated into broader intervention frameworks to support behavior change. Research also emphasizes the importance of culturally tailored interventions that consider demographic and socioeconomic differences in smartphone use patterns. Strong regulatory frameworks and institutional support are necessary to ensure the successful implementation and sustainability of these interventions.

Future Research Directions

Future research on smartphone addiction and its impact on physical activity and health outcomes should prioritize experimental and longitudinal study designs to establish clearer causal relationships and enhance the strength of the evidence in this field. Experimental studies, including randomized controlled trials, can help determine the effectiveness of specific interventions aimed at reducing smartphone use and improving physical activity. Such designs allow researchers to control for confounding variables and directly assess the impact of targeted strategies on behavioral and physiological outcomes. Longitudinal studies are equally important because they enable the tracking of changes in smartphone use patterns and health indicators over time, providing insights into long-term effects and trends. These approaches can help identify critical periods during which individuals are most vulnerable to developing smartphone addiction and experiencing its negative effects. Furthermore, longitudinal research can reveal bidirectional relationships between smartphone use and physical activity, clarifying whether one behavior influences the other. Incorporating objective measurement tools, such as wearable devices and digital tracking systems, in future studies can improve data accuracy and reduce reliance on self-reported measures. Multicenter and cross-cultural studies are also needed to enhance the generalizability of the findings across diverse populations and settings. Additionally, integrating qualitative methods with quantitative designs can provide a deeper understanding of the behavioral motivations and contextual factors influencing smartphone use. Advanced analytical techniques, including machine learning, may further support the identification of complex patterns and smartphone addiction.

DISCUSSION

The association between smartphone addiction and functional exercise capacity is multifactorial, involving a complex interplay of behavioral, physiological, and psychosocial factors that collectively influence physical performance and health outcomes. Excessive smartphone use contributes to sedentary behavior, which directly reduces cardiovascular endurance, muscular strength, and flexibility, thereby impairing functional exercise capacity [21]. Prolonged screen time is also linked to poor posture and musculoskeletal discomfort, which can limit mobility and reduce the ability to perform physical tasks efficiently. Additionally, smartphone addiction is associated with disrupted sleep patterns, leading to fatigue and decreased energy levels, which negatively impact exercise performance and recovery [22–26]. Psychological factors, such as stress, anxiety, and reduced motivation, further mediate this relationship, as individuals with higher smartphone dependency may exhibit lower engagement in physical activity. The decline in functional capacity is also influenced by reduced participation in structured exercise programs, as increased screen time often replaces opportunities for physical activity. Moreover, decreased social interaction and outdoor activities linked to smartphone overuse can further contribute to lower physical fitness and endurance levels. Addressing this multifactorial association requires integrated interventions that combine physical activity promotion, behavioral modification, and digital usage regulation to achieve optimal outcomes [27–29]. Structured exercise programs, along with cognitive behavioral strategies, can help improve physical fitness and reduce smartphone dependency. The incorporation of digital health tools, such as activity trackers and monitoring applications, can also support behavior change and enhance adherence to healthy routines. Furthermore, educational and policy-level interventions are essential to raise awareness and promote balanced technology use among different populations [30].

CONCLUSION

The association between smartphone addiction and functional exercise capacity in college students highlights a growing concern that extends beyond behavioral habits into physical health and overall well-being, as excessive smartphone use contributes to sedentary lifestyles, reduced physical activity, and impaired physiological functioning, all of which negatively influence exercise tolerance and performance. Factors such as poor posture, disrupted sleep, psychological stress, and decreased motivation further compound this relationship, making it multifactorial and complex, therefore emphasizing the need for a holistic understanding of how digital behaviors interact with physical fitness in this population. Moreover, the evidence suggests that unchecked smartphone addiction can gradually diminish cardiovascular efficiency, muscular endurance, and daily functional capacity, potentially leading to long-term health consequences if not addressed early. Consequently, there is a clear need for targeted interventions within college settings that promote balanced technology use alongside regular physical activity, including structured exercise programs, awareness initiatives, and behavioral strategies aimed at reducing screen time and encouraging active lifestyles. Additionally, institutional support and policy-level measures can play a vital role in creating environments that foster healthy habits and limit excessive smartphone dependence, while the integration of digital health tools may also offer opportunities for monitoring and improving activity levels. Ultimately, addressing smartphone addiction among college students requires a coordinated and multidisciplinary approach that combines education, behavioral change, and supportive environments to enhance functional exercise capacity and promote long-term health, ensuring that technological advancements do not compromise the physical fitness and well-being of the younger population.

REFERENCES

1. Daniele A, Lucas SJE, Rendeiro C. Detrimental effects of physical inactivity on peripheral and brain vasculature in humans: Insights into mechanisms, long-term health consequences and protective strategies. *Front Physiol.* 2022;13:998380. doi:10.3389/fphys.2022.998380. PMID: 36237532; PMCID: PMC9553009.
2. Bi S, Yuan J, Wang Y, Zhang W, Zhang L, Zhang Y, et al. Effectiveness of digital health interventions in promoting physical activity among college students: Systematic review and meta-analysis. *J Med Internet Res.* 2024;26:e51714. doi:10.2196/51714. PMID: 39566049; PMCID: PMC11618011.
3. Teychenne M, Sousa GM, Baker T, Liddelow C, Babic M, Chantry AJ, et al. Domain-specific physical activity and mental health: An updated systematic review and multilevel meta-analysis in a combined sample of 3.3 million people. *Br J Sports Med.* 2026;60(4):267–285. doi:10.1136/bjsports-2025-109806. PMID: 40903203; PMCID: PMC12702480.
4. Currier BS, D'Souza AC, Singh MAF, Lowisz CV, Rawson ES, Schoenfeld BJ, et al. American College of Sports Medicine position stand: Resistance training prescription for muscle function, hypertrophy, and physical performance in healthy adults: An overview of reviews. *Med Sci Sports Exerc.* 2026;58(4):851–872. doi:10.1249/MSS.0000000000003897. PMID: 41843416; PMCID: PMC12965823.
5. Lee DS, Jiang T, Crocker J, Way BM. Social media use and its link to physical health indicators. *Cyberpsychol Behav Soc Netw.* 2022;25(2):87–93. doi:10.1089/cyber.2021.0188. PMID: 35021894; PMCID: PMC8864418.
6. Sun Y, Chen J, Ji M, Li X. Wearable technologies for health promotion and disease prevention in older adults: Systematic scoping review and evidence map. *J Med Internet Res.* 2025;27:e69077. doi:10.2196/69077. PMID: 40553512; PMCID: PMC12238792.
7. Kidokoro T, Tomkinson GR, Lang JJ, Suzuki K. Physical fitness before and during the COVID-19 pandemic: Results of annual national physical fitness surveillance among 16,647,699 Japanese children and adolescents between 2013 and 2021. *J Sport Health Sci.* 2023;12(2):246–254. doi:10.1016/j.jshs.2022.11.002. PMID: 36343895; PMCID: PMC9635948.
8. Leung KW, Sum KR, Yang YJ. Patterns of sedentary behavior among older adults in care facilities: A scoping review. *Int J Environ Res Public Health.* 2021;18(5):2710. doi:10.3390/ijerph18052710. PMID: 33800199; PMCID: PMC7967434.

9. Mehmood M, Khan MR, Ishaque A, Tariq R, Himani G. Tech strain: The musculoskeletal impact of electronic devices on young adults: A cross-sectional study. *J Pak Med Assoc.* 2025;75(7):1086–1090. doi:10.47391/JPMA.22334. PMID: 40751618.
10. Zhu C, Li S, Zhang L. The impact of smartphone addiction on mental health and its relationship with life satisfaction in the post-COVID-19 era. *Front Psychiatry.* 2025;16:1542040. doi:10.3389/fpsy.2025.1542040. PMID: 40171312; PMCID: PMC11960497.
11. Zvolensky MJ, Garey L, Rogers AH, Schmidt NB, Vujanovic AA, Storch EA, et al. Psychological, addictive, and health behavior implications of the COVID-19 pandemic. *Behav Res Ther.* 2020;134:103715. doi:10.1016/j.brat.2020.103715. PMID: 32891956; PMCID: PMC7451060.
12. Piszka M, Kwapien E, Brasse P, Staszkiwicz K, Zerdka J, Staszkiwicz KK, et al. The impact of screen time on the health of the pediatric population: Short- and long-term consequences for lifestyle, ophthalmology, and mental health. *Cureus.* 2025;17(11):e96944. doi:10.7759/cureus.96944. PMID: 41409950; PMCID: PMC12706514.
13. National Academies of Sciences, Engineering, and Medicine; Health and Medicine Division; Board on Population Health and Public Health Practice; Committee on the Impact of Social Media on Adolescent Health; Wojtowicz A, Buckley GJ, Galea S, editors. *Social Media and Adolescent Health.* Washington (DC): National Academies Press (US); 2024 Mar 25. 4, The Relation between Social Media and Health. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK603429/>
14. Presta V, Guarnieri A, Laurenti F, Mazzei S, Arcari ML, Mirandola P, et al. The impact of digital devices on children's health: A systematic literature review. *J Funct Morphol Kinesiol.* 2024;9:236. doi:10.3390/jfmk9040236. PMID: 39584889; PMCID: PMC11587142.
15. Philippe TJ, Sikder N, Jackson A, Koblanski ME, Liow E, Pilarinos A, et al. Digital health interventions for delivery of mental health care: Systematic and comprehensive meta-review. *JMIR Ment Health.* 2022;9:e35159. doi:10.2196/35159. PMID: 35551058; PMCID: PMC9109782.
16. Alhassan MA, Al-Otaibi EM, Bin Ofaysan MS, Alarjani KH, Alzuhairi MO, Alarfaj NS, et al. Smartphone addiction and its psychosocial correlates among university students. *Cureus.* 2025;17(8):e91165. doi:10.7759/cureus.91165. PMID: 41018378; PMCID: PMC12475987.
17. Li Z, Zhao YY, Li X, Liu K, Tian DH, Li SX, et al. The relationship between anxiety and parent–child attachment and psychological resilience among Chinese college students: take 100 thousand college students as sample. *Front Psychiatry.* 2026;16:1667736. doi:10.3389/fpsy.2025.1667736. PMID: 41573047; PMCID: PMC12819800.
18. Zhang J, Zeng Y. Effect of college students' smartphone addiction on academic achievement: The mediating role of academic anxiety and moderating role of sense of academic control. *Psychol Res Behav Manag.* 2024;17:933–944. doi:10.2147/PRBM.S442924. PMID: 38464809; PMCID: PMC10924864.
19. Cox S, Dawkins L, Doshi J, Cameron J. Effects of e-cigarettes versus nicotine replacement therapy on short-term smoking abstinence when delivered at a community pharmacy. *Addict Behav Rep.* 2019;10:100202. doi:10.1016/j.abrep.2019.100202. PMID: 31338412; PMCID: PMC6626064.
20. Xiang MQ, Lin L, Wang ZR, Li J, Xu Z, Hu M. Sedentary behavior and problematic smartphone use in Chinese adolescents: The moderating role of self-control. *Front Psychol.* 2020;10:3032. doi:10.3389/fpsyg.2019.03032. PMID: 32038398; PMCID: PMC6985778.
21. Bhandari DJ, Pandya YP, Sharma DB. Smartphone use and its addiction among adolescents in the age group of 16–19 years. *Indian J Community Med.* 2021;46(1):88–92. doi:10.4103/ijcm.IJCM_263_20. PMID: 34035584; PMCID: PMC8117907.
22. Nikolic A, Bukurov B, Kocic I, Vukovic M, Ladjovic N, Vrhovac M, et al. Smartphone addiction, sleep quality, depression, anxiety, and stress among medical students. *Front Public Health.* 2023;11:1252371. doi:10.3389/fpubh.2023.1252371. PMID: 37744504; PMCID: PMC10512032.
23. Wang F. Effects of physical activity on mobile phone addiction among university students: The mediating roles of self-control and resilience. *Front Psychol.* 2025;16:1503607. doi:10.3389/fpsyg.2025.1503607. PMID: 40231003; PMCID: PMC11994967.
24. Atış Akyol N, Atalan Ergin D, Krettmann AK, Essau CA. Is the relationship between problematic mobile phone use and mental health problems mediated by fear of missing out and escapism? *Addict Behav Rep.* 2021;14:100384. doi:10.1016/j.abrep.2021.100384. PMID: 34938842; PMCID: PMC8664862.

-
25. Pinto AJ, Bergouignan A, Dempsey PC, Roschel H, Owen N, Gualano B, et al. Physiology of sedentary behavior. *Physiol Rev.* 2023;103(4):2561–2622. doi:10.1152/physrev.00022.2022. PMID: 37326297; PMCID: PMC10625842.
 26. Patel H, Desai R, Shah M. Relationship between physical activity levels and smartphone usage in university students. *J Phys Act Health.* 2020;17(6):601–608.
 27. Pirwani N, Szabo A. One-year update on physical activity and smartphone addiction in university students: A systematic review of novel research. *Prev Med Rep.* 2025;57:103178. doi:10.1016/j.pmedr.2025.103178. PMID: 40734703; PMCID: PMC12304691.
 28. Pengpid S, Peltzer K. Sedentary behaviour, physical activity and life satisfaction, happiness and perceived health status in university students from 24 countries. *Int J Environ Res Public Health.* 2019;16(12):2084. doi:10.3390/ijerph16122084. PMID: 31200427; PMCID: PMC6617209.
 29. Liu Y, Tan D, Wang P, Xiao T, Wang X, Zhang T. Physical activity moderated the mediating effect of self-control between bullying victimization and mobile phone addiction among college students. *Sci Rep.* 2024;14(1):20855. doi:10.1038/s41598-024-71797-2. PMID: 39242798; PMCID: PMC11379861.
 30. Kumpfer KL, Bluth B. Parent/child transactional processes predictive of resilience or vulnerability to substance abuse disorders. *Subst Use Misuse.* 2004;39(5):671–698. doi:10.1081/JA-120034011. PMID: 15202804.