

International Journal of Cheminformatics

Volume- 04, Issue- 01, Year- 2026

Review Article

Received Date: January 22, 2026

Accepted Date: February 19, 2026

Published Date: February 28, 2026

Vitamin D Mitigates Inflammation and Downregulates Importin $\alpha 3$ in Non-Alcoholic Fatty Liver Disease (NAFLD)

(Research Scholar) Ashok Kujur^{1,2,3}, (Assistant Professor) S. John Mary^{1,3*}, (Research Scholar) Delinta D^{1,3} and (Research Scholar) Ajila A^{1,3}, (Research Scholar) Jesuraj P^{1,3}, (Research Scholar) Christina C^{1,3}, (Assistant Professor) Gunasekar C J³

¹Loyola Institute of Frontier Energy (LIFE), Loyola College, Sterling Road, Chennai, India.

²Creighton University, School of Medicine, 2500 California Plz, Omaha, USA.

³Department of Chemistry, Loyola College, Sterling Road, Chennai, India.

Correspondence Mail: aku77764@creighton.edu

Abstract

Non-alcoholic fatty liver disease (NAFLD) affects about one-third of the population in the United States and Western countries, reaching up to 90% among obese individuals and those undergoing bariatric surgery. Characterized by hepatic fat accumulation, NAFLD progresses from simple steatosis to non-alcoholic steatohepatitis (NASH), fibrosis, and, in rare cases, hepatocellular carcinoma. Linked intricately with metabolic syndrome, including diabetes and obesity, NAFLD development involves insulin resistance and hyperinsulinemia. Experimental evidence consistently associates low vitamin D levels with NAFLD severity, independent of age, gender, and BMI. Pro-inflammatory cytokines, such as TNF- α and IL-6, play critical roles in NAFLD, with positive correlations between liver fibrosis and TNF- α levels in NASH patients. The inflammatory cascade involves NF- κ B activation, Importin $\alpha 3$, and Importin $\alpha 4$. Vitamin D emerges as a key player, up-regulating the NF- κ B inhibitor I κ B α and reducing pro-inflammatory activity. Studies in NASH models highlight vitamin D's positive impact on inflammation, fibrosis, and insulin resistance. Recent research underscores calcitriol's role in decreasing Importin $\alpha 3$ expression, influencing NF- κ B translocation. This

study provides novel insights into calcitriol's effects on Importin α 3 and I κ B α in HepG-2 cells, unraveling cellular and molecular mechanisms in the intricate interplay between vitamin D and NAFLD pathology.

Keywords: NAFLD, vitamin D, Importin α 3, pro-inflammatory cytokines, calcitriol, inflammation.

Background

Pro-inflammatory cytokines, such as TNF- α , IL-1 β , IL-6, and IL-8, seem to play a crucial role in the progression of NAFLD as they activate the transcription factor NF- κ B. The activated NF- κ Bp50/RelA subunits are translocated to the nucleus by Importin α 3 and Importin α 4. Numerous studies have indicated a negative association between NAFLD and vitamin D levels. Low vitamin D levels have been correlated with histological severity, necro-inflammation, and fibrosis in NAFLD.

Objective

Vitamin D may resolve the ongoing inflammation observed in NAFLD by reducing the effects of pro-inflammatory cytokines that are extensively produced in NAFLD.

Methods

"HepG-2 cell cultures were subjected to calcitriol stimulation, both in the presence and absence of pro-inflammatory cytokines alongside the anti-inflammatory cytokine IL-10. Quantitative polymerase chain reaction (qPCR) was employed to assess mRNA expression levels of Importin α 3 and Importin α 4. Additionally, immunofluorescence was utilized for antibody detection."

Results

Pro-inflammatory cytokines significantly increase the mRNA and protein expression of Importin- α 3 in HepG2 cells. Treatment of HepG-2s with calcitriol significantly decreases the

mRNA and protein expression of Importin- α 3 and attenuates the effects of pro-inflammatory cytokines.

Conclusion

Our data suggest that, under inflammatory conditions, calcitriol plays an inhibitory role in the regulation of NF- κ B by decreasing the expression of Importin α 3. Thus, calcitriol reduces the translocation of NF- κ B and thereby lowers its transcriptional activities.

Introduction

Non-alcoholic fatty liver disease (NAFLD) stands as the predominant chronic liver disorder in the United States and other Western countries, impacting approximately one-third of the population (1). The surge in obesity rates has intensified the prevalence of NAFLD, particularly in obese individuals, reaching up to 90% in subjects undergoing bariatric surgery (2).

Histologically, NAFLD is characterized by the accumulation of hepatic fat, driven by the excessive deposition of free fatty acids (FFA) and triglycerides (TG) in over 5% of the liver parenchyma, culminating in steatosis and hepatic inflammation (3). The pathological progression often commences as simple steatosis, with 20% of cases evolving into nonalcoholic steatohepatitis (NASH), stimulating an inflammatory response. In 9% of cases, this inflammatory state progresses to fibrosis or liver cirrhosis, and in 1% of cases, hepatocellular carcinoma may manifest. NAFLD is intricately linked with metabolic syndrome (MS), encompassing diabetes, obesity, hypertension, hypercholesterolemia, and hypertriglyceridemia, solidifying its status as the hepatic manifestation of metabolic syndrome (4). Insulin resistance and associated hyperinsulinemia play pivotal roles in the development of both NAFLD and metabolic syndrome (5).

Experimental evidence has consistently demonstrated a negative correlation between NAFLD and vitamin D levels (6). Low vitamin D levels exhibit significant associations with

histological severity, necro-inflammation, and fibrosis in NAFLD, independent of age, gender, BMI, and the presence of other metabolic syndrome features (7). These findings extend to pediatric cases of NAFLD, emphasizing the robust association between NAFLD and chronic inflammation across all stages (8, 3). Inflammatory cytokines, such as TNF- α , IL-1 β , IL-6, and IL-8, are implicated in the pathogenesis of NAFLD, with positive associations observed between liver fibrosis amounts and circulating TNF- α levels in NASH patients (9-12).

The inflammatory cascade in NAFLD is closely linked to nuclear factor $\kappa\beta$ (NF- κB) activation, particularly its association with pro-inflammatory cytokines and the stimulation of Kupffer cells (13,14). The regulation of NF- κB relies on factors like I κB family proteins and Importin $\alpha 3$ and Importin $\alpha 4$ (15). Vitamin D has emerged as a key player in mitigating inflammation, as evidenced by its ability to up-regulate I $\kappa\text{B}\alpha$, an NF- κB inhibitor, and decrease pro-inflammatory NF- κB activity (16). Additionally, studies in NASH models have demonstrated the positive impact of vitamin D on hepatic inflammation, fibrosis, and insulin resistance (17). A recent investigation highlighted calcitriol's role in decreasing Importin $\alpha 3$ expression, shedding light on its potential to influence NF- κB translocation (15-18).

In this study, we provide novel insights into the effects of calcitriol on Importin $\alpha 3$, in HepG-2 cells, aiming to unravel the cellular and molecular mechanisms underlying the complex interplay between vitamin D and NAFLD pathology (19, 20).

Materials and Methods

Cell Culture

The hepatocellular carcinoma cell line (HepG-2, ATCC #: HB-8065) was acquired from the ATCC cell bank. Cells were cultured in a 25 cm² cell flask containing Eagle's Minimum

Essential Medium with L-Glutamine (EMEM), supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin. The cell cultures were maintained at 37°C in a humidified incubator with a 5% CO₂ and 95% air environment.

Cell Stimulation

All experiments were conducted using three biologically independent samples. The growth of cultured HepG-2 cells (maintained at 70-80% confluency) was arrested through serum starvation for 24 hours, achieved by replacing FBS-containing EMEM with FBS-free EMEM. Subsequently, cells were stimulated with calcitriol (100 nM) (D1530 Sigma-Aldrich, St. Louis, MO) in fresh EMEM for 24 hours. Recombinant TNF- α , IL-1 β , IL-6, IL-8, and IL-10 (Pepro Tech, Inc. NJ) were administered at a dose of 10 ng/ml. Cell viability and cytotoxicity were assessed using the trypan blue dye method. HepG-2 cells were exposed to varying cytokine concentrations (5 ng/ml, 10 ng/ml, 20 ng/ml, and 50 ng/ml) with viability assessed through cell counting. Multiple readings were obtained, and 10 ng/ml was identified as the most effective dose for HepG-2 cells.

RNA isolation, reverse transcription, and real-time PCR

RNA extraction was performed to obtain total RNA, followed by Real-Time RNA quantitation using previously established methods (21). Relative gene expression was determined by calculating differences in threshold cycles (Ct). The obtained results were normalized against the expression of GAPDH (Table 1).

Table 1. The primer sequences utilized in this analysis are as follows:

Gene	Forward primer	Reverse primer
GAPDH	5'- GGG AAG GTG AAG GTC GGA GT-3'	5'- TTG AGG TCA ATG AAG GGG TCA-3'
VDR	5'- CTT CAG GCG AAG CAT GAA GC-3'	5'- CCT TCA TCA TGC CGA TGT CC-3'

IκBα	5'-ACA CTA GAA AAC TTC AGA TGC-3'	5'- ACA CAG TCA TCA TAG GGC AG-3'
Importin α3	5'- TGT GAG CAA GCA GTG TGG GCA-3'	5'- TGG TGG TGG GTC TTT GTG GCG-3'
Cyp27B1	5'- TGG CCC AGA TCC TAA CAC ATT T-3'	5'- GTC CGG GTC TTG GGT CTA ACT-3'
Cyp24A1	5'- TGT GAG CAA GCA GTG TGG GCA-3'	5'- TGG TGG TGG GTC TTT GTG GCG-3'

Immunofluorescence (IF)

Immunofluorescence staining was conducted to assess the expression of VDR, IκBα, and importin α3, with a negative control that omitted the primary antibody. Briefly, cells were cultured in a T25 flask, and 30,000 cells were plated in a chamber (4 chambers per slide). Subsequently, fixation was performed using 4% formalin for 10 minutes, followed by washing and the addition of 0.1% Triton for 15 minutes. After washing, cells were subjected to a blocking solution for 30 minutes (5% fetal bovine serum in PBS). The subsequent step involved incubation with the primary antibody for 2 hours at room temperature or overnight at 4°C.

Following primary antibody incubation, cells underwent three washes with PBS, each lasting five minutes. Subsequently, cells were incubated with the corresponding secondary antibody (Alexa Fluor 594 and 488 at a 1:1000 dilution) for 30 minutes, followed by additional washes. Counterstaining with DAPI (4,6-diamidino-2 phenyl indole) was performed. Immunofluorescence microscopy was conducted using an Olympus inverted fluorescent microscope (Olympus BX51). Image J software was employed to measure immunofluorescence intensity for stained slides. Ten cells per slide were randomly selected for fluorescence intensity measurement, and the mean fluorescence intensity for each gene of interest was calculated. Each experiment was independently repeated three times, utilizing distinct batches of cells.

Statistical Analysis

The data are presented as Mean \pm Standard Deviation (SD). Statistical analyses were conducted using GraphPad Prism 4.0 biochemical statistical software (GraphPad Software, Inc., San Diego, CA) for both data analysis and graph plotting. One-way Analysis of Variance (ANOVA) was employed to identify statistically significant differences among groups. Post-hoc tests, including Dunnett or Bonferroni's test, were subsequently applied for pairwise comparisons between groups. A significance threshold of $p < 0.05$ (* $p < 0.05$, *** $p < 0.001$, **** $p < 0.0001$) was established to determine statistical significance.

Results

1. Modulatory Effects of TNF- α and Calcitriol in the Regulation of Importin $\alpha 3$

Expression

We further tested if the importin $\alpha 3$ expression was modulated by pro-inflammatory cytokines. HepG2 cells were cultured and treated with 10 ng/ml of TNF- α for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. After treatment, cells were washed and fixed for immunofluorescence. The immunofluorescence staining of Importin $\alpha 3$ expression in HepG-2 cells treated with 10 ng/ml TNF- α showed the highest immunoreactivity. HepG-2 cells treated with 100 nM calcitriol exhibited the lowest immune sensitivity for Importin $\alpha 3$ gene. Importin $\alpha 3$ showed cytoplasmic expression. *Immunofluorescence and RT-PCR analyses demonstrated that TNF- α significantly increased Importin $\alpha 3$ expression, whereas calcitriol treatment markedly reduced this effect (Figure 1).*




Figure 1: Immunofluorescence staining for the effects of TNF- α on importin α 3 in HepG-2.

The RT-PCR analysis revealed that unstimulated HepG-2 cells express Importin α 3. Treatment with 10 ng of TNF- α for 24 hours significantly increased mRNA expression of Importin α 3 ($P < 0.05$). Addition of 25 and 100 nM of calcitriol showed a significant decrease ($P < 0.05$) in mRNA expression of Importin α 3 in HepG-2. Calcitriol also showed a reduction in mRNA expression of Importin α 3, and concomitantly, calcitriol-treated cells exhibited higher immunoreactivity.

Immunofluorescence studies were conducted to assess the expression of importin α 3 in HepG-2 cells following treatment with TNF- α , with or without concurrent administration of calcitriol. Alexa Fluor 488 (green) served as the secondary antibody. The expression of importin α 3 is visualized in images A (untreated HepG-2), D (HepG-2 treated with TNF- α), G (HepG-2 treated with TNF- α and calcitriol), and J (HepG-2 treated with calcitriol only). Nuclei were counterstained with DAPI (B, E, H, and K). Merged images of Alexa Fluor and DAPI are presented in images C, F, I, and L. This figure represents a typical image from

three independent experiments. M and N show RT-PCR analysis demonstrating the effects of TNF- α and Vit D on importin $\alpha 3$ gene expression. HepG-2 cells underwent 24-hour serum starvation followed by a 24-hour treatment with TNF- α in the presence and absence of different doses of calcitriol (25-100 nm). The extracted RNA was analyzed via RT-PCR to evaluate importin $\alpha 3$ expression. The figure illustrates the fold change in mRNA expression of importin $\alpha 3$. Results are presented as mean \pm SD from three individual samples, with statistical significance indicated by $P < 0.05$. *The quantitative RT-PCR data further confirmed these findings, showing a dose-dependent reduction in Importin $\alpha 3$ mRNA expression following calcitriol treatment (Figure 2).*




Figure 2: Immunofluorescence staining for effect of IL-1 β and vitamin D on Importin $\alpha 3$ in HepG-2 cells.

2. Effect of IL-1 β and Vitamin D on Importin $\alpha 3$

We further examined the effects of IL-1 β and Vitamin D on Importin $\alpha 3$ gene expression. HepG2 cells were cultured and exposed to 10 ng/ml of IL-1 β for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. Following treatment, cells were washed and fixed for

immunofluorescence staining. The immunofluorescence results revealed the lowest immunoreactivity for importin $\alpha 3$ expression in HepG-2 cells treated with 100 nM/ml calcitriol, while the highest immunoreactivity was observed in HepG-2 cells treated with 10 ng/ml IL-1 β . Importin $\alpha 3$ exhibited cytoplasmic expression.

The RT-PCR analysis demonstrated that unstimulated HepG-2 cells express Importin $\alpha 3$. Treatment with 10 ng IL-1 β for 24 hours significantly increased mRNA expression of importin $\alpha 3$ ($P < 0.05$). The addition of 25 and 100 nM calcitriol significantly ($P < 0.05$) decreased mRNA expression of Importin $\alpha 3$ in HepG-2 cells. Calcitriol was shown to reduce mRNA expression of importin $\alpha 3$ in HepG-2 cells. Our data suggest that IL-1 β induces an increase, while calcitriol results in a decrease in Importin $\alpha 3$ expression in HepG-2 cells.

Immunofluorescence studies were conducted to assess the expression of I κ B α in HepG-2 cells after treating cells with IL-1 β , with or without calcitriol. Alexa Fluor 488 (green) served as the secondary antibody. The expression of I κ B α is depicted in [image A (untreated HepG-2), D (HepG-2 treated with IL-1 β), G (HepG-2 treated with IL-1 β and calcitriol), and J (HepG-2 treated with calcitriol only)]. DAPI (B, E, H, and K) was used to counterstain the nucleus. Images C, F, I, and L represent the merged images of Alexa Fluor and DAPI. This is a representative image from three separate samples. RT-PCR analysis was performed to investigate the effects of IL-1 β and Vitamin D on Importin $\alpha 3$ in HepG-2 cells (M and N). The cells were subjected to serum starvation for 24 hours, followed by treatment with IL-1 β in the presence and absence of different doses of calcitriol (25-100 nM) for an additional 24 hours. The extracted RNA was then analyzed using RT-PCR to assess importin $\alpha 3$ gene expression. The figure illustrates the fold change in mRNA expression of Importin $\alpha 3$. Data are presented as mean \pm SD from three individual experimental samples; $P < 0.05$. ***IL-1 β treatment significantly enhanced Importin $\alpha 3$ expression at both protein and mRNA levels, while calcitriol markedly***

suppressed this response (Figure 3).




Figure 3: Immunofluorescence staining for the effects of IL-6 on Importin α 3 in HepG-2.

3. Effect of IL-6 and Vitamin D on Importin α 3

We further tested if IL-6 modulated Importin α 3 gene expression. HepG2 cells were cultured and exposed to 10 ng/ml of IL-6 for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. Subsequently, cells were subjected to washing and fixation for immunofluorescence staining. The immunofluorescence results revealed the highest immunoreactivity for importin α 3 in HepG-2 cells treated with 10 ng/ml IL-6, while the lowest immunoreactivity was observed in HepG-2 cells treated with 100 nM calcitriol. Importin α 3 exhibited cytoplasmic expression. The RT-PCR analysis demonstrated that unstimulated HepG-2 cells express Importin α 3; treatment with 10 ng IL-6 for 24 hours led to an insignificant increase in mRNA expression of importin α 3 ($P > 0.05$). The addition of 25 or 100 nM calcitriol with 10 ng IL-6 showed a statistically insignificant ($P > 0.05$) decrease in mRNA expression of Importin α 3 in HepG-2 cells. Calcitriol, as indicated, reduces the

mRNA expression of importin $\alpha 3$, and concomitantly, calcitriol treatment resulted in higher immunoreactivity. These findings suggest that calcitriol reduces the expression of Importin $\alpha 3$ in HepG-2 cells. Together our data suggest that IL-6 induces an increase, while calcitriol results in a decrease in Importin $\alpha 3$ expression in HepG-2 cells.

Immunofluorescence studies were conducted to assess the expression of I κ B α in HepG-2 cells following treatment with IL-6, with or without calcitriol. Alexa Fluor 488 (green) served as the secondary antibody. The expression of I κ B α is illustrated in [image A (untreated HepG-2), D (HepG-2 treated with IL-6), G (HepG-2 treated with IL-6 and calcitriol), and J (HepG-2 treated with calcitriol only)]. DAPI (B, E, H, and K) was used to counterstain the nucleus. Images C, F, I, and L represent the merged images of Alexa Fluor and DAPI. This serves as a representative image from three separate experiments. RT-PCR analysis was conducted to investigate the effects of IL-6 and Vitamin D on Importin $\alpha 3$ (M and N). HepG-2 cells were serum-starved for 24 hours, followed by treatment with IL-6 in the presence and absence of different doses of calcitriol (25-100 nM) for an additional 24 hours. The extracted RNA was subjected to RT-PCR for Importin $\alpha 3$ expression. The figure illustrates the fold change in mRNA expression of Importin $\alpha 3$. Data are presented as mean \pm SD from three individual samples; $P < 0.05$. ***IL-6 moderately increased Importin $\alpha 3$ expression, which was attenuated by calcitriol treatment, as demonstrated by immunofluorescence and RT-PCR analyses (Figure 4).***




Figure 4: Immunofluorescence staining for the effects of IL-8 on importin α3 in HepG-2.

4. Effect of IL-8 and Vitamin D on Importin α3

HepG2 cells were cultured and treated with 10 ng/ml of IL-8 for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. After treatment, cells were washed and fixed for immunofluorescence. Immunofluorescence staining showed the highest immunoreactivity for importin α3 in HepG2 cells treated with 10 ng/ml IL-8, with the lowest observed in HepG2 cells treated with 100 nM calcitriol. Importin α3 exhibited cytoplasmic expression.

The RT-PCR analysis demonstrated that unstimulated HepG-2 cells express Importin α3. Treatment with 10 ng IL-18 for 24 hours significantly increased the mRNA expression of Importin α3 ($P < 0.05$). The addition of 25 or 100 nM of calcitriol significantly ($P < 0.05$) decreased the mRNA expression of Importin α3 in HepG-2. Calcitriol has shown the capability to reduce the mRNA expression of Importin α3 in HepG-2 cells. These data

suggest that calcitriol, by reducing the expression of Importin $\alpha 3$, diminishes the pro-inflammatory effects of IL-8 in HepG-2. Our data show that IL-8 increases Importin $\alpha 3$ expression where as calcitriol decreases.

Immunofluorescence studies were conducted to assess the expression of importin $\alpha 3$ in HepG-2 cells following treatment with IL-8, with or without calcitriol. Alexa Fluor 488 (green) served as the secondary antibody. The expression of Importin $\alpha 3$ is depicted in Image A (untreated HepG-2), D (HepG-2 treated with IL-8), G (HepG-2 treated with IL-8 and calcitriol), and J (HepG-2 treated with calcitriol only). DAPI (B, E H, and K) was used to counterstain the nucleus. Images C, F, I, and L represent the merged images of Alexa Fluor and DAPI. This image is a representative snapshot of three distinct experiments. RT-PCR analysis was performed to investigate the effects of IL-8 and Vitamin D on Importin $\alpha 3$ (M and N). HepG-2 cells were serum-starved for 24 hours, followed by treatment with IL-8 in the presence and absence of different doses of calcitriol (25-100 nM) for 24 hours. The extracted RNA was subjected to RT-PCR to assess the expression of Importin $\alpha 3$. The figure illustrates the fold change in mRNA expression of Importin $\alpha 3$. Data are presented as mean \pm SD from three individual experimental samples; $P < 0.05$. ***IL-8 significantly upregulated Importin $\alpha 3$ expression, whereas calcitriol treatment reversed this effect at both transcriptional and protein levels (Figure 5).***




Figure 5: Immunofluorescence staining for effect of IL-10 and vitamin

5. Effect of IL-10 and Vitamin D on Importin α 3

IL-10, an anti-inflammatory cytokine mainly produced by the liver, regulates inflammation in various organs (19). In the liver, IL-10 eases hepatocellular injury by inhibiting T-cells, monocytes, and macrophages (20). Its role in NAFLD's pathogenesis, insulin resistance, and progression is unclear. A study on mice lacking IL-10, fed a high-fat diet, suggested that IL-10 prevents hepatic steatosis but isn't linked to insulin resistance (21). Another study using IL-10 interventions in an NAFLD mouse model increased pro-inflammatory cytokines and reduced insulin signal transduction and steatosis (22). The role of IL-10 in preventing metabolic syndromes, including NAFLD, shows mixed results across studies, raising critical questions about its effectiveness in different population.

Further investigations were conducted to examine if IL-10 and calcitriol had modulated Importin α 3 expression. HepG2 cells were cultured and subjected to a 24-hour treatment with 10 ng/ml of IL-10 in the presence and absence of 25 or 100 ng/ml calcitriol. Following

treatment, cells were washed and fixed for immunofluorescence staining. HepG2 cells treated with 10 ng/ml IL-10 and 100 nM calcitriol exhibited the lowest immunoreactivity for importin $\alpha 3$ protein expression compared to the control. Importin $\alpha 3$ demonstrated a cytoplasmic expression pattern.

RT-PCR analysis of unstimulated HepG-2 cells revealed the expression of importin $\alpha 3$ mRNA. Treatment with IL-10 (10 ng/ml) significantly decreased the mRNA expression of importin $\alpha 3$ ($P < 0.05$). Similarly, calcitriol significantly ($P < 0.05$) reduced the mRNA expression of importin $\alpha 3$ in HepG-2 cells. Notably, this study highlights how calcitriol augments the anti-inflammatory effects of IL-10 by reducing the mRNA expression of importin $\alpha 3$. Our data indicate that IL-10 and calcitriol collectively diminish importin $\alpha 3$ mRNA expression in HepG-2 cells.

D on Importin $\alpha 3$ in HepG-2 cells. Immunofluorescence studies were conducted to evaluate the expression of Importin $\alpha 3$ in HepG-2 cells following treatment with IL-10, with or without calcitriol. A secondary antibody, Alexa Fluor 488 (green), was utilized. The expression of Importin $\alpha 3$ is illustrated in Image A (untreated HepG-2), D (HepG-2 treated with IL-10), G (HepG-2 treated with IL-10 and calcitriol), and J (HepG-2 treated with calcitriol only). DAPI (B, E, H, and K) was employed to counterstain the nucleus. Images C, F, I, and L depict the merged images of Alexa Fluor and DAPI. This representation is derived from three distinct samples. RT-PCR analysis was conducted to investigate the impact of IL-10 and Vitamin D on importin $\alpha 3$ (M and N). HepG-2 cells were subjected to a 24-hour serum starvation period, followed by treatment with IL-10 in the presence and absence of different doses of calcitriol (25-100 nM) for an additional 24 hours. Subsequently, RNA was extracted and subjected to RT-PCR to assess the gene expression of importin $\alpha 3$. The figure illustrates the fold change in mRNA expression of Importin $\alpha 3$. The data are presented as mean \pm SD from three individual experimental samples; $P < 0.05$. *IL-10 alone and in*

combination with calcitriol significantly reduced *Importin $\alpha 3$* expression in HepG-2 cells (Figure 6).




Figure 6: Immunofluorescence staining of the effects of TNF- α on I κ B α in HepG-2.

6. Effect of TNF- α and Vitamin D on I κ B α expression

We have further assessed the effects of pro-inflammatory cytokine TNF- α on I κ B α expression. HepG2 cells were cultured and treated with 10 ng/ml of TNF- α for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. After treatment, cells were washed and fixed for immunofluorescence. Accordingly, immunofluorescence staining showed the highest immunoreactivity of I κ B α in calcitriol-treated HepG-2 cells. HepG-2 cells treated with 10 ng/ml TNF- α showed the lowest I κ B α expression. I κ B α exhibited cytoplasmic expression in HepG-2 cells.

To further explore the role of calcitriol in the regulation of the effects of TNF- α on I κ B α , RT-PCR was conducted. Unstimulated HepG-2 cells exhibited mRNA expression of

I κ B α . Following TNF- α (10 ng/L) treatment, there was a significant decrease in mRNA expression of I κ B α in HepG-2 cells. The addition of calcitriol at concentrations of 25-100 nM significantly ($P < 0.05$) increased mRNA expression of I κ B α . These results suggest that calcitriol is capable of up-regulating the transcription of I κ B α in HepG-2 cells and attenuating the effects of TNF- α on I κ B α gene expression in HepG-2 cells.

Immunofluorescence studies were conducted to examine the expression of I κ B α in HepG-2 cells after treating the cells with TNF- α with or without calcitriol. Alexa Fluor 594 (red) was used as the secondary antibody. The expression of I κ B α is depicted in [image A (untreated HepG-2), D (HepG-2 treated with TNF- α), G (HepG-2 treated with TNF- α and calcitriol), and J (HepG-2 treated with calcitriol only)]. DAPI (B, E, H, and K) was used to counterstain the nucleus. Images C, F, I, and L represent the merged images of Alexa Fluor and DAPI. This is a representative image from three separate experiments. RT-PCR analysis for the effects of TNF- α and Vitamin D on I κ B α expression (M and N): HepG-2 cells were serum starved for 24 hour followed by treatment with TNF- α in the presence and absence of different doses calcitriol (25-100 nm) for 24 hours. The extracted RNA was subjected to RT-PCR for I κ B α expression. Figure shows fold change in mRNA expression of I κ B α . Data are shown as mean \pm SD from three individual samples; $P < 0.05$. ***TNF- α markedly suppressed I κ B α expression, while calcitriol significantly restored its expression, as shown by immunofluorescence and RT-PCR analyses (Figure 7).***




Figure 7: Immunofluorescence staining for effect of IL-1 β and vitamin D on I κ B α in HepG-2 cells.

7. Effects of IL-1 β and Vitamin D on I κ B α

We further investigated the effects of IL-1 β and Vitamin D on I κ B α . HepG2 cells were cultured and treated with 10ng/ml of IL-1 β for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. After treatment, cells were washed and fixed for immunofluorescence staining. Accordingly, immunofluorescence staining for I κ B α expression in HepG-2, cells treated with 10ng/ml IL-1 β showed the lowest immunoreactivity compared to other treatments. I κ B α showed cytoplasmic expression in hepG-2.

The unstimulated HepG-2s expressed mRNA of I κ B α . Following IL-1 β (10ng/L) treatment, there was significant decrease in mRNA expression of I κ B α in HepG-2s ($P < 0.05$); addition of calcitriol (100nM) significantly ($P < 0.05$) increase mRNA expression of I κ B α in HepG-2. Our data suggest that IL-1 β decreases I κ B α expression while calcitriol increase in HepG-2 cells. Immunofluorescence studies were completed for the expression of I κ B α in HepG-2 cells after treating cells with IL-1 β with or without calcitriol. Alexa fluor 594 (red)

was used as secondary antibody. Expression of I κ B α [image A (untreated HepG-2), D (HepG-2 treated with IL-1 β), G (HepG-2 treated with IL-1 β and calcitriol) and J (HepG-2 treated with calcitriol only)]. DAPI (B, E, H and K) was used to counterstain nucleus. Images C, F, I and L are the merged images of images of Alexa Fluor and DAPI. This is a representative image of three separate samples. RT-PCR analysis for the effects of IL-1 β and Vitamin D on I κ B α (M and N). HepG-2 cells were starved for 24 hour followed by treatment with IL-1 β in the presence and absence of different doses calcitriol (25-100 nm) for 24 hours. The extracted RNA was subjected to RT-PCR for I κ B α gene expression. Figure shows fold change in mRNA expression of I κ B α . Data are shown as mean \pm SD from three individual experimental samples; $P < 0.05$. *IL-1 β treatment resulted in a significant reduction of I κ B α expression, which was reversed upon calcitriol treatment (Figure 8).*




Figure 8: Immunofluorescence staining for the effects of IL-6 on I κ B α in HepG-2.

8. Effect of IL-6 and Vitamin D on Importin α 3

Interleukin-6 (IL-6), a versatile cytokine, is released by various cells such as fibroblasts, endothelial cells, adipocytes, monocytes, and T-cells in response to acute inflammation (23).

Its intricate role in liver pathogenesis and its contribution to NAFLD development are not fully elucidated. Nevertheless, IL-6 has systemic effects, including inflammation stimulation, immune defense regulation (B- and T-cell functions), support of hematopoiesis, and oncogenesis (24).

Elevated IL-6 levels in the serum of obese individuals serve as a robust predictor for the onset of Type 2 diabetes, metabolic syndrome, and cardiovascular diseases (25, 26). Moreover, IL-6 exhibits slightly inhibitory effects on adipogenesis, impacting adiponectin production and lipolysis. In the context of NAFLD, IL-6 from visceral fat enters the liver, activating signaling transducer protein JAK/STAT3 in hepatocytes. This cascade triggers the transcription of various genes, including suppressor cytokine signaling 3 (SOCS-3), affecting insulin signaling and contributing to hepatic insulin resistance, thereby fostering NAFLD development (27). While initial studies suggested a protective mechanism of IL-6 in liver steatosis by promoting hepatocyte proliferation and mitigating oxidative stress, more recent research demonstrates a dual role.

Here we have assessed the effects of IL-6 and Vitamin D on Importin $\alpha 3$ expression. HepG2 cells were cultured and treated with 10 ng/ml of IL-6 for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. After treatment, cells were washed and fixed for immunofluorescence staining. Immunofluorescence staining showed the highest immunoreactivity for importin $\alpha 3$ in HepG-2 cells treated with 10 ng/ml IL-6, with the lowest in HepG-2 cells treated with 100 nM calcitriol. Importin $\alpha 3$ exhibited cytoplasmic expression.

The RT-PCR analysis showed that unstimulated HepG-2 cells express Importin $\alpha 3$. Treatment with 10 ng IL-6 for 24 hours insignificantly increased mRNA expression of Importin $\alpha 3$ ($P > 0.05$). The addition of 25 or 100 nM calcitriol with 10 ng IL-6 showed insignificantly ($P > 0.05$) decreased mRNA expression of Importin $\alpha 3$ in HepG-2 cells.

Calcitriol, as shown, reduces mRNA expression of Importin $\alpha 3$, along with having higher immunoreactivity. These data suggest that IL-6 upregulates Importin $\alpha 3$ expression whereas calcitriol downregulates in HepG-2 cells.

Immunofluorescence studies were conducted to analyze the expression of I κ B α in HepG-2 cells after treating the cells with IL-6 with or without calcitriol. Alexa Fluor 488 (green) was used as the secondary antibody. The expression of I κ B α is depicted in [image A (untreated HepG-2), D (HepG-2 treated with IL-6), G (HepG-2 treated with IL-6 and calcitriol), and J (HepG-2 treated with calcitriol only)]. DAPI (B, E, H, and K) was used to counterstain the nucleus. Images C, F, I, and L represent the merged images of Alexa Fluor and DAPI. This is a representative image from three separate experiments. RT-PCR analysis for the effects of IL-6 and Vitamin D on Importin $\alpha 3$ (M and N) was done. HepG-2 cells were serum-starved for 24 hours, followed by treatment with IL-6 in the presence and absence of different doses of calcitriol (25-100 nM) for 24 hours. The extracted RNA was subjected to RT-PCR for Importin $\alpha 3$ expression. The figure shows the fold change in mRNA expression of Importin $\alpha 3$. Data are presented as mean \pm SD from three individual samples; P < 0.05. ***Calcitriol counteracted IL-6-mediated modulation of I κ B α expression in HepG-2 cells (Figure 9).***




Figure 9: Immunofluorescence staining for effect of IL-8 and vitamin D on IκBα in HepG-2 cells.

9. Effect of IL-8 and Vitamin D on IκBα

IL-8, a chemokine secreted by various cells, including inflammatory and endothelial cells, primarily functions to coordinate the attraction of neutrophils within inflamed tissue (28). Despite limited data supporting its involvement in NAFLD development, recent research indicated elevated circulating levels of IL-8 in NAFLD patients, particularly in the NASH group, compared to obese or lean individuals (29). Furthermore, serum IL-8 levels were significantly higher in the NASH group compared to those with hepatic steatosis or in a healthy control group (30, 31). However, studies reported no discernible association between circulating IL-8 levels and NAFLD (32).

To assess the effects of IL-8 and Vitamin D on IκBα, HepG2 cells were cultured and treated with 10ng/ml of IL-8 for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. After treatment, cells were washed and fixed for immunofluorescence.

Accordingly, HepG-2 cells treated with calcitriol shown the highest immunoreactivity.

HepG-2 cells treated with 10ng/ml IL-8, the control, and 10ng/ml IL-8+100 nM shown the similar immunoreactivity. I κ B α showed cytoplasmic expression in HepG-2.

To further explore the role of calcitriol in the regulation of the effects of IL-8 on I κ B α gene expression, RT-PCR was done, and the analysis shown unstimulated HepG-2s expressed mRNA of I κ B α . Following IL-8 (10ng/L) treatment, there was insignificantly decreased in mRNA expression of I κ B α in HepG-2s ($P > 0.05$); adding of calcitriol (100nM) significantly ($P < 0.05$) increase mRNA expression of I κ B α . These data indicate that IL-8 has a limited effect on the suppression of I κ B α gene expression in HepG-2. Together our data indicate that IL-8 decreases whereas calcitriol increases I κ B α expression in HepG-2.

Immunofluorescence studies were completed for the expression of I κ B α in HepG-2 cells after treating cells with IL-8 with or without calcitriol. Alexa fluor 594 (red) was used as secondary antibody. Expression of I κ B α [image A (untreated HepG-2), D (HepG-2 treated with IL-8), G (HepG-2 treated with IL-8 and calcitriol) and J (HepG-2 treated with calcitriol only)]. DAPI (B, E, H and K) was used to counterstain nucleus. Images C, F, I and L are the merged images of Alexa Fluor and DAPI. This is a representative image of three separate experiments. RT-PCR analysis for the effects of IL-8 and Vitamin D on I κ B α HepG-2 cells were serum starved for 24 hour followed by treatment with IL-8 in the presence and absence of different doses calcitriol (25-100 nM) for 24 hours. The extracted RNA was subjected to RT-PCR for I κ B α expression. Figure shows fold change in mRNA expression of I κ B α . Data are shown as mean \pm SD from three individual experimental samples; $P < 0.05$. ***IL-8 exerted minimal suppressive effects on I κ B α expression, whereas calcitriol significantly enhanced I κ B α levels (Figure 10).***




Figure 10: Immunofluorescence staining for effect of IL-10 and vitamin D on IκBα in HepG-2 cells.

10. Effect of IL-10 and Vitamin D on IκBα

HepG2 cells were cultured and treated with 10ng/ml of IL-10 for 24 hours in the presence and absence of 25 or 100 ng/ml calcitriol. After treatment, cells were washed and fixed for immunofluorescence. Accordingly, immunofluorescence staining of HepG-2 cells treated with 10ng/ml IL-10 showed the highest immunoreactivity expression of IκBα, followed by IL-10 + 100nM. IκBα showed cytoplasmic expression in hepG-2.

RT-PCR analysis of unstimulated HepG-2 cells shown expression of IκBα mRNA; treatment with IL-10 (10ng/ml) insignificantly increased mRNA expression of IκBα; adding 25 nM calcitriol to 10ng/ml IL-10 significantly ($P < 0.05$) increased the mRNA expression of IκBα in HepG-2. It is protrusive in this study how calcitriol enhances the anti-inflammatory effects IL-10 on IκBα expression.

Immunofluorescence studies were completed for the expression of IκBα in HepG-2 cells, after treating cells with IL-10 with or without calcitriol. Alexa fluor 488 (green) was used as

secondary antibody. Expression of I κ B α [image A (untreated HepG-2), D (HepG-2 treated with IL-10), G (HepG-2 treated with IL-10 and calcitriol) and J (HepG-2 treated with calcitriol only)]. DAPI (B, E, H and K) was used to counterstain nucleus. Images C, F, I and L are the merged images of Alexa Fluor and DAPI. This image is a representative of three separate samples. RT-PCR analysis for the effects of IL-10 and Vitamin D on I κ B α (M and N). HepG-2 cells were serum starved for 24 hours followed by treatment with IL-10 in the presence and absence of different doses calcitriol (25-100 nm) for 24 hours. The extracted RNA was subjected to RT-PCR for I κ B α gene expression. Figure shows fold change in mRNA expression of I κ B α . *Data are shown as mean \pm SD from three individual experimental samples; P < 0.05. Calcitriol potentiated the anti-inflammatory effects of IL-10 by significantly increasing I κ B α expression.*

Discussion

NF- κ B, a ubiquitous cytoplasmic transcription factor, intricately governs more than 500 genes, predominantly those associated with inflammation (33). The activation cascade involves IKK activation subsequent to I κ B α degradation, culminating in NF- κ B translocation to the nucleus (17). This study unveils, for the first time, the impact of pro-inflammatory cytokines (TNF- α , IL-1 β , IL-6, IL-8) on I κ B α mRNA and protein expression in HepG-2 cells, revealing a significant reduction induced by these cytokines. Intriguingly, calcitriol, administered alone or concomitantly with these cytokines, markedly enhances I κ B α mRNA and protein expression. Our findings align with prior research illustrating calcitriol's direct role in mitigating NF- κ B activity. Specifically, in human keratinocytes, calcitriol diminished NF- κ B DNA binding activity by elevating I κ B α protein levels and curtailing IL-8 production (34). Furthermore, in pancreatic islet cells, calcitriol down-regulated pro-inflammatory chemokine production while augmenting I κ B α expression, ultimately attenuating NF- κ B translational activity (35). Given the limited understanding of the sustained inflammation

implicated in NAFLD development, our observed mitigating effects of calcitriol on proinflammatory cytokines activating NF- κ B are consequential. The observed upregulation of I κ B α expression is particularly noteworthy, as it curtails the nuclear translocation of the NF- κ B transcription factor (36).

In this study, we present, for the first time, the impact of pro-inflammatory cytokines (TNF- α , IL-1 β , IL-6, IL-8) in the presence and absence of calcitriol on both mRNA and protein expression of Importin α 3 in HepG-2 cells. Notably, these proinflammatory cytokines induced a significant increase in both mRNA and protein levels of Importin α 3 (37). Our findings reveal that calcitriol exerts a down-regulatory effect on the mRNA and protein expression of Importin α 3. Importin α 3, a constituent of the nuclear pore complex and harboring the nuclear localization sequence akin to ReLA and P50 subunits of NF- κ B, plays a pivotal role in NF- κ B stimulatory activities (15). A previous study from our laboratory has established the role of calcitriol in diminishing the mRNA expression of Importin α 3 in Primary Human Bronchial Smooth Muscle Cells (38). Knockdown of Importin α 3 in these cells attenuated TNF- α -induced expression of ReLA. Consequently, calcitriol, through the reduction of Importin α 3 expression, inhibits the translocation of stimulated ReLA from the cytoplasm to the nucleus in HepG-2 cells.

Conclusion

In conclusion, our study sheds light on the intricate regulatory mechanisms involving NF- κ B, revealing its susceptibility to modulation by pro-inflammatory cytokines and the regulatory role of calcitriol. The observed reduction in I κ B α expression induced by cytokines underscores their potent inflammatory impact, while calcitriol demonstrates a mitigating effect. Additionally, our novel findings elucidate the influence of pro-inflammatory cytokines on Importin α 3 in HepG-2 cells, with calcitriol exerting a down-regulatory influence. This

signifies a potential therapeutic avenue for addressing NF- κ B-mediated inflammation, the full spectrum of these regulatory pathways.

References

1. Sung KC, Kim SH. Interrelationship between fatty liver and insulin resistance in the development of type 2 diabetes. *J Clin Endocrinol Metab.* 2011;96(4):1093–1097.
2. Boza C, et al. Predictors of non-alcoholic steatohepatitis in obese patients undergoing gastric bypass. *Obes Surg.* 2005;15(8):1148–1153.
3. Tarantino G. Hepatic steatosis, low-grade chronic inflammation and hormone/growth factor/adipokine imbalance. *World J Gastroenterol.* 2010;16(38):4773–4783.
4. Boppidi H. Non-alcoholic fatty liver disease: hepatic manifestation of obesity and the metabolic syndrome. *Postgrad Med.* 2008;120(2):E01–E07.
5. Oh MS, et al. Associations among the degree of non-alcoholic fatty liver disease, metabolic syndrome, degree of obesity in children, and parental obesity. *Pediatr Gastroenterol Hepatol Nutr.* 2016;19(3):199–206.
6. Targher G, et al. Associations between serum 25-hydroxyvitamin D3 concentrations and liver histology in patients with non-alcoholic fatty liver disease. *Nutr Metab Cardiovasc Dis.* 2007;17(7):517–524.
7. Younossi ZM, et al. Changes in the prevalence of the most common causes of chronic liver diseases in the United States from 1988 to 2008. *Clin Gastroenterol Hepatol.* 2011;9(6):524–530.e1.
8. Manco M, Ciampalini P, Nobili V. Low levels of 25-hydroxyvitamin D3 in children with biopsy-proven non-alcoholic fatty liver disease. *Hepatology.* 2010;51(6):2229.
9. Alkhoury N, et al. The inflamed liver and atherosclerosis: a link between histologic severity of non-alcoholic fatty liver disease and increased cardiovascular risk. *Dig Dis Sci.* 2010;55(9):2644–2650.

10. Day CP. Non-alcoholic fatty liver disease: a massive problem. *Clin Med (Lond)*. 2011;11(2):176–178.
11. Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science*. 1993;259(5091):87–91.
12. Xu A, et al. The fat-derived hormone adiponectin alleviates alcoholic and non-alcoholic fatty liver diseases in mice. *J Clin Invest*. 2003;112(1):91–100.
13. Stienstra R, et al. Kupffer cells promote hepatic steatosis via interleukin-1 β -dependent suppression of peroxisome proliferator-activated receptor α activity. *Hepatology*. 2010;51(2):511–522.
14. Subramaniyan V, Lubau NSA, Mukerjee N, Kumarasamy V. Alcohol-induced liver injury in signalling pathways and curcumin's therapeutic potential. *Toxicol Rep*. 2023;11:355–367.
15. Agrawal T, Gupta GK, Agrawal DK. Calcitriol decreases expression of importin α 3 and attenuates RelA translocation in human bronchial smooth muscle cells. *J Clin Immunol*. 2012;32(5):1093–1103.
16. Alexander MR, Murgai M, Moehle CW, Owens GK. Interleukin-1 β modulates smooth muscle cell phenotype to a distinct inflammatory state relative to PDGF-DD via NF- κ B-dependent mechanisms. *Physiol Genomics*. 2012;44(7):417–429.
17. Cohen-Lahav M, Shany S, Tobvin D, Chaimovitz C, Douvdevani A. Vitamin D decreases NF- κ B activity by increasing I κ B α levels. *Nephrol Dial Transplant*. 2006;21(4):889–897.
18. Fagerlund R, Melén K, Cao X, Julkunen I. NF- κ B p52, RelB and c-Rel are transported into the nucleus via a subset of importin α molecules. *Cell Signal*. 2008;20(8):1442–1451.

19. Moore KW, de Waal Malefyt R, Coffman RL, O'Garra A. Interleukin-10 and the interleukin-10 receptor. *Annu Rev Immunol.* 2001;19:683–765.
20. Dhingra S, Sharma AK, Arora RC, Slezak J, Singal PK. IL-10 attenuates TNF- α -induced NF- κ B pathway activation and cardiomyocyte apoptosis. *Cardiovasc Res.* 2009;82(1):59–66.
21. Beppu LY, et al. Tregs facilitate obesity and insulin resistance via a Blimp-1/IL-10 axis. *JCI Insight.* 2021;6(5):e140644.
22. Fontes-Cal TCM, et al. Crosstalk between plasma cytokines, inflammation, and liver damage as a new strategy to monitoring NAFLD progression. *Front Immunol.* 2021;12:708959.
23. Gwechenberger M, et al. Cardiac myocytes produce interleukin-6 in culture and in viable border zone of reperfused infarctions. *Circulation.* 1999;99(4):546–551.
24. Kishimoto T. IL-6: from its discovery to clinical applications. *Int Immunol.* 2010;22(5):347–352.
25. Ridker PM, Rifai N, Stampfer MJ, Hennekens CH. Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. *Circulation.* 2000;101(15):1767–1772.
26. Spranger J, et al. Inflammatory cytokines and the risk to develop type 2 diabetes. *Diabetes.* 2003;52(3):812–817.
27. Ueki K, Kondo T, Kahn CR. Suppressor of cytokine signaling 1 and 3 cause insulin resistance through inhibition of tyrosine phosphorylation of insulin receptor substrate proteins. *Mol Cell Biol.* 2004;24(12):5434–5446.
28. Bickel M. The role of interleukin-8 in inflammation and mechanisms of regulation. *J Periodontol.* 1993;64(5 Suppl):456–460.

29. Bahcecioglu IH, et al. Levels of serum hyaluronic acid, TNF- α and IL-8 in patients with non-alcoholic steatohepatitis. *Hepatogastroenterology*. 2005;52(65):1549–1553.
30. Torer N, Ozenirler S, Yucel A, Bukan N, Erdem O. Importance of cytokines, oxidative stress and expression of BCL-2 in the pathogenesis of non-alcoholic steatohepatitis. *Scand J Gastroenterol*. 2007;42(9):1095–1101.
31. Seo YY, et al. Tumor necrosis factor- α as a predictor for the development of non-alcoholic fatty liver disease: a 4-year follow-up study. *Endocrinol Metab (Seoul)*. 2013;28(1):41–45.
32. Duan Y, et al. Association of inflammatory cytokines with non-alcoholic fatty liver disease. *Front Immunol*. 2022;13:880298.
33. Gupta SC, Sundaram C, Reuter S, Aggarwal BB. Inhibiting NF- κ B activation by small molecules as a therapeutic strategy. *Biochim Biophys Acta*. 2010;1799(10–12):775–787.
34. Janjetovic Z, et al. 20-Hydroxycholecalciferol decreases NF- κ B activity by increasing I κ B α levels in human keratinocytes. *PLoS One*. 2009;4(4):e5988.
35. Ebert R, et al. Down-regulation by nuclear factor κ B of human 25-hydroxyvitamin D3 1 α -hydroxylase promoter. *Mol Endocrinol*. 2004;18(10):2440–2450.
36. Tse AKW, et al. 1,25-Dihydroxyvitamin D3 induces biphasic NF- κ B responses during HL-60 leukaemia cell differentiation. *Exp Cell Res*. 2007;313(8):1722–1734.
37. Zhuo Y, et al. African swine fever virus MGF360-12L inhibits type I interferon production by blocking importin α -NF- κ B signaling. *Virology*. 2021;36(2):176–186.
38. Diaz C, Thankam FG, Agrawal DK. Karyopherins in the remodelling of extracellular matrix: implications in tendon injury. *J Orthop Sports Med*. 2023;5:1–12.