

Legume-Induced Frothy Bloat in Tropical Cattle: Pathophysiology, Consequences, and Preventative Measures

Md. Emran Hossain¹, ASM Lutful Ahasan^{2,*}

Abstract

Legume-induced bloat in tropical cattle is a complex and significant health challenge that affects livestock productivity and welfare. This review aims to elucidate the multifaceted causes of bloat associated with the consumption of leguminous forages, highlighting both the physiological mechanisms and environmental factors that contribute to this condition. Key factors include the high soluble protein content of legumes, rapid fermentation rates in the rumen, and the formation of stable foam, which obstructs normal gas expulsion. Additionally, environmental conditions, such as rainfall and pasture management practices, exacerbate the risk of bloat in grazing systems. The consequences of bloat are profound, leading to reduced feed intake, impaired nutrient absorption, and, in severe cases, death. The economic implications for farmers are substantial, resulting in increased veterinary costs and decreased milk and meat production. To mitigate the risk of bloat, this review discusses various preventative measures that can be implemented in tropical cattle management systems. These include strategic feeding practices, such as providing dry forages before grazing on legumes, incorporating a balanced mix of grasses and legumes, and utilizing feed additives like poloxalene and ionophores. Furthermore, grazing management strategies, including rotational grazing and monitoring pasture conditions, play a crucial role in reducing bloat incidence. By understanding the underlying causes and consequences of legume-induced bloat, as well as implementing effective preventative strategies, cattle producers can enhance animal welfare and improve the sustainability of livestock production systems in tropical environments. This review serves as a comprehensive resource for researchers, veterinarians, and livestock managers seeking to address the challenges posed by legume-induced bloat in cattle.

Keywords: Bloat, causes, cattle, consequences, legume, preventive measures

*Author for Correspondence

ASM Lutful Ahasan

E-mail: lutful.ahasan@cvasu.ac.bd

¹Professor, Department of Animal Science and Nutrition, Faculty of Veterinary Medicine, Chattogram Veterinary and Animal Sciences University, Khulshi, Chattogram, Bangladesh

²Professor, Department of Anatomy and Histology, Faculty of Veterinary Medicine, Chattogram Veterinary and Animal Sciences University, Khulshi, Chattogram, Bangladesh

Received Date: May 20, 2026

Accepted Date: July 04, 2026

Published Date: July 06, 2026

Citation: Md. Emran Hossain, ASM Lutful Ahasan. Legume-Induced Frothy Bloat in Tropical Cattle: Pathophysiology, Consequences and Preventative Measures. Research & Reviews: Journal of Veterinary Science and Technology. 2026; 15(2): 38–59p.

INTRODUCTION

Legume-induced bloat in tropical cattle is a pressing concern that significantly impacts livestock health and productivity [1]. Bloat, characterized by the excessive accumulation of gas in the rumen, can arise from the rapid fermentation of leguminous forages, leading to severe distension and, in extreme cases, death [2]. This condition is particularly prevalent in tropical regions, where high-protein legume species, such as alfalfa, clover, and various tropical legumes, are often incorporated into grazing systems to enhance animal nutrition and forage quality [1]. Despite their nutritional benefits, these legumes pose considerable risks for bloat, necessitating a comprehensive understanding of the underlying mechanisms and effective management strategies [1].

The susceptibility of legumes to inducing bloat can be attributed to several physiological and biochemical factors [3, 4]. High concentrations of soluble proteins and non-protein nitrogen in legumes lead to rapid fermentation, resulting in the production of gases like carbon dioxide and methane. In addition, the structural properties of leguminous forages contribute to foam formation, preventing normal gas expulsion through eructation [2]. This is compounded by the tropical environment, where factors – such as high humidity and temperature – can exacerbate the risks associated with bloat [5]. The interaction between these variables creates a complex scenario for cattle producers, particularly in tropical regions where legume-based pastures are commonly utilized.

The consequences of bloat extend beyond immediate health concerns, with significant economic implications for livestock producers. Cattle experiencing bloat may exhibit reduced feed intake, decreased weight gain, and impaired reproductive performance, ultimately impacting overall herd productivity [6, 7]. Moreover, the need for veterinary intervention in severe cases adds additional financial burdens on farmers, underscoring the importance of preventative measures. Effective management practices are essential to mitigate the risks associated with legume-induced bloat. These include strategic grazing management, the incorporation of fiber-rich forages, and the use of dietary supplements that inhibit foam formation in the rumen [4, 8]. Educating farmers and livestock managers on the risks and management strategies associated with legume-induced bloat is crucial for improving animal welfare and enhancing the sustainability of tropical livestock production systems.

This study offers a novel perspective on legume-induced bloat in tropical cattle by integrating recent research findings with practical management strategies tailored to the unique challenges of tropical environments. Unlike previous studies that primarily focused on bloat mechanisms, this review emphasizes a holistic approach, examining the interplay between nutritional composition, environmental factors, and grazing practices. Additionally, it identifies underexplored legume species with lower bloat risk and provides evidence-based recommendations for integrating these species into pasture systems. By bridging the gap between scientific knowledge and practical application, this study aims to enhance the sustainability and profitability of tropical livestock production while improving animal welfare.

PATHOPHYSIOLOGY OF BLOAT

Composition of Forage

High Soluble Protein Content

Legumes are rich in soluble proteins, which are rapidly broken down by rumen microorganisms. This leads to a fast release of nitrogen compounds and subsequent fermentation [9, 10]. The rapid degradation of proteins results in a surge of gas, primarily methane and carbon dioxide. The excess nitrogen in the rumen also increases ammonia production, contributing to metabolic imbalances. Without adequate buffering, this excess gas cannot be expelled efficiently, causing a build-up in the rumen and resulting in bloat, especially when legumes are consumed in large quantities over a short period.

Rapid Fermentation

The rapid fermentation of leguminous plants – such as alfalfa or clover – leads to quick gas production in the rumen. Compared to grasses, legumes break down more quickly due to their easily degradable components [11–13]. As fermentation accelerates, bacteria and other microorganisms proliferate, producing gases as metabolic byproducts. If gas production outpaces the cattle's ability to expel it, bloat develops. Fast fermentation also contributes to frothy bloat, as foam forms within the rumen, trapping gases and preventing the animal from belching, a natural defense against bloat.

High Saponin Levels

Legumes, particularly species like alfalfa and clover, contain saponins, compounds known for their foaming properties. These saponins stabilize foam in the rumen, making it harder for gas bubbles to combine and be expelled [14, 15]. The foam creates a layer of small gas bubbles in the rumen, leading

to frothy bloat, where cattle are unable to eructate normally. The persistence of foam can result in distension of the rumen, compressing other organs and causing discomfort or even respiratory failure. Managing legume intake can reduce the risk of saponin-induced frothy bloat.

High Nitrogen Content

Legumes are known for their nitrogen-fixing capabilities, contributing to their high nitrogen content. When consumed, this nitrogen is rapidly converted into ammonia in the rumen. Excess ammonia can disrupt the ruminal environment, leading to metabolic disturbances [16, 17]. The increased microbial activity triggered by high nitrogen levels also accelerates gas production. If the rate of gas accumulation exceeds the cattle's ability to expel it, bloat ensues. Moreover, nitrogen imbalance in the rumen can lead to inefficient digestion, further increasing the risk of bloat due to undigested food particles fermenting excessively.

High Crude Protein Content

Legumes – such as alfalfa and clover – are high in crude protein, which, while beneficial for growth and milk production, poses a risk for bloat. When leguminous plants are consumed in large quantities, the high protein content rapidly degrades in the rumen, leading to rapid gas production [18, 19]. The breakdown of proteins also releases nitrogen compounds that can form a stable foam in the rumen. This foam traps gas, preventing its expulsion through belching. As foam accumulates, the rumen swells, causing discomfort and potentially leading to severe bloat if not managed.

High Rapidly Fermentable Fiber

The fiber in legumes is highly fermentable due to its chemical composition. While fiber is essential for healthy digestion in ruminants, rapidly fermentable fiber leads to the quick production of gases, primarily methane and carbon dioxide, as a byproduct of microbial fermentation [20]. Legumes' fiber content, although lower than grasses, is broken down faster in the rumen, which can result in an overload of gases that cannot be expelled quickly enough. This phenomenon often contributes to frothy bloat, especially when legumes are consumed without adequate amounts of roughage to slow fermentation.

PHYSICAL AND STRUCTURAL CHARACTERISTICS

Cell Wall Composition

The cell walls of legumes are generally thinner than those of grass, making them more easily degradable. This rapid breakdown accelerates microbial fermentation, leading to faster gas production [20]. The reduced structural fiber in legume cell walls means that less chewing and rumination are required, which decreases saliva production. Saliva is a critical buffer that helps to prevent bloat by maintaining rumen pH and aiding in gas expulsion. Without sufficient structural fiber, the rapid fermentation of leguminous plants can overwhelm the rumen's capacity to handle gas production, resulting in bloating.

Leafy Nature of Legumes

The fine, leafy structure of legumes makes them more susceptible to frothy bloat. When legumes are chewed, their small leaf particles mix easily with the rumen fluid, creating an environment conducive to foam formation [1, 21]. The small particles allow for rapid microbial fermentation, leading to an accelerated breakdown of nutrients. As gases are produced during fermentation, they become trapped in the foam, preventing the cattle from belching. The foam also contributes to ruminal distension, compressing internal organs and potentially leading to more severe symptoms if the animal cannot release the trapped gas.

Fine Particle Size

Legumes tend to break down into finer particles compared to grasses. This fine particle size facilitates faster digestion and fermentation in the rumen, contributing to quicker gas production [1, 21]. Finer particles also promote the formation of stable foam in the rumen, leading to frothy bloat. The small particle size limits the formation of gas pockets that are easily expelled, trapping gas within the rumen.

This foam prevents cattle from belching effectively, leading to a build-up of gas. Managing the particle size of leguminous forage, such as through chopping or wilting, can help reduce the risk of bloat.

Structural Characteristics of Legumes

The physical structure of legumes plays a significant role in their propensity to cause bloat. Unlike grasses, which have a more fibrous structure that promotes slower fermentation, legumes are leafy and have thinner stems [22–24]. This structural difference allows for quicker breakdown and microbial fermentation in the rumen. The faster fermentation leads to more rapid gas production. Additionally, the finer, leafy nature of legumes makes it easier for foam to form and stabilize, trapping gas within the rumen. Cattle consuming legumes must be monitored closely, as their physical structure predisposes them to bloat.

GRAZING AND CONSUMPTION PATTERNS

Concentrated Grazing

Cattle tend to graze more intensively on lush, high-quality legumes due to their palatability. This concentrated grazing behavior leads to rapid consumption of large quantities of legumes, which the rumen cannot process efficiently all at once [25–27]. As legumes are rapidly fermented, gas production outpaces the animal's ability to expel it. This concentrated intake, particularly in pastures dominated by legumes, increases the risk of bloat, especially if there is insufficient time between grazing sessions for rumen adaptation. Careful pasture management, such as rotational grazing, can mitigate concentrated legume intake and reduce the risk of bloat.

Sudden Dietary Changes

Cattle are particularly susceptible to bloat when introduced to high-legume diets suddenly. If animals are moved from a low-legume diet to one dominated by legumes without a gradual transition, their rumen microbes cannot adjust quickly enough to handle the increased rate of fermentation [14, 28]. This abrupt dietary change leads to rapid gas production, overwhelming the animal's ability to expel it, which can cause bloat. The transition period allows the rumen microbial population to adapt to the higher soluble protein and nitrogen content of legumes, reducing the risk of fermentation-associated gas buildup.

Overgrazing on Legumes

Cattle that overgraze on legume-dominated pastures are at an increased risk of bloat. When pastures are primarily composed of legumes, cattle may consume more forage than their rumen can handle at one time, leading to rapid fermentation [29, 30]. The excess intake of legumes produces a significant amount of gas in a short period, which the cattle cannot expel fast enough. Overgrazing also reduces plant diversity in the pasture, leaving fewer fibrous plants like grasses that would normally help slow down fermentation. Rotational grazing and balanced pasture compositions can help prevent overgrazing and reduce bloat risk.

Time of Day Grazing

The time-of-day cattle graze legumes can influence their susceptibility to bloat. Early morning grazing, particularly when pastures are covered in dew, increases the moisture content of the forage, leading to more rapid fermentation in the rumen [5, 7]. Moist legumes ferment faster than dry forage, producing gas at a rate that can overwhelm the cattle's ability to expel it. Grazing management that limits legume consumption during these high-risk times, or providing alternative forages in the morning, can reduce the likelihood of bloat. Monitoring grazing patterns can significantly help prevent legume-induced bloat.

Lack of Roughage

When cattle consume a high proportion of legumes without sufficient roughage in their diet, the risk of bloat increases [2, 20, 31]. Roughage, such as grass or hay, is essential for stimulating rumination and promoting saliva production. Saliva acts as a natural buffer in the rumen, helping to neutralize acids

and facilitating gas expulsion. Without enough roughage to slow down the fermentation of legumes, gas builds up more rapidly in the rumen. Providing cattle with access to high-fiber roughage when grazing on legume-dominated pastures helps maintain rumen function and prevents the rapid fermentation that leads to bloat.

ENVIRONMENTAL AND MANAGEMENT FACTORS

Weather Conditions

Environmental conditions, particularly weather, can influence the likelihood of bloat in cattle grazing on legumes. Wet, dewy, or rainy conditions increase the moisture content of leguminous forage, making it more prone to rapid fermentation in the rumen [18]. Moist forage ferments more quickly than dry forage, leading to faster gas production. During periods of wet weather, the risk of bloat increases as cattle consume water-laden legumes that ferment at an accelerated rate. Proper grazing management, including delaying grazing after rainfall or dew, can reduce the risk of bloat during such environmental conditions. Providing alternative feed or grazing options during wet periods can also mitigate the risk of legume-induced bloat in cattle.

Warm Temperatures

Warm weather conditions can exacerbate the risk of bloat in cattle grazing on legumes. Higher temperatures accelerate the fermentation process in the rumen, leading to more rapid gas production [18]. As legumes ferment faster under warm conditions, the rate at which gas is produced can surpass the animal's ability to expel it through belching. The increased microbial activity in the rumen due to warmth also contributes to the stabilization of foam, trapping gas and leading to frothy bloat. Grazing management practices, such as providing access to shaded areas and alternative forage during hot weather, can reduce this risk.

Fluctuations in Pasture Quality

Variability in pasture quality, especially in legume-dominated pastures, can increase the likelihood of bloat [21, 32, 33]. Changes in plant maturity, moisture levels, or grazing pressure can result in inconsistent forage quality, leading to uneven fermentation rates in the rumen. For example, legumes in early growth stages tend to be more succulent and rapidly fermentable, while overmature legumes might still pose a risk if they are grazed after rain. These fluctuations make it difficult to predict when bloat might occur, necessitating careful pasture monitoring and management to balance legume intake and reduce the risk of bloat.

Poor Grazing Management

Inadequate grazing management, such as overstocking or continuous grazing of legume-rich pastures, heightens the risk of bloating in cattle. Overstocked pastures force cattle to graze more intensively, often consuming more legumes than their rumen can handle, resulting in excessive gas production [2, 5, 7]. Continuous grazing on legume-dominant pastures does not allow the rumen microbiota to adapt to changing forage conditions, leading to an increased risk of bloat during times of rapid legume growth. Implementing rotational grazing, mixing legumes with grass species, and monitoring the grazing intensity can help mitigate these risks and promote rumen health.

BIOLOGICAL AND PHYSIOLOGICAL FACTORS

Foaming Action

The proteins and saponins present in legumes contribute to the formation of stable foam in the rumen, which is a key factor in frothy bloat [22, 23]. This foam traps gas bubbles within the rumen fluid, preventing their release through eructation. Unlike free gas bloat, where gas accumulates in a large bubble that can be belched, frothy bloat is more dangerous because the gas is dispersed throughout the ruminal contents. The stability of this foam makes it difficult for cattle to expel the trapped gas, leading to ruminal distension and potentially life-threatening bloat if not addressed promptly.

Rumen Viscosity

Legumes increase the viscosity of the rumen contents, making it harder for cattle to expel gas naturally. The increased viscosity results from the breakdown of soluble proteins and other components of the legume, which can contribute to the formation of stable foam [22, 23]. This thicker rumen content can trap gas bubbles, leading to frothy bloat. High rumen viscosity also slows down rumen contractions, which are essential for moving gas toward the esophagus for expulsion. Managing the legume intake and ensuring that cattle have access to coarser forages can help reduce the viscosity and lower the risk of bloat.

Gas Trapping by Mucilage

Some legumes contain mucilage, a gelatinous substance that can trap gases in the rumen, contributing to the development of bloat [34–36]. The presence of mucilage in legumes – like alfalfa and clover – can stabilize foam and prevent gas from coalescing into larger bubbles that can be belched out. Instead, the gas remains trapped in small pockets throughout the rumen contents, leading to frothy bloat. The mucilage forms a sticky matrix that increases the difficulty of gas release, necessitating management practices – such as mixing legumes with more fibrous forages – to decrease the likelihood of foam formation.

Higher Rumen Gas Production

The fermentation process of legumes inherently produces more gas than other types of forage due to their high protein and soluble carbohydrate content. This gas, primarily methane and carbon dioxide, is a natural byproduct of microbial digestion in the rumen [31, 37]. However, the rapid fermentation of legumes can produce gas at a rate that outpaces the animal's ability to expel it through eructation. The higher rumen gas production associated with legume-rich diets, combined with the foaming action of leguminous proteins, makes cattle more susceptible to both frothy and free gas bloat, requiring careful dietary management.

Increased Microbial Activity

Legumes support rapid microbial growth and activity in the rumen, which in turn accelerates gas production. The nitrogen and protein-rich content of legumes provide an ideal environment for rumen bacteria to thrive, leading to increased fermentation rates [31, 37]. As microbial populations proliferate, they produce more gas as a byproduct of their metabolic processes. The higher levels of microbial activity can overwhelm the animal's normal mechanisms for expelling gas, leading to bloat. Managing the intake of legumes and ensuring the inclusion of slower-fermenting forages can help balance microbial activity and reduce the risk of excessive gas buildup.

Slower Rumen Contractions

Excessive gas production due to legume consumption can slow down normal rumen contractions, which are vital for moving gas to the esophagus for expulsion. These contractions, known as rumen motility, are essential for the digestion process and for preventing gas accumulation [38]. When gas builds up in the rumen due to the rapid fermentation of legumes, it can impair normal contractions, leading to a distended rumen. The distension puts pressure on other organs, potentially causing more severe symptoms of bloat such as difficulty breathing. Maintaining proper forage balance can help sustain rumen motility and prevent bloat.

Rumen pH Changes

The fermentation of legumes can lower the rumen pH slightly, creating an acidic environment that favors microbial activity and foam formation. The drop in pH occurs because the rapid fermentation of soluble carbohydrates and proteins in legumes produces volatile fatty acids (VFAs), which reduce the rumen's buffering capacity [38]. This lower pH can alter the microbial population, encouraging the growth of bacteria that produce more gas and foam. The acidic environment also impairs the normal function of rumen contractions, making it harder for cattle to expel gas. Providing adequate roughage and buffering agents can mitigate these pH-related risks.

Lack of Saliva Production

Saliva plays a crucial role in buffering the rumen and helping to prevent bloat. When cattle consume legumes, particularly in large amounts, they may not chew their cud as thoroughly as they would with more fibrous forages, resulting in less saliva production [39–41]. Saliva contains bicarbonate, which helps to neutralize the acids produced during fermentation and maintain a stable rumen pH. Without sufficient saliva, the rumen becomes more acidic, increasing the risk of frothy bloat. Ensuring that cattle have access to roughage that requires more chewing can stimulate saliva production and reduce the likelihood of legume-induced bloat.

DIETARY AND FEEDING PRACTICES

Lack of Dietary Adaptation

Introducing cattle to high-legume diets without a proper adaptation period increases the risk of bloat. The rumen microbiota requires time to adjust to the high protein and nitrogen content of legumes, and sudden dietary changes can overwhelm the system [20, 24, 42]. When cattle are transitioned too quickly to a diet dominated by legumes, the rapid fermentation of these plants leads to excessive gas production. Gradual adaptation allows the rumen microbes to adjust to the new diet and helps prevent metabolic disturbances that contribute to bloat. Implementing a slow transition to legume-rich diets can significantly reduce the risk.

Inadequate Forage Mixing

Cattle grazing in pastures that lack a balance of grasses and legumes are at higher risk of bloat. While legumes are highly nutritious, their rapid fermentation can lead to excessive gas production [20, 24, 42]. Mixing legumes with grass or other high-fiber forages slows down the fermentation process, allowing gas to be expelled more effectively. Balanced pastures provide the fiber needed to promote proper rumination and saliva production, both of which are essential for maintaining rumen health. By managing pasture composition and ensuring a diverse forage mix, farmers can reduce the risk of legume-induced bloat in cattle.

Imbalance in Fiber

Legumes tend to have a lower fiber content compared to grasses, which slows down the fermentation process. Without enough fiber, legumes ferment rapidly in the rumen, producing gas at a faster rate than the animal can expel [43]. The lack of structural fiber in legumes also means that cattle spend less time chewing their cud, which reduces saliva production and limits the buffering capacity of the rumen. Feeding cattle a diet that includes both legumes and high-fiber forages, such as hay or straw, can help balance the rate of fermentation and prevent the rapid gas buildup that leads to bloat.

METABOLIC FACTORS

High Ammonia Production

Legumes are rich in nitrogenous compounds, which convert rapidly into ammonia in the rumen. While ammonia is crucial for microbial protein synthesis, excess can be harmful. High levels of ammonia from legume consumption overwhelm the animal's natural nitrogen absorption and conversion mechanisms [44, 45]. Ammonia toxicity can cause metabolic disturbances, leading to a slower fermentation rate in the rumen and inefficient gas expulsion. The imbalance can contribute to the conditions that cause bloat, as the rumen becomes less capable of handling excessive gas. Including balanced protein and carbohydrate sources in the diet can help mitigate these effects.

Disruption of Normal Microbial Populations

Legume-rich diets can alter the balance of the microbial populations in the rumen, leading to higher levels of gas-producing bacteria. The high soluble protein content in legumes encourages the growth of specific microorganisms that thrive in nitrogen-rich environments. This altered microbial profile accelerates the fermentation of legumes, producing methane and carbon dioxide at a rate that can outpace the rumen's ability to expel gas [45–47]. The shift in microbial dynamics may also reduce the population of fiber-degrading bacteria, further contributing to bloat. Managing the dietary ratio of legumes and fiber helps maintain a stable microbial ecosystem in the rumen.

Increased VFA Production

The rapid fermentation of legumes leads to the production of volatile fatty acids (VFAs) such as acetate, propionate, and butyrate. These acids are beneficial for energy production in cattle, but their excessive accumulation can lower the rumen pH, creating an acidic environment. This drop in pH encourages foam formation and slows rumen contractions, which hinders gas expulsion [1, 5, 48]. The increase in VFAs also alters the osmotic balance in the rumen, drawing water into the rumen and compounding the bloating effect. Balancing legume intake with other forages can reduce VFA overproduction and its associated risks.

Excessive Fermentation End Products

As legumes ferment quickly, they produce more byproducts such as VFAs, ammonia, and gases like methane and carbon dioxide. These fermentation end products contribute to rumen acidification and increase the likelihood of foam formation. The excessive production of fermentation byproducts can overwhelm the natural buffering systems in the rumen, leading to bloat [35, 36]. Additionally, the high ammonia content can be toxic, impairing the efficiency of the rumen's microbial ecosystem. Feeding cattle balanced diets that combine legumes with slower-fermenting forages can help reduce the rapid accumulation of these harmful fermentation byproducts.

Rumen Dysbiosis

Legume-induced rapid fermentation can lead to an imbalance in the rumen's microbial ecosystem, known as dysbiosis. This condition occurs when the population of gas-producing bacteria proliferates while fiber-degrading microbes decline [45–47]. Dysbiosis results in inefficient digestion and increased gas production. The reduced fiber degradation slows down rumination and saliva production, both essential for preventing bloat. Once the rumen environment is altered, it becomes more difficult for cattle to recover from episodes of bloat. Preventive dietary strategies, such as gradually introducing legumes into the diet, can help maintain microbial balance and prevent dysbiosis.

RUMINAL GAS DYNAMICS

Methane Production

Legumes, due to their high protein content, encourage the production of methane in the rumen. Methane is a natural byproduct of the fermentation process, produced by methanogenic archaea in the rumen [45–47]. However, the rapid fermentation of legumes results in methane being generated at a rate that may exceed the cattle's ability to expel it through belching. Methane, when trapped in the foam of the rumen, contributes to frothy bloat. Managing legume intake and including dietary supplements, like ionophores or tannins, which reduce methane production, can help mitigate this risk and improve rumen efficiency.

Carbon Dioxide Accumulation

In addition to methane, the fermentation of legumes produces significant amounts of carbon dioxide (CO₂), which must be expelled by the animal to prevent bloating. The rapid fermentation of leguminous forages can produce CO₂ at a rate that overwhelms the rumen's natural mechanisms for expulsion [35, 36]. When CO₂ accumulates in the rumen, it becomes trapped in foam created by legume proteins and saponins, leading to frothy bloat. If the animal is unable to expel the gas effectively, pressure builds up in the rumen, causing distension. Slow introductions of legumes into the diet can prevent this rapid gas buildup.

Gas Trapping in the Rumen

One of the primary reasons legumes cause bloat is due to their tendency to trap gases in the rumen. The rapid fermentation of legume proteins and carbohydrates produces gases like methane and carbon dioxide [35, 36]. These gases can become trapped in the foam that forms in the rumen due to the presence of saponins and soluble proteins. Once trapped, the gases cannot be belched out, leading to an accumulation of pressure within the rumen. This trapped gas leads to the expansion of the rumen and, if untreated, can cause severe bloating or even death in extreme cases.

Slow Gas Release

The gases produced during the fermentation of legumes, such as methane and carbon dioxide, are typically released through belching. However, the foam produced by leguminous plants, especially in the case of alfalfa and clover, prevents the coalescence of gas bubbles into larger pockets, making it harder for cattle to expel the gas naturally [35, 36]. This slow gas release results in ruminal distension and bloat, as the trapped gas continues to build up within the rumen. Grazing practices that combine legumes with more fibrous plants can aid in promoting natural gas release and preventing bloating.

Inefficient Gas Eructation

Legumes, with their high soluble protein content, produce stable foam that prevents the efficient eructation (belching) of gas from the rumen. Normally, cattle expel gases through eructation to maintain a stable rumen environment [35, 36]. However, the foam produced by legumes traps gases, like methane and carbon dioxide, making it difficult for cattle to expel them. As gas accumulates, it increases rumen pressure, causing discomfort and potentially leading to severe bloat. Introducing anti-foaming agents, such as certain oils or surfactants, can help reduce foam formation and allow for the more efficient release of gases.

FOAM AND GAS RETENTION MECHANISMS

Stable Foam Formation

Legumes contain saponins and soluble proteins that promote the formation of stable foam in the rumen. This foam traps gas bubbles produced during fermentation, preventing them from being expelled through normal eructation. As foam continues to form, the trapped gas builds up in the rumen, leading to frothy bloat [35, 36]. This stable foam is difficult for cattle to break down naturally, making bloat a common problem when animals graze on legume-dominant pastures. Feeding practices that include anti-foaming agents or providing alternative forages alongside legumes can help reduce foam formation and prevent bloating.

Protein Foaming Action

The proteins in legumes are highly soluble, meaning they dissolve easily in the rumen and contribute to foaming action. As these proteins break down, they form a matrix that stabilizes foam within the rumen, trapping gas bubbles and preventing their release [35, 36]. This protein-induced foam creates a barrier between the gas and the esophagus, hindering the animal's ability to expel the gas through belching. The presence of protein foam is one of the primary contributors to frothy bloat, a more dangerous form of bloating compared to free gas bloat, where the gas is more easily released.

Foam Stabilization by Saponins

Saponins, naturally occurring compounds found in many legume species, have strong foaming properties that stabilize foam in the rumen. These compounds create a barrier that traps gas bubbles within the rumen fluid, leading to the development of frothy bloat [35, 36]. The stable foam formed by saponins prevents the gases produced during fermentation from escaping, leading to ruminal distension. Saponin content is particularly high in legumes, like alfalfa and clover, which are common causes of bloat in cattle. Managing the intake of saponin-rich legumes can significantly reduce the risk of bloat in grazing animals.

Gas Retention in Frothy Bloat

In frothy bloat, gas is trapped in small bubbles within the foam produced in the rumen, preventing it from being expelled. This is distinct from free gas bloat, where gas accumulates in larger pockets and can typically be belched out. Frothy bloat is more dangerous because the foam stabilizes the gas bubbles, making it difficult for cattle to release the pressure building in their rumen [35, 36]. The longer the gas remains trapped, the greater the risk of severe bloating and potentially fatal complications. Feeding practices that reduce foam formation can help prevent frothy bloat in legume-fed cattle.

Reduced Gas Bubble Coalescence

When cattle consume legumes, the gases produced in the rumen do not easily form larger bubbles due to the presence of foam, primarily caused by the soluble proteins and saponins found in these plants [35, 36]. Instead of coalescing into larger bubbles that can be expelled through eructation, the gas remains trapped in tiny bubbles within the foam. This lack of coalescence prevents the cattle from belching efficiently, leading to the accumulation of gas in the rumen and causing bloat. Feeding cattle more fibrous forage alongside legumes can encourage better gas coalescence and reduce the risk of bloat.

PREVENTATIVE MEASURES AND MANAGEMENT

Gradual Introduction to Legumes

One of the most effective ways to prevent bloat is by gradually introducing cattle to legume-rich diets. When cattle are abruptly exposed to a high proportion of legumes, their rumen microbiota have little time to adjust, which can lead to rapid fermentation and excessive gas production [1, 5, 48]. Gradual introduction allows the rumen microbes to adapt to the high levels of protein and nitrogen in legumes, reducing the risk of bloat. This can be achieved by slowly increasing the proportion of legumes in the diet over several weeks, allowing the animal's digestive system to become accustomed to the new feed.

Use of Antifoaming Agents

Antifoaming agents, such as oils or surfactants, can be added to the cattle's diet to prevent foam formation in the rumen. These agents reduce the surface tension of the ruminal contents, breaking down the foam and allowing gas to be expelled more easily through eructation [1, 5, 48]. Oils – like vegetable oil, mineral oil, or tallow – can be used effectively to prevent frothy bloat. Additionally, commercial antifoaming products are available, which can be mixed with feed or applied directly to grazing animals during periods of high legume consumption to reduce the risk of bloat.

Feeding Dry Forage Before Grazing Legumes

A preventive measure to reduce the risk of legume-induced bloat is to offer dry forage, such as hay, before cattle are turned out to graze on legume-dominated pastures. Dry forages promote rumination and saliva production, which helps buffer the rumen and prevent excessive foam formation [1, 5, 48]. The fiber in dry forages also slows down the fermentation process, allowing gas to be expelled more easily. By filling the rumen with dry forage first, cattle are less likely to consume large quantities of fresh legumes, thus reducing the likelihood of rapid gas production and bloat.

Mixing Legumes with Grasses

A well-established method for preventing bloat is to mix legumes with grasses in pastures. Grasses, being higher in fiber and lower in soluble protein, ferment more slowly in the rumen and produce less gas compared to legumes [1, 5, 48]. By planting mixed pastures with a combination of legumes and grasses, cattle are less likely to consume a large proportion of legumes at once, which can mitigate the rapid fermentation that leads to bloat. Grazing on mixed pastures also encourages more balanced intake, promoting better rumen health and reducing the likelihood of gas accumulation.

Avoid Grazing on Wet Pastures

Wet pastures, particularly those with a high legume content, can significantly increase the risk of bloat. Legumes that are wet from dew, rain, or irrigation tend to break down more rapidly in the rumen, leading to faster fermentation and more gas production. Furthermore, wet conditions can make legumes more palatable, encouraging cattle to consume them in larger quantities [1, 5, 48]. To prevent bloat, it is advisable to avoid grazing cattle on legume-rich pastures during wet conditions or to wait until the pasture has dried. Offering dry feed before grazing can also help mitigate this risk.

MANAGEMENT OF CATTLE AND GRAZING PRACTICES

Rotational Grazing

Rotational grazing is a pasture management technique that can help prevent bloat by controlling cattle intake of legumes. Instead of allowing continuous grazing on legume-rich pastures, rotational grazing

involves moving cattle between paddocks, allowing for controlled grazing times and reducing the risk of overconsumption of legumes [2, 5, 29, 30, 47]. By rotating cattle between different pastures, including those with a higher proportion of grasses, the overall intake of legumes is moderated. This practice helps maintain a balanced diet and reduces the likelihood of rapid gas production and foam formation in the rumen.

Avoid Grazing Hungry Cattle on Legumes

Allowing hungry cattle to graze on legume-dominated pastures can significantly increase the risk of bloat. When cattle are very hungry, they tend to gorge on the available forage, often consuming large amounts of legumes rapidly [1, 5, 48]. This sudden intake of high-protein, rapidly fermentable forage can overwhelm the rumen's capacity to expel gas, leading to bloat. To prevent this, it is important to offer cattle dry forage or feed before allowing them to graze on legume-rich pastures. By ensuring that cattle are not excessively hungry when they begin grazing, the risk of bloat can be minimized.

Restrict Grazing Time on Legume Pastures

Limiting the amount of time cattle spend grazing on legume-rich pastures can help reduce the risk of bloat. By controlling grazing time, farmers can prevent cattle from consuming large quantities of legumes in a short period, which would otherwise increase the likelihood of rapid fermentation and gas production [2, 5, 29, 30, 47]. Grazing periods can be restricted to a few hours a day, particularly during times when legumes are most lush and bloat-inducing such as early morning or after rain. Restricting grazing time also encourages cattle to eat more slowly, giving the rumen time to process the legumes properly.

Monitor Legume Growth Stages

The growth stage of legumes can impact their bloat potential. Legumes in their early vegetative stages contain higher levels of soluble proteins and are more likely to cause bloat, whereas more mature legumes, while still potentially bloat-inducing, pose a slightly lower risk [2, 5, 29, 30, 47]. By monitoring the growth stages of legumes and adjusting grazing practices accordingly, farmers can mitigate bloat risk. For example, grazing can be delayed until legumes reach a more mature stage, or cattle can be moved to alternate pastures when legumes are in their most bloat-inducing growth phase.

Use of Bloat Blocks or Licks

Bloat blocks or mineral licks containing anti-foaming agents can be placed in pastures to help prevent bloat in cattle grazing on legume-dominated fields. These blocks contain substances that break down foam in the rumen, allowing gas to be expelled more easily [2, 5, 29, 30, 47]. Cattle lick these blocks as they graze, which ensures they ingest small amounts of anti-foaming agents regularly. These products can be particularly useful during periods when bloat risk is high such as when legumes are in early growth stages or after rain. Ensuring cattle have access to these blocks can significantly reduce the risk of bloat.

OTHER FACTORS CONTRIBUTING TO BLOAT

Lack of Supplementation

Cattle grazing on legume-rich pastures may not receive enough supplementation of other nutrients, particularly fiber, which is crucial for proper digestion and rumen health. Legumes are high in protein but often low in fiber compared to grasses, leading to faster fermentation and increased gas production. Without adequate fiber supplementation, the rumen's natural buffering systems are compromised, leading to a higher risk of bloat. Providing supplemental hay, silage, or other fibrous feeds ensures that cattle receive a balanced diet, which helps to slow down fermentation and reduce gas buildup in the rumen.

Genetic Susceptibility

Some cattle breeds or individual animals may be genetically predisposed to developing bloat more readily than others. Factors – such as rumen size, motility, and microbial populations – can vary between

breeds, with some cattle having a higher tolerance for legume-rich diets than others. For example, some studies suggest that dairy breeds, like Holsteins, may be more prone to bloat compared to beef breeds like Angus. Selecting cattle with lower susceptibility to bloat, or monitoring those that are more at risk, can help reduce the incidence of bloat in herds grazing on legume pastures.

Stress and Disruptions in Feeding Routine

Stress and disruptions in the feeding routine can increase the risk of bloat in cattle. Sudden changes in diet, environmental stressors, or changes in the grazing schedule can disrupt normal rumen function, leading to reduced motility and increased fermentation of legumes. Stress can also reduce saliva production, which is crucial for buffering the rumen. Ensuring consistent feeding routines and minimizing stressors, such as through proper herd management and handling, can help maintain rumen health and prevent bloat. Calm environments and gradual dietary changes reduce the physiological stress that contributes to the development of bloat.

Rumen Fill Level

Cattle with a high rumen fill level are more likely to experience bloat when grazing on legumes. When the rumen is already full, there is less space for gas to accumulate, leading to increased pressure and a higher risk of frothy bloat. This is especially problematic if the cattle have recently consumed high-fiber feed or are transitioning to a legume-dominant pasture. To manage rumen fill levels, it is important to regulate grazing periods and provide balanced, moderate meals that do not overwhelm the rumen's capacity. This practice helps maintain rumen motility and reduces the risk of gas buildup.

Failure to Monitor Early Signs

Bloat can often be prevented if early signs are identified and addressed quickly. Early symptoms of bloat include distension on the left side of the animal, discomfort, restlessness, and reluctance to graze. Failure to recognize and respond to these early signs can result in the progression of bloat, which can become life-threatening if not treated promptly. Regular monitoring of cattle grazing on legume-rich pastures is essential for early detection and intervention. Timely actions, such as moving cattle to a different pasture or administering anti-foaming agents, can prevent the condition from worsening.

Use of Tannins

Tannins are plant compounds that can be added to cattle diets to reduce the risk of bloat. They work by binding to proteins in the rumen, reducing their solubility and slowing down the fermentation process. This decreases the production of gas and foam in the rumen, mitigating the risk of bloat. Tannins are naturally present in some forages and can be introduced into the diet through feed additives or by incorporating tannin-rich plants, such as certain varieties of clover or sainfoin, into pastures. When used appropriately, tannins can help create a more stable rumen environment and reduce the likelihood of excessive gas buildup in cattle grazing on legume-rich pastures.

Ionophores

Ionophores are a class of feed additives that can be used to reduce the incidence of bloat in cattle. They work by altering the rumen's microbial population, specifically by reducing the activity of bacteria that produce methane and other gases during fermentation. By shifting the balance of microbial fermentation towards pathways that produce fewer gases, ionophores help mitigate the risk of bloat. Commonly used ionophores include monensin and lasalocid, which can be added to feed to control gas production and prevent excessive foam formation. Their use is particularly helpful in cattle grazing on legume-heavy pastures.

Rumen Buffers

Rumen buffers, such as sodium bicarbonate, can be added to cattle diets to maintain a stable pH in the rumen. Legume-rich diets tend to lower rumen pH due to their high protein content and rapid fermentation, which can contribute to foam formation and bloat. By adding rumen buffers, the acidity of the rumen is neutralized, promoting a healthier fermentation process and reducing the risk of bloat.

These buffers can be administered through feed supplements, and their regular use is beneficial in herds that are consistently exposed to bloat-inducing diets such as legume-dominant pastures.

Probiotics

Probiotics are beneficial microorganisms that can be added to cattle diets to promote a healthy microbial balance in the rumen. By increasing the population of certain rumen bacteria, probiotics can help regulate fermentation processes, improving fiber digestion and reducing gas production. Probiotic supplements, such as those containing *Lactobacillus* species, can be administered to cattle to enhance the rumen's ability to process legume-rich diets without excessive gas accumulation. This approach supports overall rumen health and helps prevent the conditions that lead to frothy bloat, especially when cattle are grazing on legume-dominant pastures.

Enzyme Supplements

Enzyme supplements, such as those that target specific fiber or protein fractions, can be used to aid digestion and reduce the risk of bloat. Enzymes break down complex feed components more efficiently, leading to a more controlled fermentation process in the rumen. By enhancing the breakdown of fibrous components in the diet, enzyme supplements reduce the likelihood of gas buildup from rapid fermentation of legumes. They can be particularly useful when cattle are transitioning to legume-rich diets or during periods when bloat risk is high such as during early spring or after rainfall.

IMPACT OF SEASONAL AND ENVIRONMENTAL CONDITIONS

Spring Pastures

In spring, pastures tend to have lush growth, particularly of legumes, such as clover and alfalfa, which can increase the risk of bloat in grazing cattle. The high moisture content, combined with the rapid regrowth of legumes, promotes a fermentation environment in the rumen that is conducive to frothy bloat [18]. To mitigate this risk, it is essential to manage spring grazing carefully, including allowing cattle to graze only for short periods on high-risk pastures or mixing pastures with grasses to reduce legume intake. Early supplementation with dry forage can also help counteract the effects of lush spring pastures.

Rainfall and Dew

Heavy rainfall or dew on legume-rich pastures increases the likelihood of bloat in cattle. Wet conditions make legumes more palatable and accelerate the breakdown of plant material in the rumen, leading to faster fermentation and gas production [18]. Additionally, wet legumes can lead to a higher intake, as cattle tend to consume more during these conditions. To prevent bloat during periods of rain or heavy dew, it is advisable to limit grazing on legume-dominant pastures or to provide dry feed before cattle are allowed to graze. Grazing management during wet conditions is critical to reducing bloat risk.

Cold Weather

Cold weather can indirectly contribute to bloating by slowing down rumen motility and altering feeding patterns. During colder months, cattle may consume larger quantities of legumes to meet their energy needs, which can overwhelm the rumen's ability to process the forage efficiently [18]. Furthermore, colder temperatures can reduce the cow's ability to ruminate effectively, leading to increased risk of bloat. To mitigate these risks, it is important to adjust grazing practices during cold weather, such as providing supplemental feed, to balance the diet and ensuring cattle are not overly dependent on legume-rich forages for their energy needs.

Frost Conditions

Frost on legume-rich pastures can increase the risk of bloat in cattle. When legumes freeze, their cell walls rupture, making the plant material more easily fermentable in the rumen [18]. This results in a rapid release of soluble proteins, leading to a spike in fermentation and gas production. Grazing frost-covered legumes can, therefore, lead to a sudden onset of frothy bloat. To prevent this, cattle should be

removed from pastures with frost-damaged legumes or given dry forage to dilute their intake of highly fermentable plant material during frost conditions.

Time of Day for Grazing

The time of day when cattle are allowed to graze on legume-rich pastures can influence the risk of bloat [2, 5, 29, 30, 47]. Grazing during early morning or late evening, when legumes tend to have higher moisture content due to dew, can increase the likelihood of bloat. During these times, the soluble protein content of legumes is also higher, contributing to faster fermentation in the rumen. To reduce the risk of bloat, it is recommended to graze cattle during midday hours when the moisture content in the forage has decreased, and the legumes are less likely to cause frothy bloat.

NUTRITIONAL IMBALANCE

High Nitrogen Content in Legumes

Legumes are known for their high nitrogen content, which can contribute to the risk of bloat in cattle. Nitrogen in legumes is primarily found in the form of proteins and non-protein nitrogen, both of which are rapidly broken down in the rumen [14, 26, 49–51]. This rapid breakdown leads to excessive gas production, particularly in the form of ammonia and carbon dioxide, which can contribute to the formation of frothy bloat. To manage this risk, it is important to balance cattle diets with feeds that are lower in nitrogen, such as fibrous forages, to slow down the fermentation process and reduce gas buildup.

Low Fiber Content

While legumes are high in protein, they often have a lower fiber content compared to grasses. Fiber is essential for proper rumen function, as it stimulates chewing and saliva production, both of which help buffer the rumen and prevent foam formation [14, 50]. When cattle consume a diet high in legumes and low in fiber, fermentation occurs rapidly, leading to gas accumulation and bloat. To prevent this, it is important to supplement legume-rich diets with fibrous forages, such as hay or silage, which slow down fermentation and promote healthy rumen function.

Imbalance in Calcium and Magnesium

Legumes tend to have a high calcium-to-magnesium ratio, which can lead to an imbalance in mineral absorption and rumen function. Magnesium plays a critical role in muscle function, including the muscles responsible for rumen motility and eructation (belching) [14, 50]. A deficiency in magnesium can impair the animal's ability to expel gas, leading to an increased risk of bloat. To address this issue, it is important to provide mineral supplements that include adequate levels of magnesium, especially when cattle are grazing on legume-rich pastures, to ensure proper rumen motility and prevent gas buildup.

High Soluble Protein Content

Legumes contain high levels of soluble protein, which is quickly degraded in the rumen. This rapid degradation leads to the production of large amounts of ammonia and other gases, which can overwhelm the rumen's ability to expel them, resulting in bloat [14, 50]. Unlike insoluble proteins that are broken down more slowly, soluble proteins in legumes are rapidly fermented, leading to a sudden increase in gas production. To mitigate this, it is important to introduce legumes gradually into the diet and ensure that cattle have access to fiber-rich feeds that slow down the fermentation of soluble proteins.

Excessive Fermentation of Carbohydrates

In addition to their high protein content, legumes also contain fermentable carbohydrates, which contribute to the rapid fermentation process in the rumen [14, 50]. These carbohydrates are broken down quickly by rumen microbes, leading to the production of volatile fatty acids and gases such as carbon dioxide and methane. When excessive fermentation occurs, the risk of bloat increases due to the buildup of gases that are not expelled efficiently. To prevent this, cattle diets should be balanced with forages

that contain more complex carbohydrates, such as grasses, which are digested more slowly and produce less gas during fermentation.

Low Dry Matter Content in Legumes

Legumes typically have a lower dry matter content compared to other forages, especially when they are young and lush. This high moisture content means that cattle can consume large amounts of legume biomass quickly, without feeling full. As a result, cattle may ingest more legumes than their rumen can handle, leading to rapid fermentation and excessive gas production [14, 50]. The lower dry matter content also speeds up the digestion process, increasing the risk of bloat. To mitigate this, dry matter intake can be balanced by providing dry hay or forage before grazing on legume-rich pastures.

Imbalanced Forage to Concentrate Ratio

When cattle graze on legume-dominated pastures, the forage-to-concentrate ratio in their diet becomes skewed towards a high-protein, low-fiber intake. This imbalance leads to faster fermentation in the rumen, increasing the risk of bloat. Forage-to-concentrate ratios are critical in regulating rumen fermentation processes [14, 50]. By ensuring that cattle diets include an appropriate balance of fiber-rich forages and protein-rich legumes, rumen function remains more stable, reducing the likelihood of frothy bloat. Farmers can achieve this by supplementing legume-dominated pastures with dry forages or limiting the grazing of high-legume content areas.

High Levels of Non-Protein Nitrogen (NPN)

Legumes, especially alfalfa and clover, contain high levels of non-protein nitrogen (NPN), which can be rapidly converted to ammonia in the rumen. Excess ammonia increases the risk of bloat, as ruminal microbes quickly convert NPN into gases like methane and carbon dioxide [14]. The rapid buildup of ammonia can overwhelm the rumen's buffering systems, leading to frothy bloat. Managing the intake of NPN by reducing the amount of legume-rich forage or supplementing the diet with fiber-rich feedstuffs can help slow down the fermentation of NPN and prevent the onset of bloat.

Lack of Adequate Rumen Microflora Adaptation

The rumen microflora of cattle needs time to adjust to diets rich in legumes. When cattle are suddenly introduced to legume-dominant pastures without a period of adaptation, their rumen bacteria struggle to efficiently process the rapid influx of soluble proteins and carbohydrates, leading to excess gas production [14]. Gradual introduction to legume-rich diets allows the rumen microbial populations to adjust, facilitating more stable fermentation processes. This adaptation period can help cattle process legumes more efficiently, reducing the risk of frothy bloat. Therefore, incremental increases in legume intake are essential for maintaining healthy rumen function.

Rapid Passage of Feed

Legumes tend to pass through the digestive system more quickly than fibrous forages, leading to reduced chewing and saliva production. The quicker passage of legume material through the digestive tract limits the opportunity for thorough breakdown and fermentation, resulting in less efficient gas expulsion [14, 50]. The lack of saliva, which acts as a buffer for the rumen's acidity, further contributes to the conditions necessary for frothy bloat. By supplementing legume-heavy diets with fibrous feed, cattle spend more time chewing, which stimulates saliva production and slows down feed passage, reducing bloat risk.

EFFECTS ON RUMEN PHYSIOLOGY

Reduced Saliva Production

Saliva is a key component in regulating rumen pH and preventing bloat. When cattle consume legumes, especially in large quantities, they tend to produce less saliva compared to when they eat fibrous forages [39, 40]. Saliva helps neutralize acids produced during fermentation and prevents foam formation. Legumes, being lower in fiber, require less chewing, which reduces saliva production. The

lack of adequate saliva can lead to an acidic rumen environment that promotes frothy bloat. Increasing the fiber content in the diet, or feeding cattle dry forage before grazing on legumes, can help stimulate saliva production.

Foam Formation in the Rumen

The high soluble protein content in legumes contributes to foam formation in rumen. This foam traps gas, preventing it from being expelled through eructation, leading to the characteristic distension of frothy bloat [39, 40]. The foam forms a stable layer on top of the rumen contents, and the animal's natural expulsion mechanisms become ineffective. This trapped gas causes pressure to build, leading to discomfort and, in severe cases, death. Preventing foam formation can be achieved through dietary additives like poloxalene or oils, which act as defoaming agents, breaking down the foam and allowing gas to escape.

Inhibition of Rumen Motility

Rumen motility, the contraction and relaxation of the rumen walls, is essential for mixing ruminal contents and facilitating the expulsion of gases. Legumes can interfere with normal rumen motility due to the rapid production of gases and acids during fermentation [39, 40]. This disruption slows down the movement of rumen contents, exacerbating the buildup of gases. The resulting reduced motility increases the risk of bloat, as gas remains trapped in the rumen. Adding fibrous forage or feed additives to the diet can help maintain rumen motility and reduce the risk of bloat by promoting more effective digestion.

Alteration of Microbial Populations

The consumption of legumes can alter the balance of microbial populations in the rumen, favoring species that produce more gas such as methane and carbon dioxide. This shift in microbial activity results in excessive fermentation, increasing the risk of bloat [39, 40]. The rapid availability of soluble proteins and carbohydrates from legumes creates a fermentation environment that promotes gas-producing bacteria over other beneficial microbes. To counter this, probiotic supplements or gradual changes to the diet can help restore a balanced microbial population in the rumen, thereby preventing the overproduction of gases and reducing bloat risk.

Accumulation of Volatile Fatty Acids (VFAs)

The rapid fermentation of legumes in the rumen leads to the excessive production of volatile fatty acids (VFAs) such as acetic, propionic, and butyric acids. These acids lower the rumen pH, making the environment more acidic, which can promote foam formation and inhibit the animal's ability to expel gas [39, 40]. The accumulation of VFAs also slows down the absorption process, leading to further gas buildup. Maintaining a balanced diet with adequate fiber helps slow down fermentation, reducing the accumulation of VFAs and stabilizing rumen pH, thereby lowering the risk of frothy bloat.

PREVENTATIVE MEASURES

Poloxalene Supplementation

Poloxalene is an anti-foaming agent that can be added to cattle feed or administered as a drench to prevent bloat. It works by breaking down the foam that forms in the rumen, allowing gases to escape more easily through eructation [1, 5, 48]. Poloxalene is especially effective when cattle are grazing on pastures with a high legume content, where the risk of frothy bloat is elevated. It can be provided as a block, lick, or mixed into feed, and its regular use during high-risk periods significantly reduces the occurrence of bloat in cattle grazing on legume-dominated pastures.

Use of Ionophore Feed Additives

Ionophore feed additives, such as monensin, are commonly used to prevent bloat by modifying the rumen microbial population. These additives selectively inhibit gas-producing bacteria, thereby reducing the overall production of gases during fermentation [1, 5, 48]. By altering the microbial

environment, ionophores promote more efficient digestion and lower the risk of excessive gas accumulation that leads to bloat. When used in combination with other management practices, ionophore feed additives can be an effective strategy for preventing bloat in cattle grazing on legume-rich pastures, especially during high-risk times like early spring.

Ensuring Regular Feeding Schedules

Consistent feeding schedules help maintain stable rumen function and reduce the risk of bloat. Irregular feeding, or sudden changes in diet, can disrupt the rumen's microbial balance and fermentation processes, increasing the likelihood of bloat. When cattle are fed regularly, the rumen microbial populations are better able to adjust to the diet, including legumes, and fermentation processes remain more controlled. Ensuring that cattle are fed at the same time each day and providing a consistent mix of fibrous forages and legumes helps prevent the sudden onset of frothy bloat.

Provision of Mineral Supplements

Proper mineral supplementation, especially with magnesium and calcium, can help maintain rumen function and reduce the risk of bloat. Magnesium plays a critical role in maintaining muscle function, including the muscles responsible for rumen motility and gas expulsion. A deficiency in magnesium can impair rumen contractions and reduce the animal's ability to expel gases. Providing mineral supplements, particularly during high-risk grazing periods, ensures that cattle maintain proper muscle and digestive function, reducing the likelihood of bloat. Mineral blocks or supplements can be made available to cattle grazing on legume-dominant pastures as a preventive measure.

Avoiding High-Risk Pastures

In situations where bloat risk is high, such as after rain or during periods of rapid legume growth, it may be best to avoid grazing cattle on legume-dominated pastures altogether. By rotating cattle to different pastures or limiting access to legume-rich areas, farmers can effectively reduce the chances of bloat. High-risk conditions include wet pastures, early spring growth, or frost-damaged legumes, all of which increase the likelihood of excessive gas production. Monitoring pasture conditions and adjusting grazing strategies accordingly is a key preventative measure in managing bloat risk.

Controlled Grazing Management

Implementing controlled grazing strategies can significantly reduce the risk of bloating in cattle. By managing the timing and duration of grazing on legume-rich pastures, farmers can limit the amount of legumes cattle consume at any given time. Strip grazing, rotational grazing, or limiting access to high-legume pastures can help ensure that cattle do not overconsume legumes in a short period, which reduces the likelihood of rapid fermentation and gas buildup in the rumen. Controlled grazing also allows for better pasture utilization and the inclusion of grass species to balance the legume content.

Slow Introduction to Legume-Rich Pastures

Gradual introduction to legume-rich pastures helps the rumen microbial populations adjust to the increased levels of soluble protein and fermentable carbohydrates. By slowly increasing the time cattle spend grazing on legumes, their rumen adapts to the diet, reducing the risk of sudden gas accumulation that can lead to bloat. This practice is particularly important during early spring when legume growth is at its peak, or when cattle are moved to new pastures with high legume content. A slow transition gives the rumen microflora the opportunity to adjust and maintain stable fermentation processes.

Pre-Grazing Feeding of Dry Forage

Feeding cattle dry forage, such as hay or straw, before turning them out onto legume-dominant pastures can help reduce bloat risk. Dry forage provides extra fiber that slows down rumen fermentation and promotes saliva production, which buffers the rumen against foam formation. By filling the rumen with dry feed before grazing, cattle are less likely to overconsume legumes, and the rate of fermentation is moderate. This practice is especially useful in high-risk situations, such as after rainfall or during periods of rapid legume growth, when bloat risk is elevated.

Mixed Pasture Systems

Planting mixed pastures with a combination of legumes and grasses can reduce the risk of bloat by providing cattle with a more balanced diet. Grass, being higher in fiber and lower in soluble proteins than legumes, slow down rumen fermentation and reduce gas production. By including grasses, such as ryegrass or timothy in the pasture, cattle consume a more balanced mix of fiber and protein, which helps regulate rumen function and minimize bloat risk. Mixed pastures also improve pasture resilience and productivity, benefiting both cattle health and long-term pasture management.

Limiting Grazing During High-Risk Times

Bloat risk tends to be higher during certain times of the day, particularly early morning and late evening, when dew is present on legumes, increasing their palatability and fermentability. To reduce bloat risk, cattle should be restricted from grazing during these high-risk periods. Instead, allowing them to graze in the middle of the day, when the legumes are drier and less likely to cause bloat, can help prevent frothy bloat. Monitoring environmental conditions and adjusting grazing schedules accordingly is an important aspect of effective bloat management.

Avoiding Overgrazing

Overgrazing can exacerbate bloat risk by forcing cattle to consume more of the remaining legumes in a pasture as grass becomes depleted. When pastures are overgrazed, the proportion of legumes to grasses increases, leading to an imbalance in cattle diets that can promote rapid fermentation and gas buildup. Managing pasture stocking rates and avoiding overgrazing ensures that cattle have access to a more balanced forage mix, reducing their reliance on legumes. This practice not only reduces bloat risk but also promotes healthier pastures by preventing the overconsumption of certain plant species.

Reducing the Proportion of High-Risk Legume Species

Certain legume species, like alfalfa and clover, are more likely to cause bloat due to their high levels of soluble protein and rapid fermentability. By reducing the proportion of these high-risk legumes in pastures, the likelihood of bloat can be minimized. Instead, incorporating lower-risk legume species – such as birdsfoot trefoil, sainfoin, or cicer milkvetch – can help maintain the benefits of legumes without significantly increasing bloat risk. These legumes produce tannins, which bind to soluble proteins and reduce their ability to form foam, thereby lowering the incidence of bloat in cattle grazing on mixed pastures.

Feeding Probiotics or Yeast Supplements

Probiotic and yeast supplements can be used to help stabilize rumen fermentation and prevent excessive gas production. These additives promote a healthy balance of rumen microbes, particularly those that aid in breaking down fiber and reducing foam formation. Yeast supplements, for example, help maintain a stable rumen pH, which is essential for normal fermentation and gas expulsion. By improving microbial activity and digestion, probiotics and yeast supplements can play an important role in reducing the risk of bloat, especially when cattle are consuming a diet high in legumes.

Regular Monitoring of Cattle Health

Frequent monitoring of cattle for signs of bloat is crucial, especially during periods when they are grazing on legume-rich pastures. Early signs of bloat, such as distension of the left side, discomfort, or reduced grazing activity, should be addressed immediately to prevent more severe outcomes. Regular health checks, combined with the observation of grazing behavior, can help farmers catch bloat in their early stages and take necessary action. Quick interventions, such as administering anti-bloat agents or moving cattle to safer pastures, can prevent fatalities and reduce economic losses associated with bloat.

Adjusting Diet Based on Seasonal Risks

The risk of bloat varies throughout the year, with higher risks occurring during certain seasons – such as spring and fall – when legumes are lush and more abundant. Adjusting cattle diets based on these seasonal risks can help prevent bloat. For instance, providing more fibrous feed or hay during high-risk

periods can slow down rumen fermentation and promote the expulsion of gases. Additionally, reducing legume intake during these times by limiting access to certain pastures or rotating grazing areas ensures that cattle receive a more balanced diet, lowering the overall risk of bloat.

LIMITATIONS

Geographical focus on tropical regions may limit the applicability of findings to temperate zones, where different environmental conditions and legume species exist. While the review covers various legume species, it may not account for the specific characteristics and bloat risks associated with all regional legumes, potentially overlooking locally relevant factors. Additionally, the variability in individual cattle responses to legume consumption due to genetic, physiological, and health differences complicates the generalization of findings across diverse herds. The reliance on short-term studies restricts insights into the long-term effects of legume-induced bloat on cattle health, productivity, and pasture sustainability. Furthermore, the interactions between legumes and other dietary components may not be fully explored, limiting the understanding of how to effectively balance rations to mitigate bloat risk. Practical recommendations may also encounter implementation challenges due to economic constraints, farmer education, and regional differences in pasture management practices. Research gaps exist regarding certain management strategies or alternative legume species that could effectively mitigate bloat risk, necessitating further exploration. Finally, seasonal variations and unpredictable climate patterns can influence the growth and nutritional quality of legumes, potentially impacting their bloat-inducing properties, which may not be thoroughly addressed in the study. These limitations highlight the need for ongoing research and adaptive management strategies in the context of legume-induced bloat in tropical cattle.

FUTURE DIRECTIONS

Future research on legume-induced bloat in tropical cattle should focus on exploring alternative legume species with lower bloat potential, assessing their nutritional benefits and compatibility with existing pasture systems to enhance diversity and resilience. Longitudinal studies are essential to understand the chronic effects of legume consumption on cattle health and performance, while developing sophisticated nutritional models can optimize diets to mitigate bloat risk without compromising productivity. Investigating the rumen microbiome's role in bloat could lead to targeted interventions, such as probiotics, and researching innovative grazing management techniques will improve pasture utilization and animal welfare. Additionally, understanding environmental factors influencing bloat risk, implementing integrated pest management practices, and developing educational programs for farmers about bloat risks and best management practices are crucial. Economic impact analyses will aid in understanding the costs associated with bloat and the benefits of preventative measures. Finally, conducting cross-regional comparisons will provide insights into variability in legume-induced bloat across tropical regions, facilitating the development of tailored management strategies suited to local conditions. These future directions aim to enhance the understanding of legume-induced bloat and contribute to more sustainable cattle production in tropical environments.

CONCLUSION

Legume-induced bloat remains a major constraint to the productivity, health, and welfare of tropical cattle, particularly in grazing systems where highly digestible and protein-rich leguminous forages are extensively utilized. The development of bloat is a multifactorial process involving the interaction of forage characteristics, ruminal fermentation, animal physiology, and environmental and pasture-management conditions. High levels of soluble proteins and rapid fermentation of susceptible legumes promote the formation and stabilization of ruminal foam, thereby interfering with normal eructation and leading to excessive gas accumulation. The risk and severity of bloat may be further influenced by rainfall, pasture maturity, grazing intensity, and other management practices.

The consequences of severe bloat extend beyond immediate animal health concerns and can substantially affect livestock production and farm profitability through reduced feed intake, impaired nutrient utilization, decreased milk and meat production, increased veterinary expenditure, and, in

extreme cases, mortality. Therefore, effective management of legume-induced bloat requires an integrated approach rather than reliance on a single preventive measure. Strategic feeding practices, including the provision of dry forage before access to high-risk legumes and the maintenance of appropriate grass–legume proportions, can help moderate the risk. Feed additives – such as poloxalene and ionophores – may provide additional protection when used appropriately. Similarly, rotational grazing, regular monitoring of pasture conditions, and careful control of grazing access are important components of a comprehensive bloat-management program.

Overall, improving the management of legume-rich pastures offers an important opportunity to exploit the nutritional benefits of legumes while minimizing their associated risks. A better understanding of the interactions among forage composition, ruminal fermentation, climatic conditions, animal factors, and grazing management will be essential for developing more reliable and sustainable prevention strategies for tropical cattle production. Future research should focus on identifying practical, cost-effective, and region-specific interventions, including improved forage selection, precision pasture management, and innovative nutritional approaches. Such integrated strategies can contribute to reducing bloat-related losses, improving animal welfare and productivity, and enhancing the long-term sustainability and resilience of tropical livestock production systems.

REFERENCES

1. Clarke RT, Reid CS. Foamy bloat of cattle. A review. *Journal of dairy science*. 1974 Jul 1;57(7):753-85.
2. McMahon LR, McAllister TA, Berg BP, Majak W, Acharya SN, Popp JD, Coulman BE, Wang Y, Cheng KJ. A review of the effects of forage condensed tannins on ruminal fermentation and bloat in grazing cattle. *Canadian Journal of Plant Science*. 2000 Jul 1;80(3):469-85.
3. Thompson DJ, Brooke BM, Garland GJ, Hall JW, Majak W. Effect of stage of growth of alfalfa on the incidence of bloat in cattle. *Can J Anim Sci*. 2000;80(4):725–727. doi: 10.4141/A00-065.
4. Majak W, Hall JW, McAllister TA. Practical measures for reducing risk of alfalfa bloat in cattle. *J Range Manag*. 2001;54(4). doi: 10.2307/4003120.
5. Berg BP, Majak W, McAllister TA, Hall JW, McCartney D, Coulman BE, Goplen BP, Acharya SN, Tait RM, Cheng KJ. Bloat in cattle grazing alfalfa cultivars selected for a low initial rate of digestion: A review. *Canadian Journal of Plant Science*. 2000 Jul 1;80(3):493-502.
6. Berg BP, Majak W, McAllister TA, Hall JW, McCartney D, Coulman BE, Goplen BP, Acharya SN, Tait RM, Cheng KJ. Bloat in cattle grazing alfalfa cultivars selected for a low initial rate of digestion: A review. *Canadian Journal of Plant Science*. 2000 Jul 1;80(3):493-502.
7. Bretschneider G, Santini FJ, Fay JP, Faverin C. Effects of maize silage supplementation before lucerne grazing on the occurrence of bloat in cattle. *N Z J Agric Res*. 2001;44(4):241–251. doi: 10.1080/00288233.2001.9513481.
8. Saha D, Hossain MA, Juyena NS. Clinical management and control of clinical and induced bloat in cattle. 2006;23(2). doi: 10.5555/20083000197.
9. Yu MA, Damodaran S. Kinetics of protein foam destabilization: evaluation of a method using bovine serum albumin. *Journal of Agricultural and Food Chemistry*. 1991 Sep;39(9):1555-62.
10. Jones WT, Lyttleton JW, Clarke RTJ. Bloat in cattle: XXXIII. The soluble proteins of legume forages in New Zealand, and their relationship to bloat. *N Z J Agric Res*. 1970;13(1):149–156. doi: 10.1080/00288233.1970.10421204.
11. Jones WT, Anderson LB, Ross MD. Bloat in cattle: XXXIX. Detection of protein precipitants (flavolans) in legumes. *N Z J Agric Res*. 1973;16(3):441–446. doi: 10.1080/00288233.1973.10421128.
12. Clarke RTJ, Jones WT, Reid CSW. Bloat in cattle: XLI. Proteins of saliva as possible genetic markers in breeding programmes to reduce bloat susceptibility in cattle. *N Z J Agric Res*. 1974;17(4):411–415. doi: 10.1080/00288233.1974.10421026.
13. Mangan JL. Bloat in cattle: XI. The foaming properties of proteins, saponins, and rumen liquor. *N Z J Agric Res*. 1959. doi: 10.1080/00288233.1959.10427123.

14. Majak W, Hall JW, McCaughey WP. Pasture management strategies for reducing the risk of legume bloat in cattle. *J Anim Sci.* 1995;73(5):1493–1498. doi: 10.2527/1995.7351493x.
15. Jackson HD, DeVault RL, McNairy SA, Hatcher BW, Goetsch GD. Bloat in cattle. Its relationship to a respiratory inhibitor in Ladino clover and to phosphorus fertility of the soil. *J Anim Sci.* 1962;21(2):235–240. doi: 10.2527/jas1962.212235x.
16. Meyer RM, Bartley EE, Deyoe CW. Bloat in cattle. VII. Relation to amino acid composition of alfalfa as affected by maturity. *J Dairy Sci.* 1965;48(2):213–216. doi: 10.3168/jds.S0022-0302(65)88198-X.
17. Hall JW, Majak W, Van Ryswyk AL, Kalnin CM, Howarth RE. The relationship of rumen cations and soluble protein with predisposition of cattle to alfalfa bloat. *Can J Anim Sci.* 1988;68(2):431–437. doi: 10.4141/cjas88-048.
18. Hall JW, Majak W, Williams RJ, Howarth RE. Effect of daily weather conditions on bloat in cattle fed fresh alfalfa. *Can J Anim Sci.* 1984;64(4):943–950. doi: 10.4141/cjas84-106.
19. Wolfe EC, Lazenby A. Bloat incidence and liveweight gain in beef cattle on pastures containing different proportions of white clover (*Trifolium repens*). *Aust J Exp Agric.* 1972;12(55):119–125. doi: 10.1071/EA9720119.
20. Hall JW, Majak W. Effect of time of grazing or cutting and feeding on the incidence of alfalfa bloat in cattle. *Can J Anim Sci.* 1995;75(2):271–273. doi: 10.4141/cjas95-041.
21. Lowe LB. Prevention of bloat in pastured cattle. *Bovine Pract.* 1998;27–30. doi: 10.21423/bovine-vol1998no32.1p27–30.
22. Bretschneider G. An update on frothy bloat in cattle. 2010;42(3). doi: 10.5555/20113044590.
23. Teles Coutinho L, Augusto Bastos Afonso da Silva J, de Azevedo Costa N, Lopes de Mendonça C, Antônio da Rocha Faria P, Castro Soares P. Therapeutics evaluation in cases of frothy bloat in cattle. *Ciênc. anim. bras.(Impr.).* 2009;288-93.
24. Majak W, Garland GJ, Lysyk TJ. The effect of feeding hay before fresh alfalfa on the occurrence of frothy bloat in cattle. *Can J Anim Sci.* 2008;88(1):29–31. doi: 10.4141/CJAS07073.
25. Bartley EE, Nagaraja TG. Effect of Lasalocid or Monensin on feedlot (grain) bloat in cattle. *Kansas Agricultural Experiment Station Research Reports.* 1983 Jan 1;1(1).
26. Bartley EE, Nagaraja TG, Pressman ES, Dayton AD, Katz MP, Fina LR. Effects of lasalocid or monensin on legume or grain (feedlot) bloat. *J Anim Sci.* 1983;56(6):1400–1406. doi: 10.2527/jas1983.5661400x.
27. Hoshino S, Wakita M, Kobayashi Y. Effects of salinomycin on grain bloat in fattening Japanese Black cattle. *J Anim Physiol Anim Nutr (Berl).* 1987;57(1–5):123–129. doi: 10.1111/j.1439-0396.1987.tb00018.x.
28. Ayre-Smith RA. Pasture bloat in cattle. *Aust Vet J.* 1971;47(4). doi: 10.1111/j.1751-0813.1971.tb02130.x.
29. Bretschneider G, Peralta M, Santini FJ, Fay JP, Faverin C. Influence of corn silage supplementation before alfalfa grazing on ruminal environment in relation to the occurrence of frothy bloat in cattle. *Anim Feed Sci Technol.* 2007;136(1–2):23–37. doi: 10.1016/j.anifeeds.2006.08.019.
30. Wang Y, Berg BP, Barbieri LR, Veira DM, McAllister TA. Comparison of alfalfa and mixed alfalfa-sainfoin pastures for grazing cattle: Effects on incidence of bloat, ruminal fermentation, and feed intake. *Can J Anim Sci.* 2006;86(3):383–392. doi: 10.4141/A06-009.
31. Katz MP, Nagaraja TG, Fina LR. Ruminal changes in monensin- and lasalocid-fed cattle grazing bloat-provocative alfalfa pasture. *J Anim Sci.* 1986;63(4):1246–1257. doi: 10.2527/jas1986.6341246x.
32. Gadberry S, Lalman D, White F, Linneen S, Beck P. Meta-analysis of the effects of monensin on growth and bloat of cattle on pasture. *Transl Anim Sci.* 2022;6(2). doi: 10.1093/tas/txac031.
33. Dalto AG, Bandarra PM, Pedroso PM, Guagnini FD, Leal JD, Raymundo DL, Driemeier D. Timpanismo espumoso em bovinos leiteiros em pastagens de *Trifolium* spp. (Leg. Caesalpinoideae). *Pesquisa Veterinária Brasileira.* 2009 May;29(5):401-3.
34. Waghorn GC. Bloat in cattle. 47. Relationships between intra-ruminal pressure, distension, and the volume of gas used to simulate bloat in cows. *N Z J Agric Res.* 1991;34(2):213–220. doi: 10.1080/00288233.1991.10423362.

35. Paisley SI, Horn GW. Effect of ionophore on rumen characteristics, gas production, and occurrence of bloat in cattle grazing winter wheat pasture. 1998. Available from: <http://dasnr54.dasnr.okstate.edu:8080/beefextension2018/reports/1998/1998-3%20Paisley%20Research%20Report.pdf>
36. Kleiber M, Cole HH, Mead SW. Bloat in cattle and composition of rumen gases. *J Dairy Sci.* 1943;26(10):929–933. doi: 10.3168/jds.S0022-0302(43)92788-2.
37. Majak W, Howarth RE, Cheng KJ, Hall JW. Rumen conditions that predispose cattle to pasture bloat. *J Dairy Sci.* 1983;66(8):1683–1688. doi: 10.3168/jds.S0022-0302(83)81992-4.
38. Davis JD, Essig HW. Comparison of three bloat-preventing compounds for cattle grazing clover. *Can J Anim Sci.* 1972;52(2):329–335. doi: 10.4141/cjas72-037.
39. Meyer RM, Bartley EE, Morrill JL, Stewart WE. Salivation in cattle. I. Feed and animal factors affecting salivation and its relation to bloat. *J Dairy Sci.* 1964;47(12):1339–1345. doi: 10.3168/jds.S0022-0302(64)88915-3.
40. Bartley EE, Yadava IS. Bloat in cattle. IV. The role of bovine saliva, plant mucilages, and animal mucins. *J Anim Sci.* 1961;20(3):648–653. doi: 10.2527/jas1961.203648x.
41. Jones WT, Lyttleton JW. Bloat in cattle: XXXVII. The foaming properties of bovine salivary secretions and protozoal proteins. *N Z J Agric Res.* 1972;15(3):506–511. doi: 10.1080/00288233.1972.10430542.
42. Johns AT, McDowall FH, McGillivray WA. Bloat in cattle: XII. The effect of orally administered penicillin on body weight of lactating cows, on milk production, and on production and composition of fat. *New Zealand Journal of Agricultural Research.* 1959 Feb;2(1):62–71.
43. Wright DE. Bloat in cattle. *N Z J Agric Res.* 1961;4(3–4):203–215. doi: 10.1080/00288233.1961.10420366.
44. Mangan JL, Johns AT, Bailey RW. Bloat in cattle. *N Z J Agric Res.* 1959;2(2):342–354. doi: 10.1080/00288233.1959.10420322.
45. Sakauchi R, Hoshino S. Microbial characteristics of ruminal fluid from feedlot bloat beef cattle. *J Gen Appl Microbiol.* 1981;27(2):145–155. doi: 10.2323/jgam.27.145.
46. Clarke RTJ, Hungate RE. Bloat in cattle: XXXV. Microbial activities in the reticulo-rumens of cows differently susceptible to legume bloat. *N Z J Agric Res.* 1971;14(1):108–121. doi: 10.1080/00288233.1971.10421307.
47. Azad E, Derakhshani H, Forster RJ, Gruninger RJ, Acharya S, McAllister TA, Khafipour E. Characterization of the rumen and fecal microbiome in bloated and non-bloated cattle grazing alfalfa pastures and subjected to bloat prevention strategies. *Scientific reports.* 2019 Mar 12;9(1):4272.
48. Cheng KJ, McAllister TA, Popp JD, Hristov AN, Mir Z, Shin HT. A review of bloat in feedlot cattle. *J Anim Sci.* 1998;76(1):299–308. doi: 10.2527/1998.761299x.
49. Katz MP, Bartley EE, Pressman ES, Nagaraja TG. Effect of Rumensin on legume bloat in cattle. *Kansas Agric Exp Stn Res Rep.* 1982;(1):52–53. doi: 10.4148/2378-5977.2560.
50. Khatiwada B. Rejuvenation of depleted pasture using bloat-free legumes for high performance cattle grazing. 2018. Available from: <https://search.proquest.com/openview/da5852bf597ef9304ca87c49a5352149/1>
51. Bartley EE, Bassette R. Bloat in cattle. III. Composition of foam in legume bloat. *J Dairy Sci.* 1961;44(7):1365–1366. doi: 10.3168/jds.S0022-0302(61)89890-1.