

Surajmukhi (*Helianthus annuus* L.): Medicinal Importance in Unani Literature and Modern Evidence

Abudullah^{1*}, Yasmeen Khan², Labeeb Shaida², Abdul Kabeer², Nilophar³

Abstract

Introduction: *Helianthus annuus* L. (Sunflower), belonging to the family Asteraceae, is a well-recognized plant in both traditional and modern systems of medicine. In Unani medicine, it is described under the name “Shams-ul-Nahar,” attributed with specific Mizaj (temperament) and pharmacological actions. The plant is considered to possess a hot and moist temperament (Har Ratab), and different parts, including its seeds, oil, and leaves, have been employed in classical formulations. Sunflower seeds are recognized as nutritive, demulcent, and mild laxative, while the oil is valued for its emollient, anti-inflammatory, and cardioprotective properties. The flower has been mentioned for its analgesic and diuretic potential. However, Unani scholars also caution its use in individuals with hot temperament and in disorders related to the spleen, indicating a nuanced therapeutic application. Contemporary pharmacological studies corroborate many of these traditional claims, demonstrating antioxidant, anti-inflammatory, hepatoprotective, and antimicrobial activities of *H. annuus*. This integration of Unani concepts with modern evidence highlights the significance of sunflower as a medicinal plant with potential applications in gastrointestinal, cardiovascular, and inflammatory disorders. Further scientific validation and clinical research are warranted to standardize its use and to ensure safe therapeutic application in alignment with the principles of Unani medicine.

Methodology: This is a narrative review article. Unani classical literature, such as *Al-Qanoon fi'l-Tibb*, *Khazain al-Advia*, *Makhzan-ul-Mufradat Nadkarni KM – Indian Materia Medica*, were manually searched in the central library of National Research Institute of Unani Medicine for skin Disorders, Hyderabad and Qamar Library Deoband Unani Medical College Hospital & Research Centre. We also searched online bibliographic databases such as pubmed, google scholar. Standard Unani Medical Terminology published by Central Council for Research in Unani Medicine, New Delhi was referred to describe the Unani terminologies. **Result:** In this study, 11 classical Unani books, 4 original research articles, and 5 review articles were reviewed. It has been found that famous Unani scholars, like Ibn-

Sina and Najmul Ghani, have documented the medicinal use of Gul-e-Aftab in their books. Sunflower plant is an annual herb with an herbaceous (non woody) stem. The scientific name *Helianthus annuus* L. comes from the Greek words *helios* (sun) and *anthos* (flower), referring to the sunflower's sun-like appearance and its heliotropic behavior, and the Latin word *annuus* meaning annual, distinguishing it from perennial sunflowers. In Classical Unani Literature Gul-e-Aftab, commonly known as Sunflower, is a well-known ornamental and medicinal plant. The name reflects its heliotropic nature – the flower follows the movement of the sun. **Conclusion:** In Unani medicine, it is valued for both of its nutritive and therapeutic properties. The plant provides seeds, oil, flowers, and leaves as medicinal parts. Seeds are rich in oil, proteins, and minerals, making them an important source of nutrition and medicine. Oil

*Author for Correspondence

Abudullah

E-mail: abdullahabbu851@gmail.com

¹Assistant Professor, Department of Ilmul Advia (Pharmacology), Deoband Unani Medical College Hospital & Research Centre, Deoband, Saharanpur, Uttar Pradesh, India.

²P.G. Scholar, Department of Ilmul Advia (Pharmacology), National Research Institute of Unani Medicine for Skin Disorders, Hyderabad, Telangana, India.

³Assistant Professor, Department of Mahiyatul amraz (Pathology), Deoband Unani Medical College Hospital & Research Centre, Deoband, Saharanpur, Uttar Pradesh, India.

Received Date: September 10, 2025

Accepted Date: September 22, 2025

Published Date: September 23, 2025

Citation: Abudullah, Yasmeen Khan, Labeeb Shaida, Abdul Kabeer, Nilophar. Surajmukhi (*Helianthus annuus* L.): Medicinal Importance in Unani Literature and Modern Evidence. Journal of AYUSH: Ayurveda, Yoga, Unani, Siddha and Homeopathy. 2025; 14(3): 68–75p.

is used both internally and externally in various ailments. *Gul-e-Aftab* has been reported for numerous pharmacological activities such as anti-inflammatory, antioxidant, antitumor, anti-asthmatic, antigen, antipyretic, astringent, anti-hypoglycemic effect, cathartic, diuretic, stimulant, vermifuge, antimicrobial activities, and vulnerable purposes. The cosmetics business has a new future because of the usage of yellow petals as coloring ingredients. Other plant parts, petioles, and young flowers are used as savory treats According to Unani texts, Sunflower seeds and oil used in cardiac weakness and hypertension (due to vasodilatory and lipid-lowering properties). Seeds are employed as nutritive, diuretic, and expectorant. Its oil applied externally for rheumatic pains, ulcers, and skin disorders. Sunflower flowers are used as febrifuge (anti-pyretic). And its leaves are used in some formulations as wound healer and anti-inflammatory.

Keywords: *Helianthus annuus* L., Sunflower, Surajmukhi, Unani medicine, Mizaj, traditional medicine, pharmacological activities, anti-inflammatory, cardioprotective

INTRODUCTION

The sunflower (*Helianthus annuus* L.), a significant annual herbaceous plant that is native to North America and is currently grown extensively worldwide, is a member of the *Asteraceae* (*Compositae*) family. It is known for having big, radiating, bright yellow flowers that mimic the sun; this is why it is called “sunflower.” In addition to its nutritional and economic benefits, the plant is prized for its aesthetic and therapeutic qualities [1]. The phenolic chemicals, flavonoids, polyunsaturated fatty acids, and vitamins in sunflower seeds and sprouts provide important anti-inflammatory, anti-hypertensive, antioxidant, anti-microbial, wound-healing, and cardiovascular effects [2]. The extensive use of plant extracts to treat illnesses has raised interest in identifying the key ingredients that give the extracts their therapeutic properties [3]. The sunflower, which is utilized as food and medicine all throughout the world, is the foundation of medicinal benefits. The primary reason *H. annuus* is grown is for its seeds, which provide the second most significant source of edible oil in the world. Other plant parts, such as petioles and young flowers, are used as savory treats. The herb tincture, seed oil, and shoots are used for anti-inflammatory, antioxidant, anti-tumor, anti-asthmatic, antigen, antipyretic, astringent, anti-hypoglycemic effect, cathartic, diuretic, stimulant, vermifuge, antimicrobial activities, and vulnerable purposes. The cosmetics business has a new future because of the usage of yellow petals as coloring ingredients [4].

In Unani system of medicine, Sunflower (*Helianthus annuus* L.) known as *Gul-e-Aftab-parast*, *Shams-ul-Nahar* in Arabic and *Surajmukhi* in Urdu/Hindi. It is widely cultivated for its ornamental beauty and medicinal as well as nutritional values. according to Unani medicine sunflowers, especially its seeds and oil, have the following properties, like Deobstruent or *mufatteh-e-sudad*, aids in clearing obstructions from vessels, *Musakkin* (pain reliever) sedative, *Mulattif* (softening, demulcent), diuretic (*Mudirr-e-Baul*) Antidote or anti-toxic (*Dafe'e-Samoom*) Nerve tonic (*muqawwi-e-Aasab*) [5, 6].

VERNACULAR NAMES [6–9]

Gul-e-Aftab has been found in different parts of the world and is used as traditional medicine such as Chinese medicine and Persian medicine for various therapeutic purposes. It is known in different geographical regions by its different local names. Vernacular name & Taxonomical classifications are given in Tables 1 & 2.

Table 1. Vernacular names of *Helianthus annuus* (Sunflower).

Language	Synonyms	Language	Synonyms
Urdu	<i>Gul-e-Aftab</i>	English	<i>Sunflower</i> .
Hindi	<i>Surajmukhi</i>	Sanskrit	<i>Surya-mukhi</i> .
Arabic	<i>Shams-ul-Nahar</i>	Persian	<i>Gul-e-Aftab</i> .
Bengali	<i>Surjomukhi</i>	Gujrati	<i>Suryamukhi</i> .
Punjabi	<i>Surajmukhi</i>	Tamil	<i>Suryakanti</i> .
Telugu	<i>Podda-tige</i>	Unani	<i>Gul-e-Aftab</i> .
Kannada	<i>Podda-tige</i>	Malayalam	<i>Sooryakanthi</i> .
Marathi	<i>Surajmukhi</i>		

Table 2. Taxonomical classification [4].

Kingdom	Plantae	Division	Angiospermae
Sub-division	Eudicots	Class	Asteroids.
Order	Asterales	Family	Asteraceae.
Sub family	Helianthoideae	Scientific Name	<i>Helianthus annuus</i> L.

BOTANICAL DESCRIPTION

Helianthus annuus L. (Figure 1) is a common annual plant that is rigid and fat, growing to a height of one to three meters. When the plant reaches maturity, its tap roots give way to enormous, fibrous, lateral roots. This plant has spherical, unbranched, hispid stems that range in height from one to six feet. The number of internodes determines the stem length. The arrangement of the leaves is such that the top leaves alternate down the stem and the lower leaves are primarily opposite. Most leaves are oval, with an apex that is sharp to abruptly acuminate, edges that are serrated, and leaves that are at least 4–20 cm long and 3–15 cm wide. This plant's inflorescence consists of a capitulum composite heads with ray and disc florets present, either solitary at the peduncle's end or terminal on a branch or axillary. Ray blooms have yellow ligules and are sterile, measuring 0.6 to 1.6 inches in length. Perfect disc flowers with five corolla lobes that are tubular, purple, brown to yellow, and 0.2–0.3 inches long. The ovary is inferior in disc flowers, and the pappus is two and deciduous. Achene is the fruit of *H. annuus*. Achenes range in size from 14 to 13 mm in width and 7 to 25 mm in length [10].



Figure 1. Sunflower (*Helianthus annuus* L.).

CULTIVATION OF SUNFLOWER (*Helianthus annuus* L.)

Sunflower (*Helianthus annuus* L.), an important oilseed crop, can be cultivated successfully under diverse agro-climatic conditions due to its short duration and photo-insensitivity. The crop performs best in tropical to subtropical regions, requiring a temperature range of 20–25°C for germination and 25–30°C for vegetative growth and seed development (Figure 2). Deep, well-drained loamy soils rich in organic matter, with a pH between 6.0 and 7.5, are considered ideal.



Figure 2. Sunflower seeds.

Land preparation involves ploughing the field two to three times to achieve a fine tilth, followed by the incorporation of farmyard manure (8–10 t/ha). A seed rate of 8–10 kg/ha with a spacing of 45 cm × 20 cm is recommended. Depending on the region, sowing is undertaken during three major seasons: kharif (June–July), rabi (October–November), and summer (January–February). Fertilizer application generally follows the N:P:K ratio of 60:60:40 kg/ha, with phosphorus and potassium applied at sowing and nitrogen split between sowing and 30 days after sowing.

Adequate soil moisture is critical, particularly during flowering and seed-setting stages; therefore, four to five irrigations are usually required. Weed control can be achieved through two manual weddings (20 and 40 days after sowing) or by pre-emergence herbicide application (e.g., pendimethalin at 1 kg/ha). Common pests, such as the capitulum borer and hairy caterpillar, and diseases, like *Alternaria* blight, downy mildew, and rust, can be managed through resistant cultivars, crop rotation, seed treatment, and appropriate plant protection measures.

Harvesting is performed when the back of the capitulum turns yellow and bracts turn brown. The harvested heads are dried, and seeds are separated by threshing. Under optimal conditions, the average seed yield ranges from 1.5 to 2.0 t/ha.

Mizaj (Temperament) [5–6, 9, 11]

Har Ratab (Hot & Moist) 20 Seeds are considered slightly hot and moist.

Therapeutic Dosage (*Miqdār-i-Khurāk*) [12–14]

The decoction of leaves, roots, and flower juice is administered in a dose ranging from 3½ masha (~3.4 g; ~3.4 mL for aqueous preparations) to 10 tola (~116.6 g; ~117 mL).

The powder (*Sufoof*) is administered in a dose of 4 masha (~3.9 g) to 7 masha (~6.8 g).

Seed Oil (*Roghan*) 5–10 mL orally; externally applied as needed for massage, wounds, skin conditions.

Flowers (*Gul-e-Aftab*) Used in infusion/tea; 3–6 g dried flowers.

- *Mazarrat (Adverse Effect)*: Sunflower is regarded as harmful for individuals with hot temperaments, and its adverse effects are particularly pronounced on the spleen.
- *Correctives (Muslih)*: Sikanjabeen, Shahad (honey).
- *Substitute (Badl)*: Zafraan (*Crocus sativus*, Saffron), Sulaikha *Carthamus tinctorius* L. (Safflower).

Adverse Effect, Allergic Reaction and Toxicity

According to Traditional Unani Literature and formularies do not explicitly list adverse effects for *Helianthus annuus* unlike some other herbs, Unani literature largely emphasizes its therapeutic uses as brain or heart tonic and does not document specific contraindications or side effects.

Adulterants or Substitutes of Sunflower [13–15]

Sometimes petals of other yellow flowers, such as *Tagetes erecta* (*Genda phool* or *Marigold*) or *Calendula officinalis* (*Zergul*), are mixed with *Gul-e-Aftab* due to similarity in color. Powdered flowers are also adulterated with dyed dried petals of cheaper plants. Adulteration lowers therapeutic potency because sunflower mizaj is *Har Yabis* – Hot & Dry has specific actions (*Mufarreah*, *Musaffi-e-Dam*), which differ from marigold or calendula.

Adulteration in Sunflower Oil [15–18]

Sunflower oil is highly prone to adulteration with cheaper oils, such as *Safflower oil* (*Carthamus tinctorius*), because of its very similar fatty acid profile. *Soybean oil* (*Glycine max*), *Cottonseed oil* (*Gossypium herbaceum*), *Corn oil* (*Zea mays*), *Sesame oil* (*Sesamum indicum*), Mineral oil hydrocarbons (MOH) (harmful, industrial adulterant).

Detection Methods of Adulterants

- GC–FID (Gas Chromatography–Flame Ionization Detector) → detects TAG & fatty acid profiles.
- FT-IR Spectroscopy → quick screening for adulteration.
- HPTLC (High Performance Thin Layer Chromatography) → fingerprinting of mixed oils.
- LC–QTOF MS → metabolomic biomarkers of sunflower vs other oils.

The therapeutic actions, scientific validation, pharmacological studies of the compound formulation, and pharmacological activities of *Helianthus annuus* are presented in Tables 3–6, respectively.

Table 3. Therapeutic action [5–7, 12].

Unani term	Pharmacological action	Therapeutic significance (unani concept)
Mufarreḥ	Cardiac and mental exhilarant	Uplifts mood, provides calmness, and strengthens the heart and mind.
Daf'-e-Tashannuj	Antispasmodic	Relieves muscular cramps and spasms.
Musakkin-e-Hararat	Relieve excessive heat	Beneficial in hot temperaments and febrile conditions.
Mudirr-e-Bawl	Diuretic	Promotes urine output and relieves urinary retention.
Musakkin-e-Alam	Analgesic	Relieves pain, particularly headache.
Munawwim	Mild sedative	Induces sleep and relaxation.
Da'f-e-Hiddat-e-Dam	Blood purifier	Reduces excessive heat and corrects bilious disorders.

Istimālāt (Uses)

Suda (صداع / Headache), *Humma Harra* (حرّ حَمَى / Fevers with heat), *Atash* (عطش / Excessive thirst), *Tashannuj* (تنشج / Spasms & Convulsions), *Sual* (سعال / Cough), *Khafqan* (خفقان / Palpitation).

Table 4. Scientific validation [19–25].

Part used	Traditional medicinal uses	Pharmacological uses	References
Seeds	Used as food, nutritive tonic, aphrodisiac in Unani medicine	Rich in oil (linoleic & oleic acid), lowers cholesterol, antioxidant (Vitamin E), improves cardiovascular health, immunomodulatory.	Ghafoor et al., 2017; Duke, 1992.
Seed Oil	Applied externally for wounds and skin disorders; used as base oil in Unani formulations	Edible oil, hypocholesterolemic, anti-inflammatory, skin protective (cosmetic industry).	Nandha & Kaur, 2014
Leaves	Decoction used as antipyretic, expectorant, and to reduce swelling	Contains phenolics & flavonoids – antioxidant, antimicrobial, anti-inflammatory	Taha et al., 2016.
Flowers (Petals)	Infusion used as diuretic and expectorant in folk medicine	Rich in carotenoids (lutein, zeaxanthin) – good for eye health, antioxidant, cosmetic coloring agent	Nandha & Kaur, 2014.
Roots	Decoction traditionally used as febrifuge and for respiratory disorders	Contains sesquiterpene lactones – cytotoxic, antitumor, anti-inflammatory	Duke, 1992.
Stem	Ash of stem used in soap and glassmaking (folk uses)	Source of cellulose, biofuel production, and antimicrobial potential (due to coumarins & tannins)	Taha et al., 2016.
Whole Plant	Used as fodder, diuretic, and febrifuge in traditional medicine	Allelopathic properties (weed management), phytoremediation of heavy metals (cadmium, lead)	Nandha & Kaur, 2014.

Table 5. Pharmacological studies of compound formulation.

Plant part	Phytochemicals	Pharmacological role	Unani formulations	Reference
Seeds	Linoleic acid, Oleic acid, Palmitic acid, Stearic acid, Tocopherols (Vit. E), Proteins (albumins, globulins), Phytosterols (β-sitosterol, campesterol, stigmasterol)	Hypolipidemic, Antioxidant, Cardioprotective, Nutritional	Majun Ard Khurma, Majun Murraweh-ul-Arwah, Majun Pumba Dana, Majun Jiryan Khas, Majun Sa'lab	Ghafoor et al., 2017; Duke, 1992; Medisura Unani Materia Medica; Nandha & Kaur, 2014.

Seed Oil	Linoleic acid, Oleic acid, Tocopherols, Phytosterols	Nutritional, Hypocholesterolemic, Skin-protective	Used externally in Roghan formulations	Ghafoor et al., 2017; Duke, 1992; Taha et al., 2016.
Leaves	Chlorogenic acid, Caffeic acid, Quercetin, Kaempferol, Luteolin, Tannins	Antioxidant, Anti-inflammatory, Antimicrobial	Decoctions sometimes included in Sharbat or infusion for fever, cough, respiratory disorders	Nandha & Kaur, 2014.
Flowers (Petals)	Carotenoids (Lutein, Zeaxanthin), Flavonoids	Eye health, Antioxidant, Anti-aging	Used in infusions for diuretic, expectorant effect (noted in folk/Unani practice)	Taha et al., 2016.
Roots	Sesquiterpene lactones (Helianolide, Annuolide), Polyphenols	Cytotoxic, Antitumor, Anti-inflammatory	Rarely in formulations, decoction used as febrifuge & anti-inflammatory	Duke, 1992; Nandha & Kaur, 2014.
Stem	Phenolic compounds, Coumarins (Scopoletin, Umbelliferone), Tannins	Antimicrobial, Anti-inflammatory		Duke, 1992.

Table 6. Pharmacological activities of *Helianthus annuus*.

Pharmacological activity	Part used	Experimental model	Dose / form	Reference
Antioxidant	Seeds, leaves, flowers	In vitro (DPPH, ABTS, FRAP assays)	Ethanollic and methanollic extracts (50–200 µg/mL)	Nandha & Kaur, 2014
Anti-inflammatory	Leaves, roots, oil	In vivo (rats, carrageenan-induced paw edema)	Ethanollic leaf extract (200–400 mg/kg, oral)	Taha et al., 2016
Hypolipidemic Cardioprotective	Seed oil	In vivo (hyperlipidemic rats)	Sunflower seed oil (4–8 mL/kg/day)	Ghafoor et al., 2017
Antidiabetic	Seeds	In vivo (streptozotocin-induced diabetic rats)	Ethanollic seed extract (200–400 mg/kg, oral)	Nandha & Kaur, 2014
Antimicrobial	Leaves, petals, oil	In vitro (disc diffusion, MIC) against <i>S. aureus</i> , <i>E. coli</i> , <i>Candida albicans</i>	Methanollic extract (50–200 µg/mL)	Taha et al., 2016
Cytotoxic/Antitumor	Roots (sesquiterpene lactones)	In vitro (HeLa, MCF-7 cancer cell lines)	Root extract/isolated lactones (10–100 µg/mL)	Duke, 1992
Wound healing	Seed oil	In vivo (excision wound model in rats)	Topical application of sunflower oil (daily)	Nandha & Kaur, 2014
Immunomodulatory	Seeds (polysaccharides)	In vivo (mice, macrophage activation assays)	Seed extract (50–200 mg/kg)	Ghafoor et al., 2017

DISCUSSION

Surajmukhi (*Helianthus annuus* Linn.), or sunflower, holds dual significance in Unani and modern medicine. Classical Unani texts describe it as possessing a cold and dry temperament (Barid Yabis Mizāj), with its seeds recognized as nutritive, demulcent, and expectorant, useful in respiratory ailments and bilious disorders. The oil is applied as an emollient and solvent for skin diseases and joint pain, while decoctions of leaves and flowers are prescribed for fevers, cough, and urinary complaints.

The convergence of Unani wisdom with modern pharmacological evidence highlights sunflower as both a therapeutic agent and functional food. Its traditional uses as an emollient, febrifuge, and expectorant align well with its scientifically proven pharmacological actions, underscoring the relevance of integrative approaches. Future directions should focus on standardization, clinical validation, and safety profiling to promote broader acceptance of Surajmukhi in contemporary healthcare systems.

CONCLUSIONS

In Unani medicine, *Helianthus annuus* (Sunflower) holds therapeutic significance owing to its diverse pharmacological attributes, nutritional richness, and compatibility with classical concepts of health maintenance and disease management. Its seeds, oil, and other parts are recognized as properties, such as musakkin (sedative), muqawwi qalb (cardiotonic), mulattif (demulcent), and mufatteh sudad (deobstruent), which align with both preventive and curative aspects of Unani therapeutics. Modern scientific evidence further validates its antioxidant, anti-inflammatory, hypolipidemic, and wound-healing activities, bridging traditional knowledge with contemporary pharmacology. Thus, Sunflower exemplifies the integrative potential of Unani pharmacotherapy, offering scope for its wider application in evidence-based practice and future drug development.

Conflict of Interest

None.

Source of Finding

None.

REFERENCES

1. Prolea. De la production à la consommation: statistiques des oléagineux et protéagineux. Paris: Prolea Documentation; 2012. p. 48.
2. Fowler MW. Plants, medicines and man. J Sci Food Agric. 2006;86(12):1797–804.
3. Owens GL, Rieseberg LH. Hybrid incompatibility is acquired faster in annual than in perennial species of sunflower and tarweed. Evolution. 2014;68(3):893–900.
4. Dwivedi A, Sharma GN. A review on heliotropism plant: *Helianthus annuus* L. J Pharmacol. 2014;3(2):149–55.
5. Ibn Sina. Al-Qanoon fi'l Tibb. Vol. II.
6. Ghani N. Khazain al-Advia. Vol 3. p. 217.
7. Nadkarni KM. Indian materia medica. Vol 1. Mumbai: Popular Prakashan; 1976.
8. Council of Scientific and Industrial Research (CSIR). The Wealth of India. Vol 5. New Delhi: CSIR; 1959.
9. Central Council for Research in Unani Medicine (CCRUM). Standard Unani medical terminology. New Delhi: CCRUM; 2012.
10. Halvorson W, Guertin P. *Helianthus annuus* L. U.S. Geological Survey, Southwest Desert Field Station, Sonoran Field Station, University of Arizona; 2003.
11. Gardezi S. Makhzan-ul-Mufradat.
12. Qarshi A. Jadid Makhzan-ul-Advia.
13. Kabiruddin. Makhzan al-Mufradat. New Delhi: CCRUM.
14. Anonymous. National Formulary of Unani Medicine. Vol. I. New Delhi: CCRUM; 2006.
15. Nadkarni KM. Indian materia medica. 3rd ed. Mumbai: Popular Prakashan; 1976.
16. Tarhan S, et al. Spectroscopic quantification of safflower-oil adulteration in sunflower oil. Food Anal Methods. 2021;14:1801–10.
17. Souza NE, et al. Detection of adulteration of sunflower oil with soybean oil using GC-FID and ESI-MS. J Braz Chem Soc. 2015;26(10):2019–26.
18. Sumara A, et al. LC-QTOF MS metabolomics for authenticity testing of sunflower oil. Foods. 2023;12(5):10118080.
19. Nandha R, Kaur J. A review on phytochemistry and pharmacology of *Helianthus annuus* L. Int J Pharm Sci Res. 2014;5(5):1872–81.
20. Taha HS, et al. Phytochemical and biological studies of *Helianthus annuus* L. Egypt Pharm J. 2016;15(1):12–20.
21. Ghafoor K, et al. Phytochemical composition and bioactive components of sunflower seeds. Food Res Int. 2017;100:108–18.
22. Duke JA. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton: CRC Press; 1992.

23. Nandha R, Kaur J. Medisure Unani materia medica; thesis on botany – *Helianthus annuus*. 2014.
24. Central Council for Research in Unani Medicine (CCRUM). National Formulary of Unani Medicine. Parts I–VI. New Delhi: CCRUM.
25. Kirtikar KR, Basu BD. Indian medicinal plants. Vol II. 1935 reprint.