

The Therapeutic and Biotechnological Potential of *Pogostemon benghalensis* – A Comprehensive Review

Soumili Majumdar*

Abstract

Pogostemon benghalensis is a fragrant perennial shrub belonging to the family Lamiaceae, widely distributed across tropical and subtropical regions of Asia. It has been extensively utilized in traditional and ethnomedicinal systems for the treatment of wounds, skin infections, fever, digestive disorders, inflammation, and respiratory ailments. The plant is also valued for its antiseptic, aphrodisiac, insect-repellent, and veterinary medicinal properties. Its leaves, roots, and essential oil are incorporated into herbal formulations, natural dyes, cosmetics, and traditional healthcare practices. Recent phytochemical investigations have identified more than 140 bioactive constituents, including sesquiterpenes, monoterpenes, flavonoids, alkaloids, phenolic compounds, lignans, glycosides, tannins, and triterpenoids, which contribute to its diverse biological activities. Experimental studies have demonstrated that these phytochemicals possess significant antibacterial, antifungal, antiviral, antioxidant, anti-inflammatory, larvicidal, hepatoprotective, neuroprotective, and cytoprotective properties. The essential oil exhibits remarkable free radical scavenging activity, inhibition of lipid peroxidation, and potent antimicrobial efficacy, highlighting its therapeutic potential. In addition to its medicinal importance, *P. benghalensis* has emerged as a promising source of eco-friendly biopesticides for sustainable agriculture. Its bioactive metabolites exhibit insecticidal, larvicidal, and pest-repellent activities against several agricultural pests, offering an environmentally safe alternative to synthetic pesticides and contributing to crop protection and environmental restoration. Furthermore, the plant has considerable industrial significance due to its applications in pharmaceutical formulations, herbal cosmetics, perfumery, food preservation, natural fragrances, bioactive packaging materials, and green agricultural products. This review comprehensively summarizes the phytochemistry, pharmacological properties, therapeutic potential, bio-pesticidal applications, and industrial prospects of *Pogostemon benghalensis*, emphasizing its potential as a valuable natural resource for future drug discovery, sustainable crop management, and environmentally responsible industrial development.

Keywords: *Pogostemon benghalensis*, patchouli, ethnomedicine, sesquiterpenes, flavonoids, alkaloids, lignans, and glycosides antimicrobial, anti-fungal, anti-tumor, antioxidant, patchouli alcohol, hepatoprotective, neuroprotective, phytopharmacology, bio pesticide

*Author for Correspondence

Soumili Majumdar
E-mail: soumilimajumdar52@gmail.com

Student, KIIT School of Biotechnology, KIIT – Deemed to be University, Bhubaneswar, Odissa, India

Received Date: April 16, 2026
Accepted Date: July 10, 2026
Published Date: July 11, 2026

Citation: Soumili Majumdar. The Therapeutic and Biotechnological Potential of *Pogostemon benghalensis* – A Comprehensive Review. Research & Reviews: Journal of Botany. 2026; 15(2): 55–63p.

INTRODUCTION

Medicinal plants are an essential source of biologically active compounds and continue to contribute significantly to the pharmaceutical, cosmetic, food, and fragrance industries. Despite advances in synthetic drug development, plant-derived metabolites remain important for the discovery of new therapeutic agents because of their structural diversity and broad pharmacological potential. Consequently, medicinal plants continue to play a central role in both traditional medicine and modern healthcare systems [1].

Pogostemon benghalensis (Burm.f.) Kuntze is an aromatic shrub native to South and Southeast Asia, commonly distributed across India, Nepal, Bangladesh, Myanmar, Sri Lanka, China, Thailand, Vietnam, and Bhutan. The leaves contain an essential oil rich in sesquiterpenes, obtained mainly through steam distillation. Traditionally, the plant has been used as a medicinal herb, natural insect repellent, and fragrance source. Its essential oil is widely incorporated into perfumes, cosmetics, soaps, incense, and selected food products because of its distinctive aroma and fixative properties [2].

The therapeutic value of *P. benghalensis* is largely associated with its sesquiterpenes and phenolic constituents, particularly patchouli alcohol, the principal bioactive compound. Experimental studies have demonstrated that this compound possesses neuroprotective potential and may improve cognitive performance. In addition, the essential oil exhibits a wide range of biological activities, including antimicrobial, antioxidant, anti-inflammatory, antiviral, analgesic, immunomodulatory, and antiemetic effects. These pharmacological properties highlight the potential of *P. benghalensis* as a promising natural source for future therapeutic and functional health applications [3].

LEAF MORPHOLOGY

Pogostemon species are evergreen perennial shrubs with opposite, ovate leaves that contain abundant glandular and non-glandular trichomes, particularly on the lower leaf surface and along the veins. The *Glandular trichomes* are responsible for the production and storage of essential oil. Fresh leaves have a mild aroma that becomes stronger after drying or fermentation because of changes in volatile compounds. The plant is illustrated in Figure 1. Essential oil content varies with cultivar, plant age, and environmental conditions, typically ranging from 0.25–0.75% in fresh leaves. After drying, which removes most of the leaf moisture, oil yield generally increases to 1.8–2.5%, with optimized extraction methods producing yields of up to 5% [4].



Figure 1. Photograph of *Pogostemon patchouli*.

Floral Morphology of PP Under Stereoscope

Pogostemon produces terminal spike-like inflorescences composed of densely arranged flower clusters. The flowers have a tubular calyx and a bilabiate corolla with purple and white coloration. Four protruding stamens and a long, exerted style facilitate pollination. After flowering, small brown seeds develop inside tiny capsules along the spikes. The plant possesses a well-developed, branched root system that extends deep into the soil, while its leaves, flowers, and seeds all emit a characteristic aromatic fragrance [5].

LEAF MICROSCOPY

Microscopic observations indicate that essential oil-producing secretory structures are present in the leaf blade, margins, and petiole of *Pogostemon*. *Glandular trichomes* are far more abundant on the leaf lamina than on the petiole and occur in greater numbers on the lower leaf surface than on the upper surface. They are distributed throughout the leaf but are most concentrated along the veins, reflecting their important role in essential oil synthesis and storage (Table 1) [6].

Table 1. Secondary metabolites & their chemical constituents.

Secondary metabolites	Chemical constituents
Sesquiterpenes	Patchouli alcohol, cycloseychellene, pogostol, norpatchoulene, cadinene, α -atchoulene, β -patchoulene, seychellene α -bulnsene, α -guaiene, β -elemene, α -humulene, caryophyllene.
Sesquiterpene alkaloids	Patchouli-pyridine, epiguaipyridine, guaipyridene.
Oxygenated sesquiterpenes	1 α , 5 α -epoxy- α -guaiene, 1,10-epoxy-alphabulnesene, epoxycaryophyllene.
Phenolic compounds, Flavonoids, flavonoid glucosides	Eugenol, rhamnetin, apigenin, Lichochalcone A, ombuine, pachypodol, kumatakenin, 7, 4'-di-o-methylethiodictyol, 7, 3', 4'-tri-o-methylethiodictyol, 3, 7, 4'-tri-o-methylkaemferol.

HABITAT AND DISTRIBUTION

The genus *Pogostemon* belongs to the family Lamiaceae, one of the largest families of flowering plants. It comprises about 40 species that are mainly distributed across the tropical and subtropical regions of Asia, with China and India being important centers of diversity. Although native to tropical Asia, *Pogostemon* is cultivated in several countries, while commercial essential oil production is primarily carried out in India, Brazil, Taiwan, and the Seychelles (Figure 2) [7].



Figure 2. Phytoconstituents from *Pogostemon patchouli*.

PHYTOCHEMICAL STUDIES

Pogostemon benghalensis is an aromatic member of the Lamiaceae family that has attracted considerable attention because of its rich phytochemical composition. Its essential oil is dominated by sesquiterpenes, along with smaller amounts of monoterpenes and diterpenes, which contribute to its antimicrobial, antioxidant, anti-inflammatory, and other pharmacological properties. The plant also contains flavonoids, alkaloids, and chalcone derivatives with reported cytotoxic and antimutagenic activities. Owing to its pleasant aroma, the essential oil is widely used in perfumes, cosmetics, aromatherapy, and traditional medicine. Although cultivated for its economic value, the species has become invasive in some regions due to its rapid growth, efficient vegetative propagation, and ability to adapt to diverse environmental conditions (Table 2) [8].

Figures 3–7 depict the molecular structures of major bioactive phytochemicals present in Patchouli essential oil. These compounds are responsible for the plant's characteristic aroma and several important biological activities, including antioxidant, antimicrobial, and anti-inflammatory effects. Their structural diversity highlights the therapeutic and industrial significance of *Pogostemon cablin*.

TAXONOMICAL CLASSIFICATION

Pogostemon benghalensis (patchouli) is a flowering, vascular plant of the family Lamiaceae (mint family) and the order Lamiales. It is an aromatic dicot widely recognized for its essential oil and medicinal importance. Morphologically, it is an erect shrub or undershrub with square, hairy stems and

opposite, broad ovate leaves having serrated margins. The plant bears fragrant pink to purple or whitish flowers arranged in terminal spikes (verticillasters), while its well-developed fibrous roots enable adaptation to diverse soil conditions. The fruit consists of four small nutlets, and the leaves contain secretory structures responsible for essential oil production [2, 9].

Table 2. Major phytochemical compounds identified in the essential oil of *Pogostemon cablin* (Patchouli), including their molecular formula, molecular class, and structural skeleton/type.

S.N.	Phytochemical compound	Molecular formula	Molecular class	Structural skeleton / type
1	Patchoulol (Patchouli alcohol)	C ₁₅ H ₂₆ O	Oxygenated sesquiterpene	Patchoulane tricyclic alcohol.
2	α -Bulnesene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Bulnane skeleton.
3	α -Guaiene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Guaiane bicyclic.
4	Seychellene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Guaiane derivative.
5	Pogostol	C ₁₅ H ₂₆ O	Oxygenated sesquiterpene	Patchoulane alcohol.
6	Norpatchoulol	C ₁₄ H ₂₄ O	Oxygenated sesquiterpene	Patchoulane derivative
7	β -Patchoulene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Patchoulene skeleton.
8	α -Patchoulene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Patchoulene skeleton.
9	γ -Patchoulene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Patchoulene skeleton.
10	β -Caryophyllene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Caryophyllane bicyclic.
11	Caryophyllene oxide	C ₁₅ H ₂₄ O	Oxygenated sesquiterpene	Epoxidized caryophyllene.
12	δ -Guaiene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Guaiane skeleton.
13	β -Guaiene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Guaiane skeleton.
14	α -Humulene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Humulane monocyclic.
15	Germacrene D	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Germacrane skeleton.
16	Germacrene B	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Germacrane skeleton.
17	β -Elemene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Elemene skeleton.
18	α -Selinene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Selinane skeleton.
19	β -Selinene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Selinane skeleton.
20	Viridiflorene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Eremophilane derivative.
21	Ledene	C ₁₅ H ₂₄	Sesquiterpene hydrocarbon	Ledane skeleton.
22	Cubebol	C ₁₅ H ₂₆ O	Oxygenated sesquiterpene	Cubebane alcohol.
23	Nerolidol	C ₁₅ H ₂₆ O	Sesquiterpene alcohol	Acyclic sesquiterpene.
24	Farnesol	C ₁₅ H ₂₆ O	Sesquiterpene alcohol	Acyclic terpene alcohol.
25	Spathulenol	C ₁₅ H ₂₄ O	Oxygenated sesquiterpene	Aromadendrane skeleton.
26	Globulol	C ₁₅ H ₂₆ O	Oxygenated sesquiterpene	Eudesmane alcohol.
27	Eudesmol (α -Eudesmol)	C ₁₅ H ₂₆ O	Oxygenated sesquiterpene	Eudesmane alcohol.
28	β -Eudesmol	C ₁₅ H ₂₆ O	Oxygenated sesquiterpene	Eudesmane alcohol.
29	Patchouli epoxide	C ₁₅ H ₂₄ O	Oxygenated sesquiterpene	Epoxidized patchoulene.
30	Azulene derivatives	~C ₁₅ H ₁₈	Aromatic sesquiterpene	Azulene bicyclic.

Cultivation

Pogostemon benghalensis is mainly cultivated for its essential oil, which is obtained from the leaves. The plant grows best in warm, humid climates with fertile, well-drained soil. It is commonly propagated through stem cuttings because they establish more quickly and uniformly than seeds. Appropriate irrigation, regular pruning, and effective pest management are important for maintaining healthy growth and maximizing essential oil yield [10].

CHEMICAL COMPOSITION

The essential oil derived from patchouli leaves is the most valuable component of the plant. It comprises numerous chemical constituents such as patchouli alcohol, α -bulnesene, β -patchoulene, and seychellene. Patchouli alcohol, in particular, is the main component responsible for the characteristic

aroma and plays a vital role in its various applications. Vernacular names of *Pogostemon benghalenese* are given in Table 3 [11].




Figure 3. Norpatchoulenol.




Figure 4. β -Patchoulene.






Figure 5. B-Caryophyllene.

**Figure 6.** Patchoulol.**Figure 7.** Eudesmol (α -Eudesmol).**Table 3.** Vernacular names of *Pogostemon benghalensis*.

Category	Details
Scientific Name	<i>Pogostemon benghalensis</i> (= <i>Pogostemon frutescens</i> J. Garham, <i>Pogostemon indicus</i> (Roth) Kuntze, <i>Pogostemon purpuricaulis</i> Dalzell., <i>Origanum benghalense</i> Burm.f., <i>Origanum indicum</i> Roth, <i>Mentha integra</i> Buch.-Ham. Ex Benth.).
Family	Lamiaceae.
Order	Lamiales.
Common Names	Bengali <i>Pogostemon</i> , Kukesene Patchouli.
Vernacular Names	
Kala Basing	
Lujrya	
Ishwar jata	Hindi.
Jui-lata	Bengali.
Sukloti	Assamese.
Lamgi thoiding	
Pangli	Marathi.
Dumobadotoko	Oriya.
Poksunga	Oriya.
Bhoothachedayan	Malayalam.
Arissikai	Tamil.
Naati Pachhe thene	Kannada.
Pedda Tulasi	Telugu.
Gondripula	Telugu.
Kasurijang	Telugu.
Niam ja ngu	
Rguutili (Dwarpaani)	
Kali-bant	Tharu (Nepalese).
Kali Suhali	Pakistan.

- *Leaves*: Fresh leaves are used as poultices to clean wounds and promote healing. The essential oil extracted from the leaves exhibits notable antifungal activity and is also used as a styptic to control minor bleeding. The leaves and their juice have traditionally been used to manage fever in various indigenous medicinal systems. Additionally, leaf juice mixed with dried gum of *Sterculia urens* and fried in sesame (til) oil until it turns black is traditionally consumed on an empty stomach for the treatment of piles [12].
- Shoots are used as Blackish blue Dye [9].
- *Stems*: Ashes of stems are reported to be used as manure for paddy crops [10].
- *Roots*: Root used in Hemorrhage especially in uterine hemorrhage [13].

Traditional Ethnopharmacological Uses

The essential oil of *Pogostemon benghalensis* is obtained mainly from dried leaves by steam distillation. It contains sesquiterpenes, with patchoulol being the major constituent responsible for its characteristic earthy aroma and long-lasting fragrance. Because of its excellent fixative properties, the oil is widely used in perfumes, cosmetics, soaps, hair care products, and aromatherapy formulations.

Traditionally, the plant has been used to manage respiratory disorders, digestive complaints, skin diseases, wounds, fever, and insect bites. Modern studies also suggest that its essential oil possesses antimicrobial, anti-inflammatory, antifungal, antioxidant, and wound-healing properties, supporting many of its traditional applications [14].

Besides its medicinal importance, *P. benghalensis* has economic value as a source of essential oil, natural dyes, and aromatic products. In some regions, however, it behaves as an invasive species because of its rapid growth and adaptability, making proper cultivation and management necessary [15].

NUTRITIVE VALUE

Nutritional analysis of *Pogostemon benghalensis* leaves indicates that they contain carbohydrates, proteins, dietary fibre, fatty acids, tannins, and high moisture content. Fatty acid profiling has identified palmitic acid along with essential unsaturated fatty acids, including oleic, linoleic, and linolenic acids, which contribute to the plant's nutritional and potential health benefits [16].

Ethnobotanical Uses

Different parts of *Pogostemon benghalensis*, particularly the leaves and roots, have long been used in traditional medicine by indigenous communities across Asia. The plant is prepared as decoctions, fresh extracts, infusions, or poultices to treat digestive disorders, fever, skin ailments, and various common illnesses. It is also valued in aromatherapy for its calming, antiseptic, and skin-protective properties, reflecting its importance in traditional healthcare practices [17].

- *Pharmaceutical Activity*: Various medicinal properties have been studied with the *P. benghalensis* plant. Some of the activities of the plant – such as antibacterial, antiviral, antioxidant, anti-fungal, and anti-inflammatory – are discussed below.
- *Antibacterial Activity*: Extracts and essential oils of *Pogostemon benghalensis* have demonstrated notable antibacterial activity against both Gram-positive and Gram-negative bacteria. Studies have shown inhibitory effects against pathogens such as *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Salmonella typhi*. The essential oil has exhibited particularly strong antibacterial activity, supporting the traditional use of the plant in managing infections, wounds, dysentery, and other bacterial diseases. Differences in antimicrobial efficacy among studies are likely related to variations in plant source, extraction methods, and chemical composition [18].
- *Antivirus Effect*: *Pogostemon benghalensis* has traditionally been used to treat colds, coughs, and dysentery, suggesting possible antiviral properties. Experimental studies indicate that methanolic extracts of the aerial parts exhibit selective antiviral activity, showing inhibitory effects against Sindbis virus, while limited or no significant activity has been reported against certain other

mammalian viruses. These findings suggest that the plant may possess antiviral potential, although further studies are needed to confirm its clinical significance [19].

- **Antifungal Activity:** *Pogostemon benghalensis* has been traditionally used to treat fungal skin infections such as scabies and ringworm. Scientific studies have confirmed its antifungal potential, with methanolic extracts and essential oil showing activity against several pathogenic fungi, including *Microsporum gypseum*, *Trichophyton mentagrophytes*, *Fusarium* species, and *Candida albicans*. These findings support its traditional use as a natural antifungal agent and highlight its potential for developing plant-based antifungal therapies [20].
- **Antioxidant Activity:** Some study evaluated the antioxidant activity of the crude essential oil of ten species of *Lamiaceae*, including *P. benghalensis*, using the 1,1-diphenyl-2-picrylhydrazyl radical (DPPH) scavenging method. activity and total antioxidant activity (TAA). *P. benghalensis* essential oil showed scavenging activity for DPPH and TAA with an EC₅₀ value (half maximum effective concentration) of $171.3 \pm 5.74 \mu\text{g/mL}$, $89.5 \pm 2.33 \mu\text{g/mL}$, where the corresponding values of DPPH, pH standard butylhydroxytoluene (BHT) and standard reducing acid of ascorbic acid for TAA were 156.4 and 165.7 $\mu\text{g/mL}$. The EC₅₀ value of pogostemon oil was almost close to the standard value for DPPH scavenging activity, and the EC₅₀ value of TAA was lower than the standard value [6, 10].
- **Larvicidal Activity:** Some Researchers evaluated the larvicidal potential of four types of patchouli essential oils, including *P. benghalensis*, against four-stage larvae of *Aedes albopictus*. *P. benghalensis* essential oil showed 100% larval mortality at concentrations of 100 and 200 ppm as a function of the sesquiterpene-rich essential oil. Some reports suggest larvicidal effects [12].
- **Anti-Tumor Effect:** Researcher reported the anti-Tumor effect of *P. benghalensis* and reported that the MST (median survival time) of mouse tumors when treated with HEEP (hydro ethanol extract) was significant. AEP (aqueous extract) and 5-FU (5 fluorouracil) for tumor control. At one month, mice treated with 5-FU (20 mg/kg), HEEP (500 mg/kg), and nAEP showed a tumor volume of $\pm 1.90 \text{ mL}$, whereas mice treated with 5FU (20 mg/kg), HEEP. The number of rats treated with (500 mg/kg) was 1.67 mL and 1.62 mL (500 mg/kg). Compared with tumor control mice with tumor volume $\leq 3 \text{ mL}$.

CONCLUSION

Pogostemon benghalensis is a medicinally important aromatic plant with a rich phytochemical profile, including sesquiterpenes, flavonoids, phenolics, tannins, steroids, and essential oils. Its chemical composition varies with geographical location, climate, plant maturity, and extraction methods. Numerous experimental studies have demonstrated antibacterial, antifungal, antiviral, antioxidant, anti-inflammatory, anticancer, and insecticidal activities of its extracts and essential oil, supporting its traditional medicinal use. Despite these promising findings, evidence from animal studies and clinical trials remains limited. Future research should focus on isolating bioactive compounds, understanding their mechanisms of action, and conducting well-designed preclinical and clinical studies to validate their therapeutic potential. Sustainable cultivation and efficient utilization of *P. benghalensis* could further enhance its value in the pharmaceutical, cosmetic, agricultural, and nutraceutical industries.

REFERENCES

1. Bhuiyan Md NI, Varshney VK, Shiam CV, Arvind T, Farhana. Composition of essential oil of the leaf and inflorescence of *Pogostemon benghalensis* (Burm.f.) Kuntze. *Int Res J Plant Sci*. 2011;2(9):271–5.
2. Barker DH, Mitra CR. The Indian Chemical Industry-Its Development and Needs. In: *Advances in Chemical Engineering*. 1981 Jan 1;11:135–197. Academic Press.
3. *Pogostemon benghalensis* - Bengal Shrub-Mint [Internet]. *Flowersofindia.net*. 2026 [cited 2026 Sept 2]. Available from: <https://www.flowersofindia.net/catalog/slides/Bengal%20Shrub-Mint.html>
4. ST, Mahbuba K, Husain SZ. A Micromorphological study of *Pogostemon* des f. Species (Lamiaceae) from Bangladesh. *Pak J Bot*. 1995;27(1):73–82.

5. Joshi RK, Satyal P, Setzer WN. Himalayan aromatic medicinal plants: A review of their ethnopharmacology, volatile phytochemistry, and biological activities. *Medicines*. 2016 Feb 19;3(1):6.
6. Kumar B, Vijayakumar M, Govindarajan R, Pushpangadan P. Ethnopharmacological approaches to wound healing—Exploring medicinal plants of India. *J Ethnopharmacol*. 2007 Nov 1;114(2):103–13.
7. Unni BG, Borah A, Wann SB, Singh HR, Devi B, Bhattacharjee M. Phytochemical and antibacterial study of traditional medicinal plants of North East India on *Escherichia coli*. *Asian J Exp Sci*. 2009 Mar 4;23(1):103–8.
8. Padal SB, Murty PP, Rao DS, Venkaiah M. Ethnomedicinal plants from Paderu division of Visakhapatnam district, AP, India. *J Phytol*. 2010 Dec 5;2(8).
9. Babu NC, Naidu MT, Venkaiah M. Ethnomedicinal plants of Kotia hills of Vizianagaram district, Andhra Pradesh, India. *J Phytol*. 2010 Nov 4;2(6).
10. Kamble SY, Patil SR, Sawant PS, Sawant S, Pawar SG, Singh EA. Studies on plants used in traditional medicine by Bhilla tribe of Maharashtra. *Indian J Tradit Knowl*. 2010 Jul 1;9(3):591–8.
11. Nath A, Nath A, Maiti GG. Shuktani-a new ethno-medico recipe among the Sylheti Bengali community of Barak valley, Southern Assam, India. 2012:156–160.
12. Dangol DR. Traditional uses of plants of commonland habitats in Western Chitwan, Nepal. *J Inst Agric Anim Sci*. 2008;29:71.
13. Bhattarai S, Chaudhary RP, Taylor RS. Ethno-medicinal plants used by the people of Nawalparasi District, Central Nepal. *Our Nature*. 2009;7(1):82–99.
14. Sen P, Dollo M, Choudhury MD, Choudhury D. Documentation of traditional herbal knowledge of Khamptis of Arunachal Pradesh. *Indian J Tradit Knowl*. 2008 Jul 1;7(3):438–42.
15. Harvey AL. Natural products in drug discovery. *Drug Discov Today*. 2008 Oct 1;13(19–20):894–901.
16. Meena AK, Bansal P, Kumar S. Plants-herbal wealth as a potential source of ayurvedic drugs. *Asian J Tradit Med*. 2009 Aug 20;4(4):152–70.
17. Sucher NJ, Carles MC. Genome-based approaches to the authentication of medicinal plants. *Planta Med*. 2008 May;74(6):603–23.
18. Ganie SH, Upadhyay P, Das S, Sharma MP. Authentication of medicinal plants by DNA markers. *Plant Gene*. 2015 Dec 1;4:83–99.
19. Sharma A, Namdeo A, Mahadik K. Molecular markers: New prospects in plant genome analysis. *Pharmacogn Rev*. 2008;2(3):23.
20. Heubl G. New aspects of DNA-based authentication of Chinese medicinal plants by molecular biological techniques. *Planta Med*. 2010 Dec;76(17):1963–74.