

**This article is in formatting stage final PDF Will change**

**Journal:** Research & Reviews: A Journal of Biotechnology

**ISSN:** 2231-3826

**Title:** Cucumis Sativus: A Review

**Research Article**

**Volume:** 15

**Issue:** 01

**Year:** 2025

**Article Received:** 30 April 2024

**Article Accepted:** 24 February 2025

**Article Published:**

**1.Akhil Dhar, 2. Pratham Sharma, 3. Avi Singh**

1,2,3B.Tech Student, Department of Biotechnology, Meerut Institute of Engineering and Technology, Meerut-250005, Uttar Pradesh, India.

**4.Surya Prakash D.V**

4.Assistant Professor, Department of Biotechnology, Meerut Institute of Engineering and Technology, Meerut-250005, Uttar Pradesh, India.

**Email:** [prakashsurya55@gmail.com](mailto:prakashsurya55@gmail.com)

**Abstract**

The family *Cucurbitaceae* includes the cucumber (*Cucumis sativus* L.). 30 species of *Cucumis* are distributed in Africa and Asia. One of the first vegetables to be farmed, cucumbers are native to the tropics. It has almost 3,000 years of historical documentation (Yawalkar, 1985). Widespread

cultivation of *Cucumis sativus* (CS) is found in Bangladesh (including the Chittagong Hill Tracts, Gazipur, Sherpur, Cox's Bazar, Tangail, and other forests), the Indian subcontinent, and Sri Lanka. Cucumbers are said to have originated in the region that is now mostly occupied by Bangladesh, in the foothills of the Himalayas, north of the Bay of Bengal. Its uses in Ayurveda involve a stinging feeling, enlarged eyes, burns, dermatitis, anuria, skin whitening, emollient, and moisturising. Children's dysentery is treated with raw, ripe fruit in China and India. Another powerful anthelmintic is the cucumber plant. Frontal headaches and sleeplessness can both be treated with seed oil. The main compounds in plants that carry out different functions include flavonoids, glycosides, and tannins.

Keywords: *Cucumis sativus*, Phenolics, Antioxidant, Antidiabetic.

## Introduction

As a result of this area is home to the majority of the recognised medicinal plants worldwide. The resources used to prepare medicine were plants, plant parts, and plant products; these medicinal plants and plant parts are a significant source of natural riches for a nation [1]. Considering herbal remedies have been used for a very long time in many different cultures, they are generally thought to be safe. The entire herbal market in the world is valued at 62.0 billion dollars [2]. With 45% of the global herbal market, the European Union holds the largest market share, with India contributing just \$1 billion. Being fed organically, cucumbers produce more and better because they are heavy users of organic materials. If there are essential element deficiencies in the soil, a balanced fertiliser should be applied. A minimum of 5% nitrogen and around 20% of organic elements such as dried blood, pulverised cottonseed, dehydrated manures, bone, and fish should be included in this meal. Additional disorders such as pitta, hyperdipsia, fever, sleeplessness, cephalgia, bronchitis, jaundice, haemorrhages, strangury [3], and general debility are also treated with it. Additionally, a plethora of academic publications have demonstrated the fruit extracts' anthelmintic, anti-diabetic, anti-ulcer, moisturising, and antibacterial properties at varying dosages. The pulp and seeds of cucumbers are used as a folk remedy for skin conditions like hyperpigmentation and to clean the face. In addition, *C. sativus* is still widely suggested for use in the cosmetics business today, having been used for centuries. As demonstrated by the diverse pharmacological activity and toxicities of its many sections, every plant, including the wise *Cucumis sativus*, has a number of phytoconstituents [4]. Because it contained chemicals that have

therapeutic potential, it was also exhibiting a wide range of pharmaceutical action. Cucumber fruit is mostly composed of water; roughly 96% of the edible, unpeeled fruit is water. *Cucumis sativus* is also said to include vitamins, minerals, amino acids, phytosterols, phenolic acids, fatty acids, and curcubitacin, according to one source. Cucumbers are said to include trace levels of vitamin C, amino acids, pectins, carbs, sugars, and essential oils. [5]. The following substances were found in a water-soluble extract of cucumber fruit: glycosides, triterpenoid, flavonoids, carbs, and tannins. Fruits are used as a hair tonic, to treat cirrhosis bronchitis, asthma, dyspepsia, piles, diarrhoea, coughing, hoarseness of voice, and eye diseases. They are also astringent, laxative, anthelmintic, and antipyretic. The green fruit's preparation is used to alleviate coughs. Fruit pulp assists with piles, dysenteric diarrhoea, leprosy, and dropsy. Half-ripe fruit is used as a purgative. There are drug-like kernels in the fruit. Fruits are utilised in Khagrachari medicine to cure menstrual problems. A treatment for rheumatism is seed oil [6]. Bark gum has a purgative and soothing effect. The fruits contain a triterpenoid that exhibits strong antibacterial properties. Mice that were given kernel oil for an extended period of time tolerated it well and it had purgative effects.

## Phytochemicals

**Carotene** : These phytochemicals are found in large amounts in tomatoes and pumpkins and have antioxidant qualities [7]. They give fruits and vegetables their colors—yellow, red, and orange.

**Anthocyanins**: Plants include a set of pigments that are deep red, purple, and blue in colour. They belong to a class of polyphenols that give many plants their colour.

**Polyphenols** : These are derived from the phenol molecule, which is made up of six carbon atoms bonded together in a ring configuration. They comprise the broad categories of stilbenes, phenolic acids [8].

**Phenolics** : These are examples of metabolites called secondary metabolites which develop after the breakdown of phenylpropanoid in pentose phosphate and the shikimic acid of plants. The vast majority of plant tissues [9], particularly those of fruits and vegetables like thyme and rosemary, contain them.

## Uses

This vegetable has extremely few calories but high water content. Strong antioxidant, anti-diabetic, and antilipidemic properties are seen in it. It is frequently added to fermented pickles or salads. Antioxidant chemicals that are consumed through food play a major role in protecting human health, and the Mediterranean diet, which is high in natural antioxidants, is well known to prevent cancer and cardiovascular disease [10]. Fruit contains extremely little proteins (0.6%), lipids (0.1%), and carbs (2.2%), and its main components are made up of water (95%) and fats. Different types of *C. sativus* seedlings were found to have cotyledons containing Cucurbitacins A, B, C, D, E, and I. These are the cause of the bitter taste in fruits. This fruit hydrolyze to a non-bitter state upon maturity. Cucumbers seem to have different levels of bitterness from year to year and from place to place. This could be because specific environmental conditions, such cool temperatures, can increase or decrease the formation of elaterase, which in turn can increase the bitterness of fruit [11]. As stated in, another study reveals that cucurbitacin C, which gives cucumbers their bitter taste, is present in very small amounts (0.001 mg). It is traditional to cut the fruit's stem end off before using it since the stem contains more cucurbitacin C than the fruit's core or bloom end.

## Pharmacological Properties

### Anti Cancer

The anticancer potential of *C. sativus* fruit methanol and acetone extracts was assessed. The findings showed that the fruit of *C. sativus* exhibited a high concentration of bioactive chemicals with anticancer activity when tested on IC50 cell lines (MCF 714.5 ± 1.3 and HeLa 26.1 ± 1). *C. sativus* cucurbitacins inhibited cell proliferation [12], prevented migration and invasion, promoted apoptosis, and promoted cell cycle arrest, among other methods of action, to operate as an anticancer agent. The Wnt, PI3K/Akt, JAK-STAT3, and MAPK pathways were among the signalling pathways that curcumins also blocked. The survival and apoptosis of cancer cells are significantly influenced by those signalling pathways. Combining cucurbitacins with chemotherapeutic medications (such methotrexate and docetaxel) can have synergistic anticancer effects.

### Anti oxidant

The antioxidant activity of *C. sativus* fruit was assessed using the superoxide radical scavenging assay and the 2,2-diphenyl-1-picrylhydrazyl (DPPH) method, which employed butylated

hydroxytoluene as a reference. In comparison to butylated hydroxytoluene [13], which had IC<sub>50</sub> values of  $31.37 \pm 1.42$  and  $51.78 \pm 1.04$   $\mu\text{g/ml}$ , Significant DPPH free radical and superoxide radical scavenging activity was shown by *C. sativus* fruit extract, with IP values consisting  $14.71 \pm 1.41$  and  $35.29 \pm 1.30$   $\mu\text{g/ml}$ . In the fruit, nonphenolic substances such uracil and methylenecycloartenol shown strong antioxidant activity.

### **Anti Diabetic**

Male mice were given an ethanolic extract of *C. sativus* peels at doses of 250 and 500 mg/kg for 15 days to observe changes in hepatic lipid peroxidation (LPO) and blood glucose levels. The study's findings indicated that an ethanol extract of *C. sativus* peels might control the changes in the thyroid hormone and blood lipid levels that were caused by alloxan-induced diabetes mellitus. [14]. The 500 mg/kg dose was tried for improved effects. Rats with streptozotocin-induced diabetes were used to test the antidiabetic potential of an aqueous extract of *Cucumis trigonus* fruit. The outcome showed statistical data that showed a significant drop in blood glucose, serum triglycerides, total cholesterol, and glycosylated haemoglobin levels and a rise in serum insulin level.

### **Anti Inflammatory**

Two dosages (150 and 250 mg/kg bw) of the methanolic extract of *C. sativus* leaves were found to exhibit anti-inflammatory action in the Long Evans rat model. The standard indomethacin (10 mg/kg bw) was used to compare the effects [17]. The outcome demonstrated that, in comparison to the conventional medication (78.41%), Studies conducted on endothelial cells have demonstrated that *C. sativus* extract can reduce the inflammatory response triggered by lipopolysaccharide [15]. Additionally, the aqueous fraction of *C. sativus* may lessen inflammatory factors caused by Angiotensin II. By preventing sialidase from producing functionally active HA sticky CD44, iminosugar idoBR1, an iminosugar amino acid derived from *C. sativus* fruit, may have anti-inflammatory properties.

### **Anti Bacterial**

*Cucumis sativus* potency as an antibacterial against four microbiological illnesses in humans. The diffusion technique using agar wells was employed to conduct the antibacterial assay [18]. At a

particular dose of seed extract, the highest zone of inhibition against *S. aureus* was observed. Each and every one of these pathogens, safeguard for *P. aeruginosa*, has been identified and *E. coli*, have been identified [16], were more susceptible to the extract of methanol. In final analysis, they came to the conclusion the potential for broad-spectrum antibacterial activity was present in CS seeds.

## Conclusion

The study indicated that the *Cucumis sativus* has the potential to be an excellent substitute for the present generation of synthetic anti-inflammatory medicines. It may be used as an adjunct to artificial anti-inflammatory medications, with the potential to eliminate side effects, reduce total costs, and mitigate drug resistance. It acts as antioxidants. So it shows high level of pharmacological properties for control of disorders. *Cucumis sativus* shown potent actions over a broad range of parameters, indicating the presence of molecules with a wide range of activity.

## References

1. Shah, Pritesh, et al. A review on *Cucumis sativus* (Cucumber). *Research Journal of Pharmacognosy and Phytochemistry*, 2013, 5(2): 49-53.
2. Sahu, Tripti, and Jyoti Sahu. *Cucumis sativus* (cucumber): A review on its pharmacological activity. *Journal of Applied Pharmaceutical Research*, 2015, 3(1): 04-09.
3. Kumari K, Topno SE. Performance of Different F1 Hybrids of Cucumber (*Cucumis sativus* L.) on Growth, Yield and Quality. *Int. J. Plant Soil Sci.* 2023, 35 (22):690-96.
4. Mallik, Jony, Priyanka Das and Sourav Das. Pharmacological activity of *Cucumis sativus* L.–a complete overview. *Asian Journal of Pharmaceutical Research and Development*, 2013: 1-6.
5. Kaur, Manpreet, and Parveen Sharma. Recent advances in cucumber (*Cucumis sativus* L.). *The Journal of Horticultural Science and Biotechnology*, 2022, 97(1): 3-23.
6. Khan, Anayatullah, et al. Biological and medicinal application of *Cucumis sativus* Linn.–review of current status with future possibilities. *Journal of Complementary and Integrative Medicine*, 2022, 19(4): 843-854.
7. Feng, Shengjun, et al. Recent progress on the molecular breeding of *Cucumis sativus* L. in China. *Theoretical and Applied Genetics*, 2020, 133: 1777-1790.

8. Zhang, Juping, et al. The formation of fruit quality in *Cucumis sativus* L. *Frontiers in Plant Science*, 2021, 12: 729-48.
9. Burza, Wojciech, et al. Cucumber (*Cucumis sativus* L.). *Agrobacterium protocols*, 2006: 427-438.
10. Weng, Yiqun. *Cucumis sativus* chromosome evolution, domestication, and genetic diversity: Implications for cucumber breeding. *Plant Breeding Reviews*, 2021, 44: 79-111.
11. K.Patil, A.Kandhare, D.Bhise. Pharmacological evaluation of ameliorative effect of aqueous extracts of *Cucumis sativus* L.fruit formation on wound healing in Wister rats. *Chronicles of young scientists*, 2012, 2(4):207- 213.
12. S. Sharma., J. Dwivedi., S.Paliwal. Evaluation of antacid and carminative properties of *Cucumis sativus* under simulated conditions. *Scholars Research Library Der Pharmacia Lettre*, 2012, 4(1): 234-239.
13. Z. Khan, A. H. Shah, R. Gul, A. Majid, U. Khan, H. Ahmad. Morpho-agronomic characterization of cucumber germplasm for yield and yield associated traits. *International Journal of Agronomy and Agricultural Research*, 2015, 6(1):1-6.
14. N.S.Gill and M.Bali. Isolation of antiulcer Cucurbitane type triterpenoid from the seeds of *Cucurbita pepo*. *Research Journal of Phytochemistry*, 2011, 5(2): 70-79.
15. JA.Sood, P. Kaur and R. Gupta. Phytochemical Screening and Antimicrobial Assay of Various Seeds Extract of *Cucurbitaceae* Family. *International Journal of Applied Biology and Pharmaceutical Technology*, 2012, 3(3): 401-409.
16. H. Saeed. Traditional and Medicinal Uses of Plants of Duniyapur District Lodhran. *International Journal of Advance Research*, 2017, 5(1): 1823-1827.
17. Harshitha CK, Shyamamma S. Morphological characterization of local cucumber (*Cucumis sativus*) Genotypes for fruit quality traits. *Mysore Journal of Agricultural Sciences*, 2021, 55(4):130-141.
18. Ene CO, Ogbonna PE, Agbo CU, Chukwudi UP. Heterosis and combining ability in cucumber (*Cucumis sativus*). *Information Processing in Agriculture*. 2019, 6(1):150-7.

**This article is in formatting stage final PDF Will change**